



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 01:03 PM UTC

PDB ID : 2RBL / pdb_00002rbl
Title : High resolution design of a protein loop
Authors : Hu, X.; Wang, H.; Ke, H.; Kuhlman, B.
Deposited on : 2007-09-19
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	
1	B	104	
1	M	104	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2009 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tenascin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	89	693	428	110	153	2	0	0	0
1	B	89	693	428	110	153	2	0	0	0
1	M	80	623	386	99	137	1	0	0	0

There are 57 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	801	MET	-	initiating methionine	UNP P24821
A	824	SER	PHE	engineered mutation	UNP P24821
A	825	MET	LYS	engineered mutation	UNP P24821
A	826	GLN	PRO	engineered mutation	UNP P24821
A	828	SER	ALA	engineered mutation	UNP P24821
A	829	GLN	GLU	engineered mutation	UNP P24821
A	830	LEU	ILE	engineered mutation	UNP P24821
A	831	GLU	ASP	engineered mutation	UNP P24821
A	894	ALA	-	expression tag	UNP P24821
A	895	ALA	-	expression tag	UNP P24821
A	896	ALA	-	expression tag	UNP P24821
A	897	LEU	-	expression tag	UNP P24821
A	898	GLU	-	expression tag	UNP P24821
A	899	HIS	-	expression tag	UNP P24821
A	900	HIS	-	expression tag	UNP P24821
A	901	HIS	-	expression tag	UNP P24821
A	902	HIS	-	expression tag	UNP P24821
A	903	HIS	-	expression tag	UNP P24821
A	904	HIS	-	expression tag	UNP P24821
B	801	MET	-	initiating methionine	UNP P24821
B	824	SER	PHE	engineered mutation	UNP P24821
B	825	MET	LYS	engineered mutation	UNP P24821
B	826	GLN	PRO	engineered mutation	UNP P24821

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	828	SER	ALA	engineered mutation	UNP P24821
B	829	GLN	GLU	engineered mutation	UNP P24821
B	830	LEU	ILE	engineered mutation	UNP P24821
B	831	GLU	ASP	engineered mutation	UNP P24821
B	894	ALA	-	expression tag	UNP P24821
B	895	ALA	-	expression tag	UNP P24821
B	896	ALA	-	expression tag	UNP P24821
B	897	LEU	-	expression tag	UNP P24821
B	898	GLU	-	expression tag	UNP P24821
B	899	HIS	-	expression tag	UNP P24821
B	900	HIS	-	expression tag	UNP P24821
B	901	HIS	-	expression tag	UNP P24821
B	902	HIS	-	expression tag	UNP P24821
B	903	HIS	-	expression tag	UNP P24821
B	904	HIS	-	expression tag	UNP P24821
M	801	MET	-	initiating methionine	UNP P24821
M	824	SER	PHE	engineered mutation	UNP P24821
M	825	MET	LYS	engineered mutation	UNP P24821
M	826	GLN	PRO	engineered mutation	UNP P24821
M	828	SER	ALA	engineered mutation	UNP P24821
M	829	GLN	GLU	engineered mutation	UNP P24821
M	830	LEU	ILE	engineered mutation	UNP P24821
M	831	GLU	ASP	engineered mutation	UNP P24821
M	894	ALA	-	expression tag	UNP P24821
M	895	ALA	-	expression tag	UNP P24821
M	896	ALA	-	expression tag	UNP P24821
M	897	LEU	-	expression tag	UNP P24821
M	898	GLU	-	expression tag	UNP P24821
M	899	HIS	-	expression tag	UNP P24821
M	900	HIS	-	expression tag	UNP P24821
M	901	HIS	-	expression tag	UNP P24821
M	902	HIS	-	expression tag	UNP P24821
M	903	HIS	-	expression tag	UNP P24821
M	904	HIS	-	expression tag	UNP P24821


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tenascin



- Molecule 1: Tenascin



- Molecule 1: Tenascin



4 Data and refinement statistics i

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	137.20Å 137.20Å 86.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.10 50.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.10) 85.8 (50.00-2.10)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.250 , 0.300 0.307 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	46.9	Xtriage
Anisotropy	0.435	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.005 for $-1/3^*h+1/3^*k+4/3^*l,-k,2/3^*h+1/3^*k+1/3^*l$ 0.024 for $-2/3^*h-1/3^*k-4/3^*l,-1/3^*h-2/3^*k+4/3^*l,-1/3^*h+1/3^*k+1/3^*l$ 0.017 for $-h,1/3^*h-1/3^*k-4/3^*l,-1/3^*h-2/3^*k+1/3^*l$	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2009	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	2.36	33/702 (4.7%)	2.01	21/955 (2.2%)
1	B	2.38	28/702 (4.0%)	2.04	31/955 (3.2%)
1	M	2.44	25/630 (4.0%)	1.83	20/854 (2.3%)
All	All	2.39	86/2034 (4.2%)	1.97	72/2764 (2.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	M	0	1
All	All	0	3

The worst 5 of 86 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	M	813	ASP	CG-OD2	22.08	1.67	1.25
1	M	813	ASP	CG-OD1	16.35	1.56	1.25
1	B	839	ILE	C-O	-12.32	1.11	1.24
1	M	856	ASN	CG-OD1	12.32	1.47	1.23
1	A	874	ILE	N-CA	9.28	1.56	1.46

The worst 5 of 72 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	842	VAL	N-CA-C	10.70	119.44	108.95
1	M	844	GLY	N-CA-C	10.04	126.62	113.27
1	A	885	ALA	N-CA-C	-9.16	95.02	109.50
1	A	864	LYS	CA-C-N	8.65	130.65	119.84
1	A	864	LYS	C-N-CA	8.65	130.65	119.84

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	885	ALA	Peptide
1	B	858	TYR	Mainchain
1	M	856	ASN	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	693	0	671	44	0
1	B	693	0	671	40	0
1	M	623	0	599	55	0
All	All	2009	0	1941	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 119 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:813:ASP:CG	1:M:813:ASP:OD2	1.67	1.38
1:B:829:GLN:HG3	1:B:877:ARG:NH2	1.55	1.22
1:A:831:GLU:HB3	1:B:829:GLN:NE2	1.56	1.20
1:B:829:GLN:CG	1:B:877:ARG:HH21	1.58	1.15
1:A:831:GLU:HB3	1:B:829:GLN:HE21	0.93	1.07

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	87/104 (84%)	77 (88%)	6 (7%)	4 (5%)	2	0
1	B	87/104 (84%)	77 (88%)	6 (7%)	4 (5%)	2	0
1	M	74/104 (71%)	59 (80%)	10 (14%)	5 (7%)	1	0
All	All	248/312 (80%)	213 (86%)	22 (9%)	13 (5%)	1	0

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	809	ILE
1	A	828	SER
1	A	846	ARG
1	M	808	GLN
1	M	809	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/92 (88%)	67 (83%)	14 (17%)	2	1
1	B	81/92 (88%)	69 (85%)	12 (15%)	3	1
1	M	72/92 (78%)	51 (71%)	21 (29%)	0	0
All	All	234/276 (85%)	187 (80%)	47 (20%)	1	0

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	811	VAL
1	M	824	SER
1	M	812	LYS
1	M	818	THR
1	M	841	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	829	GLN
1	A	856	ASN
1	B	826	GLN
1	B	829	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	89/104 (85%)	1.45	20 (22%) 2 2	40, 54, 70, 74	0
1	B	89/104 (85%)	1.77	30 (33%) 1 1	45, 55, 69, 77	0
1	M	80/104 (76%)	1.82	28 (35%) 1 1	42, 53, 62, 69	0
All	All	258/312 (82%)	1.68	78 (30%) 1 1	40, 54, 69, 77	0

The worst 5 of 78 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	803	LEU	7.1
1	B	818	THR	4.5
1	M	803	LEU	4.4
1	M	814	VAL	4.2
1	B	827	LEU	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.