



wwPDB X-ray Structure Validation Summary Report

Mar 8, 2026 – 07:32 AM UTC

PDB ID : 4RB4 / pdb_00004rb4
Title : Crystal structure of dodecameric iron-containing heptosyltransferase TibC in complex with ADP-D-beta-D-heptose at 3.9 angstrom resolution
Authors : Yao, Q.; Lu, Q.; Shao, F.
Deposited on : 2014-09-12
Resolution : 3.88 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1186 (4.06-3.70)
Clashscore	190562	1012 (4.04-3.72)
Ramachandran outliers	187476	1167 (4.06-3.70)
Sidechain outliers	187428	1160 (4.06-3.70)
RSRZ outliers	180081	1185 (4.06-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	 4% 71% 19% • 6%
1	B	406	 4% 73% 16% • • 6%
1	C	406	 4% 72% 19% • •
1	D	406	 4% 69% 20% • 6%
1	E	406	 5% 67% 25% • •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
1	F	406	3%	72%	19%	• 5%
1	G	406	6%	68%	22%	5% • •
1	H	406	5%	71%	19%	5% 5%
1	I	406	4%	71%	20%	• •
1	J	406	5%	71%	20%	• • 5%
1	K	406	5%	72%	21%	• •
1	L	406	12%	61%	23%	7% • 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FE	A	501	-	-	X	-
2	FE	D	501	-	-	X	-
2	FE	E	501	-	-	-	X
2	FE	G	501	-	-	-	X
2	FE	J	501	-	-	X	-
2	FE	K	501	-	-	-	X
3	AQH	D	502	-	-	X	-
3	AQH	F	502	-	-	X	-
3	AQH	H	502	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 37611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyltransferase tibC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	K	390	3132	2016	546	557	13	0	0	0
1	L	374	3008	1938	524	533	13	0	0	0
1	C	389	3121	2007	545	556	13	0	0	0
1	D	381	3061	1970	536	542	13	0	0	0
1	E	390	3129	2013	546	557	13	0	0	0
1	H	385	3093	1990	541	549	13	0	0	0
1	B	383	3082	1987	538	544	13	0	0	0
1	F	384	3084	1985	539	547	13	0	0	0
1	G	389	3121	2007	545	556	13	0	0	0
1	I	389	3121	2007	545	556	13	0	0	0
1	J	385	3097	1996	541	547	13	0	0	0
1	A	380	3058	1969	535	541	13	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
K	110	ALA	ASP	engineered mutation	UNP H5MH13
L	110	ALA	ASP	engineered mutation	UNP H5MH13
C	110	ALA	ASP	engineered mutation	UNP H5MH13
D	110	ALA	ASP	engineered mutation	UNP H5MH13
E	110	ALA	ASP	engineered mutation	UNP H5MH13

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	110	ALA	ASP	engineered mutation	UNP H5MH13
B	110	ALA	ASP	engineered mutation	UNP H5MH13
F	110	ALA	ASP	engineered mutation	UNP H5MH13
G	110	ALA	ASP	engineered mutation	UNP H5MH13
I	110	ALA	ASP	engineered mutation	UNP H5MH13
J	110	ALA	ASP	engineered mutation	UNP H5MH13
A	110	ALA	ASP	engineered mutation	UNP H5MH13

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	K	1	Total Fe 1 1	0	0
2	L	1	Total Fe 1 1	0	0
2	C	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0
2	E	1	Total Fe 1 1	0	0
2	H	1	Total Fe 1 1	0	0
2	B	1	Total Fe 1 1	0	0
2	F	1	Total Fe 1 1	0	0
2	G	1	Total Fe 1 1	0	0
2	I	1	Total Fe 1 1	0	0
2	J	1	Total Fe 1 1	0	0
2	A	1	Total Fe 1 1	0	0

- Molecule 3 is [(2R,3S,4R,5R)-5-(6-amino-9H-purin-9-yl)-3,4-dihydroxytetrahydrofuran-2-yl] methyl (2R,3R,4R,5R,6S)-6-[(1R)-1,2-dihydroxyethyl]-3,4,5-trihydroxytetrahydro-2H-pyran-2-yl dihydrogen diphosphate (CCD ID: AQH) (formula: C₁₇H₂₇N₅O₁₆P₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 4 2 2	0	0
4	I	1	Total C O 4 2 2	0	0
4	J	1	Total C O 4 2 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Glycosyltransferase tibC




- Molecule 1: Glycosyltransferase tibC




- Molecule 1: Glycosyltransferase tibC




• Molecule 1: Glycosyltransferase tibC




• Molecule 1: Glycosyltransferase tibC




• Molecule 1: Glycosyltransferase tibC



• Molecule 1: Glycosyltransferase tibC




• Molecule 1: Glycosyltransferase tibC




• Molecule 1: Glycosyltransferase tibC







- Molecule 1: Glycosyltransferase tibC



- Molecule 1: Glycosyltransferase tibC



- Molecule 1: Glycosyltransferase tibC



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	88.29Å 313.20Å 164.69Å 90.00° 101.26° 90.00°	Depositor
Resolution (Å)	20.04 – 3.88 20.04 – 3.88	Depositor EDS
% Data completeness (in resolution range)	98.7 (20.04-3.88) 98.8 (20.04-3.88)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.74 (at 3.94Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.282 , 0.295 0.293 , 0.300	Depositor DCC
R_{free} test set	3995 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	98.2	Xtrriage
Anisotropy	0.113	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.22 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	37611	wwPDB-VP
Average B, all atoms (Å ²)	149.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AQH, FE, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	7/3154 (0.2%)	0.67	0/4298
1	B	0.88	7/3181 (0.2%)	0.65	0/4338
1	C	0.81	5/3219 (0.2%)	0.68	1/4390 (0.0%)
1	D	0.77	4/3158 (0.1%)	0.66	1/4305 (0.0%)
1	E	0.75	4/3227 (0.1%)	0.67	0/4401
1	F	0.88	8/3181 (0.3%)	0.67	1/4336 (0.0%)
1	G	0.82	7/3219 (0.2%)	0.68	0/4390
1	H	0.79	5/3191 (0.2%)	0.70	3/4351 (0.1%)
1	I	0.81	7/3219 (0.2%)	0.70	2/4390 (0.0%)
1	J	0.83	5/3196 (0.2%)	0.72	6/4358 (0.1%)
1	K	0.79	3/3231 (0.1%)	0.69	1/4406 (0.0%)
1	L	0.80	8/3102 (0.3%)	0.69	1/4225 (0.0%)
All	All	0.81	70/38278 (0.2%)	0.68	16/52188 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	3
1	D	0	1
1	E	0	3
1	F	0	4
1	G	0	2
1	H	0	2
1	I	0	1
1	J	0	2
1	K	0	2

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	2
All	All	0	26

The worst 5 of 70 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	10	ILE	CA-CB	-10.08	1.44	1.55
1	B	105	THR	CA-C	-9.58	1.41	1.52
1	A	82	GLY	CA-C	8.85	1.64	1.51
1	J	60	ASP	C-O	-8.20	1.14	1.24
1	F	30	ASP	C-O	-8.11	1.17	1.24

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	109	GLY	N-CA-C	10.11	128.81	115.36
1	K	106	GLY	N-CA-C	8.48	121.28	111.36
1	J	108	LEU	N-CA-C	7.83	119.81	111.28
1	J	105	THR	N-CA-C	-7.57	97.36	109.39
1	I	103	PHE	CA-C-N	-7.34	113.16	120.21

There are no chirality outliers.

5 of 26 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	79	PHE	Peptide
1	K	251	ARG	Sidechain
1	K	357	TRP	Peptide
1	L	79	PHE	Peptide
1	L	80	ARG	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3058	0	2956	96	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3082	0	2979	89	0
1	C	3121	0	3016	100	0
1	D	3061	0	2952	144	0
1	E	3129	0	3027	135	0
1	F	3084	0	2979	100	0
1	G	3121	0	3016	130	0
1	H	3093	0	2988	121	0
1	I	3121	0	3015	124	0
1	J	3097	0	2993	106	0
1	K	3132	0	3024	123	0
1	L	3008	0	2882	273	0
2	A	1	0	0	2	0
2	B	1	0	0	1	0
2	C	1	0	0	1	0
2	D	1	0	0	3	0
2	E	1	0	0	1	0
2	F	1	0	0	1	0
2	G	1	0	0	1	0
2	H	1	0	0	1	0
2	I	1	0	0	1	0
2	J	1	0	0	3	0
2	K	1	0	0	1	0
2	L	1	0	0	1	0
3	A	40	0	25	13	0
3	B	40	0	25	8	0
3	C	40	0	25	12	0
3	D	40	0	25	23	0
3	E	40	0	25	10	0
3	F	40	0	25	21	0
3	G	40	0	25	12	0
3	H	40	0	25	24	0
3	I	40	0	25	9	0
3	J	40	0	25	8	0
3	K	40	0	25	17	0
3	L	40	0	25	17	0
4	C	4	0	6	1	0
4	I	4	0	6	1	0
4	J	4	0	6	0	0
All	All	37611	0	36145	1551	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 1551 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:296:PHE:CZ	1:L:298:GLY:HA3	1.21	1.68
1:L:107:THR:HG22	1:L:108:LEU:CD2	1.19	1.61
1:L:296:PHE:CE2	1:L:298:GLY:HA3	1.41	1.51
1:D:110:ALA:CB	1:D:137:ILE:HD12	1.07	1.50
1:D:110:ALA:HB1	1:D:137:ILE:CG2	1.43	1.49

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/406 (92%)	360 (96%)	10 (3%)	4 (1%)	11	43
1	B	381/406 (94%)	364 (96%)	12 (3%)	5 (1%)	9	40
1	C	387/406 (95%)	367 (95%)	15 (4%)	5 (1%)	9	40
1	D	377/406 (93%)	356 (94%)	15 (4%)	6 (2%)	7	36
1	E	388/406 (96%)	366 (94%)	18 (5%)	4 (1%)	12	45
1	F	380/406 (94%)	355 (93%)	20 (5%)	5 (1%)	9	40
1	G	387/406 (95%)	365 (94%)	16 (4%)	6 (2%)	7	36
1	H	383/406 (94%)	370 (97%)	7 (2%)	6 (2%)	7	36
1	I	387/406 (95%)	364 (94%)	21 (5%)	2 (0%)	24	59
1	J	383/406 (94%)	368 (96%)	11 (3%)	4 (1%)	12	45
1	K	388/406 (96%)	377 (97%)	9 (2%)	2 (0%)	24	59
1	L	364/406 (90%)	338 (93%)	14 (4%)	12 (3%)	3	24
All	All	4579/4872 (94%)	4350 (95%)	168 (4%)	61 (1%)	9	40

5 of 61 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	10	ILE
1	L	231	TYR
1	L	233	ASN
1	L	234	ASN
1	L	299	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/355 (93%)	313 (95%)	16 (5%)	22	47
1	B	331/355 (93%)	315 (95%)	16 (5%)	23	48
1	C	335/355 (94%)	318 (95%)	17 (5%)	21	46
1	D	328/355 (92%)	313 (95%)	15 (5%)	24	48
1	E	336/355 (95%)	316 (94%)	20 (6%)	17	44
1	F	331/355 (93%)	316 (96%)	15 (4%)	24	49
1	G	335/355 (94%)	310 (92%)	25 (8%)	12	38
1	H	332/355 (94%)	313 (94%)	19 (6%)	18	45
1	I	335/355 (94%)	313 (93%)	22 (7%)	15	41
1	J	332/355 (94%)	316 (95%)	16 (5%)	23	48
1	K	336/355 (95%)	314 (94%)	22 (6%)	15	42
1	L	322/355 (91%)	297 (92%)	25 (8%)	11	36
All	All	3982/4260 (94%)	3754 (94%)	228 (6%)	18	45

5 of 228 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	320	PHE
1	A	170	TYR
1	F	270	ILE
1	A	136	ASP
1	J	80	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	337	HIS
1	G	349	ASN
1	A	337	HIS
1	G	188	HIS
1	G	367	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 12 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	AQH	I	503	-	42,43,43	2.50	15 (35%)	62,66,66	2.38	20 (32%)
3	AQH	E	502	-	42,43,43	2.50	16 (38%)	62,66,66	2.38	21 (33%)
3	AQH	C	503	-	42,43,43	2.51	16 (38%)	62,66,66	2.38	20 (32%)
3	AQH	A	502	-	42,43,43	2.51	16 (38%)	62,66,66	2.37	20 (32%)
3	AQH	D	502	-	42,43,43	2.50	16 (38%)	62,66,66	2.43	21 (33%)
3	AQH	J	503	-	42,43,43	2.52	16 (38%)	62,66,66	2.45	21 (33%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AQH	G	502	-	42,43,43	2.56	16 (38%)	62,66,66	2.49	24 (38%)
4	EDO	J	502	-	3,3,3	0.25	0	2,2,2	0.36	0
3	AQH	H	502	-	42,43,43	2.51	16 (38%)	62,66,66	2.38	20 (32%)
3	AQH	L	502	-	42,43,43	2.55	16 (38%)	62,66,66	2.48	24 (38%)
4	EDO	C	502	-	3,3,3	0.28	0	2,2,2	0.35	0
3	AQH	K	502	-	42,43,43	2.53	16 (38%)	62,66,66	2.45	21 (33%)
3	AQH	F	502	-	42,43,43	2.55	16 (38%)	62,66,66	2.49	25 (40%)
3	AQH	B	502	-	42,43,43	2.52	16 (38%)	62,66,66	2.46	22 (35%)
4	EDO	I	502	-	3,3,3	0.20	0	2,2,2	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AQH	I	503	-	-	13/27/63/63	0/4/4/4
3	AQH	E	502	-	-	14/27/63/63	0/4/4/4
3	AQH	C	503	-	-	13/27/63/63	0/4/4/4
3	AQH	A	502	-	-	10/27/63/63	0/4/4/4
3	AQH	D	502	-	-	11/27/63/63	0/4/4/4
3	AQH	J	503	-	-	9/27/63/63	0/4/4/4
3	AQH	G	502	-	-	11/27/63/63	0/4/4/4
4	EDO	J	502	-	-	0/1/1/1	-
3	AQH	H	502	-	-	10/27/63/63	0/4/4/4
3	AQH	L	502	-	-	16/27/63/63	0/4/4/4
4	EDO	C	502	-	-	1/1/1/1	-
3	AQH	K	502	-	-	15/27/63/63	0/4/4/4
3	AQH	F	502	-	-	8/27/63/63	0/4/4/4
3	AQH	B	502	-	-	7/27/63/63	0/4/4/4
4	EDO	I	502	-	-	0/1/1/1	-

The worst 5 of 191 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	K	502	AQH	PB-O2B	-9.54	1.49	1.59
3	J	503	AQH	PB-O2B	-9.49	1.49	1.59
3	G	502	AQH	PB-O2B	-9.44	1.49	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	502	AQH	PB-O2B	-9.40	1.49	1.59
3	H	502	AQH	PB-O2B	-9.33	1.49	1.59

The worst 5 of 259 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	J	503	AQH	O-C2'-C1'	-9.95	86.37	110.08
3	K	502	AQH	O-C2'-C1'	-9.91	86.45	110.08
3	L	502	AQH	O-C2'-C1'	-9.79	86.74	110.08
3	F	502	AQH	O-C2'-C1'	-9.78	86.76	110.08
3	B	502	AQH	O-C2'-C1'	-9.78	86.77	110.08

There are no chirality outliers.

5 of 138 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	K	502	AQH	C5D-O5D-PA-O1A
3	K	502	AQH	C1'-O3B-PB-O2B
3	K	502	AQH	O6'-C6'-C7'-O7'
3	K	502	AQH	C5'-C6'-C7'-O7'
3	K	502	AQH	O5'-C5'-C6'-C7'


There are no ring outliers.



14 monomers are involved in 176 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	503	AQH	9	0
3	E	502	AQH	10	0
3	C	503	AQH	12	0
3	A	502	AQH	13	0
3	D	502	AQH	23	0
3	J	503	AQH	8	0
3	G	502	AQH	12	0
3	H	502	AQH	24	0
3	L	502	AQH	17	0
4	C	502	EDO	1	0
3	K	502	AQH	17	0
3	F	502	AQH	21	0
3	B	502	AQH	8	0
4	I	502	EDO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	380/406 (93%)	0.47	16 (4%) 40 29	150, 150, 150, 150	0
1	B	383/406 (94%)	0.45	17 (4%) 39 28	150, 150, 150, 150	0
1	C	389/406 (95%)	0.67	15 (3%) 43 31	150, 150, 150, 150	0
1	D	381/406 (93%)	0.49	18 (4%) 36 27	30, 150, 150, 150	0
1	E	390/406 (96%)	0.56	20 (5%) 33 25	124, 150, 150, 150	0
1	F	384/406 (94%)	0.52	14 (3%) 46 32	150, 150, 150, 150	0
1	G	389/406 (95%)	0.57	23 (5%) 28 23	150, 150, 150, 150	0
1	H	385/406 (94%)	0.52	21 (5%) 30 24	25, 150, 150, 150	0
1	I	389/406 (95%)	0.56	16 (4%) 41 30	85, 150, 150, 150	0
1	J	385/406 (94%)	0.59	20 (5%) 33 25	150, 150, 150, 150	0
1	K	390/406 (96%)	0.59	20 (5%) 33 25	150, 150, 150, 150	0
1	L	374/406 (92%)	0.82	48 (12%) 7 11	23, 150, 150, 150	0
All	All	4619/4872 (94%)	0.57	248 (5%) 31 24	23, 150, 150, 150	0

The worst 5 of 248 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	57	CYS	11.8
1	D	110	ALA	11.4
1	E	342	CYS	5.9
1	I	30	ASP	5.7
1	E	370	CYS	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	AQH	L	502	40/40	0.66	0.23	150,150,150,150	0
4	EDO	I	502	4/4	0.70	0.33	150,150,150,150	0
4	EDO	C	502	4/4	0.71	0.38	150,150,150,150	0
2	FE	E	501	1/1	0.73	0.47	150,150,150,150	0
2	FE	K	501	1/1	0.75	0.50	150,150,150,150	0
3	AQH	I	503	40/40	0.75	0.19	150,150,150,150	0
3	AQH	A	502	40/40	0.77	0.18	150,150,150,150	40
2	FE	G	501	1/1	0.78	0.49	150,150,150,150	0
3	AQH	H	502	40/40	0.79	0.18	150,150,150,150	0
3	AQH	F	502	40/40	0.79	0.18	150,150,150,150	0
4	EDO	J	502	4/4	0.81	0.33	150,150,150,150	0
3	AQH	D	502	40/40	0.82	0.18	150,150,150,150	40
2	FE	C	501	1/1	0.82	0.43	150,150,150,150	0
3	AQH	E	502	40/40	0.83	0.15	150,150,150,150	0
3	AQH	B	502	40/40	0.84	0.15	150,150,150,150	0
3	AQH	G	502	40/40	0.86	0.15	150,150,150,150	40
3	AQH	K	502	40/40	0.87	0.13	150,150,150,150	0
3	AQH	J	503	40/40	0.87	0.14	150,150,150,150	0
3	AQH	C	503	40/40	0.87	0.14	150,150,150,150	0
2	FE	I	501	1/1	0.88	0.40	150,150,150,150	0
2	FE	J	501	1/1	0.96	0.37	150,150,150,150	0
2	FE	A	501	1/1	0.97	0.33	150,150,150,150	0
2	FE	L	501	1/1	0.97	0.35	150,150,150,150	0
2	FE	B	501	1/1	0.97	0.32	150,150,150,150	0
2	FE	F	501	1/1	0.97	0.32	150,150,150,150	0
2	FE	H	501	1/1	0.98	0.34	150,150,150,150	0
2	FE	D	501	1/1	0.98	0.32	150,150,150,150	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AQH L 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AQH I 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AQH A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AQH H 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AQH F 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AQH D 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AQH E 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AQH B 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AQH G 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AQH K 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AQH J 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around AQH C 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.