



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 01:40 AM UTC

PDB ID : 2RCR / pdb_00002rcr
Title : STRUCTURE OF THE MEMBRANE-BOUND PROTEIN PHOTOSYN-
THETIC REACTION CENTER FROM RHODOBACTER SPHAEROIDES
Authors : Chang, C.-H.; Norris, J.; Schiffer, M.
Deposited on : 1991-02-04
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	281	
2	M	307	
3	H	260	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BCL	L	350	-	-	X	-
4	BCL	M	400	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BCL	M	601	-	-	X	-
5	BPH	M	500	-	-	X	-
6	UQ	L	800	-	-	X	-
6	UQ	M	750	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7049 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOTOSYNTHETIC REACTION CENTER (L SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	278	Total	C	N	O	S	0	0	0
			2203	1491	349	355	8			


- Molecule 2 is a protein called PHOTOSYNTHETIC REACTION CENTER (M SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	305	Total	C	N	O	S	0	0	0
			2428	1620	397	400	11			

- Molecule 3 is a protein called PHOTOSYNTHETIC REACTION CENTER (H SUBUNIT).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	255	Total	C	N	O	S	73	0	0
			1927	1232	330	354	11			

- Molecule 4 is BACTERIOCHLOROPHYLL A (CCD ID: BCL) (formula: $C_{55}H_{74}MgN_4O_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	L	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
4	L	1	Total 66	C 55	Mg 1	N 4	O 6	6	0
4	M	1	Total 66	C 55	Mg 1	N 4	O 6	7	0
4	M	1	Total 66	C 55	Mg 1	N 4	O 6	3	0

- Molecule 5 is BACTERIOPHEOPHYTIN A (CCD ID: BPH) (formula: $C_{55}H_{76}N_4O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	L	1	Total	C	N	O	0	0
			65	55	4	6		
5	M	1	Total	C	N	O	7	0
			65	55	4	6		

- Molecule 6 is Coenzyme Q10, (2Z,6E,10Z,14E,18E,22E,26Z)-isomer (CCD ID: UQ) (formula: $C_{59}H_{90}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	L	1	Total	C	O	16	0
			48	44	4		
6	M	1	Total	C	O	16	0
			48	44	4		

- Molecule 7 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	M	1	Total	Fe	0	0
			1	1		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PHOTOSYNTHETIC REACTION CENTER (L SUBUNIT)

Chain L: 





4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	142.20Å 139.60Å 78.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 3.10	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-3.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.220 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7049	wwPDB-VP
Average B, all atoms (Å ²)	0.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, BPH, UQ, BCL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	L	1.18	16/2291 (0.7%)	1.90	45/3137 (1.4%)
2	M	1.49	25/2521 (1.0%)	1.96	70/3442 (2.0%)
3	H	1.14	10/1977 (0.5%)	1.80	32/2689 (1.2%)
All	All	1.29	51/6789 (0.8%)	1.89	147/9268 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	M	0	5

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	M	93	LEU	C-O	25.26	1.54	1.23
2	M	93	LEU	CA-C	-24.86	1.19	1.53
2	M	250	TRP	CA-CB	-21.86	1.14	1.53
2	M	112	LEU	C-N	14.62	1.54	1.33
2	M	93	LEU	C-N	13.10	1.57	1.33

The worst 5 of 147 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	166	ASN	CA-CB-CG	42.38	154.98	112.60
2	M	112	LEU	O-C-N	-36.86	84.11	122.07
1	L	166	ASN	N-CA-CB	19.32	142.88	111.58
2	M	109	GLU	N-CA-C	18.11	144.58	108.18
3	H	142	VAL	N-CA-CB	-14.08	96.69	111.90

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	M	109	GLU	Mainchain,Peptide
2	M	112	LEU	Mainchain,Peptide
2	M	265	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	2203	0	2156	497	1
2	M	2428	0	2342	608	1
3	H	1927	0	1927	383	2
4	L	132	0	148	39	0
4	M	132	0	148	63	0
5	L	65	0	76	16	0
5	M	65	0	76	31	0
6	L	48	0	59	22	0
6	M	48	0	63	33	0
7	M	1	0	0	0	0
All	All	7049	0	6995	1411	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 103.

The worst 5 of 1411 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:M:750:UQ:C12	6:M:750:UQ:C13	1.84	1.50
1:L:183:ASN:ND2	1:L:236:LEU:HB2	1.24	1.48
2:M:96:PRO:HG2	2:M:170:SER:CA	1.57	1.33
1:L:183:ASN:HD22	1:L:236:LEU:CB	1.44	1.30
2:M:96:PRO:CG	2:M:170:SER:HA	1.75	1.17

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:M:293:TYR:OH	3:H:166:ASP:OD1[4_445]	1.59	0.61
1:L:265:TRP:CB	3:H:49:PRO:O[1_556]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	276/281 (98%)	187 (68%)	52 (19%)	37 (13%)	0	1
2	M	303/307 (99%)	189 (62%)	72 (24%)	42 (14%)	0	1
3	H	253/260 (97%)	144 (57%)	65 (26%)	44 (17%)	0	0
All	All	832/848 (98%)	520 (62%)	189 (23%)	123 (15%)	0	0

5 of 123 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	7	ARG
1	L	20	ASN
1	L	55	LEU
1	L	60	ASN
1	L	74	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	216/220 (98%)	170 (79%)	46 (21%)	1	5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	M	238/240 (99%)	179 (75%)	59 (25%)	1	2
3	H	203/208 (98%)	151 (74%)	52 (26%)	0	2
All	All	657/668 (98%)	500 (76%)	157 (24%)	1	3

5 of 157 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	45	GLU
3	H	194	GLN
3	H	68	HIS
3	H	147	ASN
3	H	223	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
2	M	257	ASN
3	H	68	HIS
3	H	206	ASN
3	H	129	ASN
3	H	9	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BPH	L	550	-	59,70,70	2.06	11 (18%)	59,101,101	1.80	14 (23%)
4	BCL	M	601	2	69,74,74	1.35	11 (15%)	79,115,115	1.63	10 (12%)
4	BCL	L	350	-	69,74,74	1.53	11 (15%)	79,115,115	1.79	13 (16%)
5	BPH	M	500	-	59,70,70	2.04	13 (22%)	59,101,101	1.75	12 (20%)
6	UQ	L	800	-	48,48,63	2.41	15 (31%)	60,61,79	2.38	25 (41%)
4	BCL	M	400	2	69,74,74	1.65	12 (17%)	79,115,115	1.63	13 (16%)
6	UQ	M	750	-	48,48,63	2.31	11 (22%)	60,61,79	1.92	18 (30%)
4	BCL	L	450	-	69,74,74	1.31	12 (17%)	79,115,115	1.40	12 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BPH	L	550	-	-	13/37/105/105	0/5/6/6
4	BCL	M	601	2	-	20/41/137/137	-
4	BCL	L	350	-	-	20/41/137/137	-
5	BPH	M	500	-	-	15/37/105/105	0/5/6/6
6	UQ	L	800	-	-	14/45/69/87	0/1/1/1
4	BCL	M	400	2	-	24/41/137/137	-
6	UQ	M	750	-	-	19/45/69/87	0/1/1/1
4	BCL	L	450	-	-	14/41/137/137	-

The worst 5 of 96 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	M	750	UQ	C12-C13	11.08	1.84	1.50
5	L	550	BPH	C2C-C3C	-10.60	1.45	1.54
5	M	500	BPH	C2C-C3C	-9.91	1.46	1.54
6	L	800	UQ	C16-C17	-8.98	1.24	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	M	400	BCL	MG-NA	8.79	2.27	2.06

The worst 5 of 117 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	M	400	BCL	C4A-NA-C1A	7.89	110.28	106.68
5	L	550	BPH	C4D-CHA-CBD	-7.22	104.99	108.45
4	L	350	BCL	C4A-NA-C1A	7.01	109.88	106.68
4	L	350	BCL	C4D-CHA-C1A	6.98	129.57	121.24
4	M	601	BCL	C4A-NA-C1A	6.46	109.63	106.68

There are no chirality outliers.

5 of 139 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	350	BCL	C6-C7-C8-C9
4	L	450	BCL	C1A-C2A-CAA-CBA
4	L	450	BCL	C3A-C2A-CAA-CBA
4	L	450	BCL	C2C-C3C-CAC-CBC
4	L	450	BCL	C4C-C3C-CAC-CBC


There are no ring outliers.



8 monomers are involved in 183 short contacts:













Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	L	550	BPH	16	0
4	M	601	BCL	25	0
4	L	350	BCL	30	0
5	M	500	BPH	31	0
6	L	800	UQ	22	0
4	M	400	BCL	44	0
6	M	750	UQ	33	0
4	L	450	BCL	20	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand UQ L 800	
 Bond lengths	 Bond angles
 Torsions	 Rings
Ligand BCL M 400	
 Bond lengths	 Bond angles
 Torsions	 Rings
Ligand UQ M 750	
 Bond lengths	 Bond angles
 Torsions	 Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.