



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 01:42 AM UTC

PDB ID : 7RC9 / pdb_00007rc9
Title : Crystal structure of human TLR8 ectodomain bound to small molecule antagonist 21
Authors : Critton, D.A.
Deposited on : 2021-07-07
Resolution : 2.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	811	3% 83% 7% 9%
1	B	811	7% 83% 7% 9%
2	C	3	100%
2	D	3	67% 33%
3	E	5	20% 80%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	5	 60% 40%
3	H	5	 40% 60%
4	G	4	 75% 25%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12205 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toll-like receptor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	737	Total	C	N	O	S	0	2	0
			5725	3683	956	1067	19			
1	B	737	Total	C	N	O	S	0	2	0
			5729	3685	957	1068	19			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	23	ARG	-	expression tag	UNP Q9NR97
A	24	SER	-	expression tag	UNP Q9NR97
A	25	PRO	-	expression tag	UNP Q9NR97
A	26	TRP	-	expression tag	UNP Q9NR97
A	828	GLU	-	expression tag	UNP Q9NR97
A	829	PHE	-	expression tag	UNP Q9NR97
A	830	LEU	-	expression tag	UNP Q9NR97
A	831	VAL	-	expression tag	UNP Q9NR97
A	832	PRO	-	expression tag	UNP Q9NR97
A	833	ARG	-	expression tag	UNP Q9NR97
B	23	ARG	-	expression tag	UNP Q9NR97
B	24	SER	-	expression tag	UNP Q9NR97
B	25	PRO	-	expression tag	UNP Q9NR97
B	26	TRP	-	expression tag	UNP Q9NR97
B	828	GLU	-	expression tag	UNP Q9NR97
B	829	PHE	-	expression tag	UNP Q9NR97
B	830	LEU	-	expression tag	UNP Q9NR97
B	831	VAL	-	expression tag	UNP Q9NR97
B	832	PRO	-	expression tag	UNP Q9NR97
B	833	ARG	-	expression tag	UNP Q9NR97

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			39	22	2	15			
2	D	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	5	Total	C	N	O	0	0	0
			61	34	2	25			
3	F	5	Total	C	N	O	0	0	0
			61	34	2	25			
3	H	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is 2-(2,6-dimethylpyridin-4-yl)-5-(piperidin-4-yl)-3-(propan-2-yl)-1H-indole (CCD ID: RQY) (formula: C₂₃H₂₉N₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	N	0	0
			26	23	3		
6	B	1	Total	C	N	0	0
			26	23	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	110	Total	O	0	0
			110	110		
7	B	82	Total	O	0	0
			82	82		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:  3% 83% 7% 9%



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%


NAG1
NAG2
BMA3

- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  67% 33%

NAG1
NAG2
BMA3

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  20% 80%

NAG1
NAG2
BMA3
MAN4
MAN5

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  60% 40%

NAG1
NAG2
BMA3
MAN4
MAN5

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  40% 60%

NAG1
NAG2
BMA3
MAN4
MAN5

- Molecule 4: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  75% 25%

NAG1
NAG2
BMA3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	160.89Å 83.88Å 153.52Å 90.00° 119.72° 90.00°	Depositor
Resolution (Å)	40.41 – 2.76 40.41 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.0 (40.41-2.76) 99.4 (40.41-2.76)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.77Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.234 , 0.262 0.247 , 0.276	Depositor DCC
R_{free} test set	2331 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	63.7	Xtriage
Anisotropy	0.633	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 61.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.009 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12205	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, RQY, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/5852	1.16	12/7979 (0.2%)
1	B	0.73	0/5856	1.16	9/7984 (0.1%)
All	All	0.73	0/11708	1.16	21/15963 (0.1%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	604	ASN	CA-C-N	7.27	134.78	121.70
1	A	604	ASN	C-N-CA	7.27	134.78	121.70
1	B	604	ASN	CA-C-N	7.25	134.75	121.70
1	B	604	ASN	C-N-CA	7.25	134.75	121.70
1	B	377	TYR	N-CA-C	-6.40	104.52	112.90
1	A	377	TYR	N-CA-C	-6.33	104.60	112.90
1	A	633	SER	CA-C-N	5.82	128.74	120.53
1	A	633	SER	C-N-CA	5.82	128.74	120.53
1	B	633	SER	CA-C-N	5.80	128.70	120.53
1	B	633	SER	C-N-CA	5.80	128.70	120.53
1	A	524	THR	CA-C-N	5.40	127.46	120.44
1	A	524	THR	C-N-CA	5.40	127.46	120.44
1	B	524	THR	CA-C-N	5.40	127.46	120.44
1	B	524	THR	C-N-CA	5.40	127.46	120.44
1	A	605	LEU	CA-C-N	5.21	131.50	121.54
1	A	605	LEU	C-N-CA	5.21	131.50	121.54
1	A	395	ASN	CA-CB-CG	5.10	117.70	112.60
1	A	187	CYS	CA-C-N	5.03	130.75	121.70
1	A	187	CYS	C-N-CA	5.03	130.75	121.70
1	B	187	CYS	CA-C-N	5.00	130.70	121.70
1	B	187	CYS	C-N-CA	5.00	130.70	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5725	0	5496	17	0
1	B	5729	0	5502	19	0
2	C	39	0	34	0	0
2	D	39	0	34	0	0
3	E	61	0	52	0	0
3	F	61	0	52	0	0
3	H	61	0	52	0	0
4	G	50	0	43	0	0
5	A	98	0	91	0	0
5	B	98	0	91	0	0
6	A	26	0	0	0	0
6	B	26	0	0	1	0
7	A	110	0	0	0	0
7	B	82	0	0	0	0
All	All	12205	0	11447	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:187:CYS:N	1:B:188:GLU:HA	2.07	0.69
1:A:187:CYS:N	1:A:188:GLU:HA	2.07	0.68
1:B:409:ILE:HD12	1:B:430:ILE:HD11	1.86	0.58
1:A:732:LEU:HD12	1:A:756:LEU:HA	1.84	0.57
1:B:732:LEU:HD12	1:B:756:LEU:HA	1.85	0.57
1:A:409:ILE:HD12	1:A:430:ILE:HD11	1.88	0.56
1:B:382:LEU:HB3	1:B:409:ILE:HG12	1.92	0.52
1:A:382:LEU:HB3	1:A:409:ILE:HG12	1.92	0.52
1:A:532:VAL:HB	1:A:553:LEU:HD22	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:532:VAL:HB	1:B:553:LEU:HD22	1.94	0.48
1:B:154:ILE:HD11	1:B:176:TYR:HB3	1.97	0.47
1:A:86:LEU:HB3	1:A:89:LEU:HD12	1.97	0.46
1:A:179:TRP:NE1	1:A:463:PRO:HB3	2.31	0.46
1:B:572:GLY:C	1:B:600:THR:HG21	2.41	0.46
1:B:86:LEU:HB3	1:B:89:LEU:HD12	1.97	0.45
1:B:179:TRP:NE1	1:B:463:PRO:HB3	2.31	0.45
1:B:520:VAL:HA	1:B:543:ASP:HB3	1.99	0.45
1:A:67:GLU:HG2	1:A:91:LYS:HB3	1.99	0.45
1:A:89:LEU:HD13	1:A:92:ILE:HD11	1.99	0.45
1:B:67:GLU:HG2	1:B:91:LYS:HB3	1.99	0.44
1:A:612:GLU:HG3	1:A:643:ARG:HB3	1.98	0.44
1:A:154:ILE:HD11	1:A:176:TYR:HB3	1.98	0.44
1:B:373:HIS:HA	1:B:400:ASN:HB3	1.99	0.44
1:A:207:SER:HA	1:A:228:PHE:HB2	2.00	0.44
1:A:373:HIS:HA	1:A:400:ASN:HB3	1.99	0.44
1:B:89:LEU:HD13	1:B:92:ILE:HD11	2.00	0.44
1:B:302:ILE:HD11	1:B:323:LEU:HD13	2.00	0.44
1:A:670:HIS:HA	1:A:694:ASP:HB3	1.99	0.43
1:A:572:GLY:C	1:A:600:THR:HG21	2.43	0.43
1:B:670:HIS:HA	1:B:694:ASP:HB3	1.99	0.43
1:B:239:GLU:HA	1:B:282:ALA:HA	2.01	0.43
1:A:296:SER:HA	1:A:320:PHE:O	2.19	0.42
1:B:207:SER:HA	1:B:228:PHE:HB2	2.01	0.41
1:A:496:ILE:HD12	1:A:519:GLN:HG2	2.04	0.40
1:B:261:PHE:HZ	6:B:908:RQY:C24	2.35	0.40
1:B:133:GLU:HG2	1:B:154:ILE:HB	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	727/811 (90%)	673 (93%)	51 (7%)	3 (0%)	30	50
1	B	727/811 (90%)	671 (92%)	54 (7%)	2 (0%)	36	56
All	All	1454/1622 (90%)	1344 (92%)	105 (7%)	5 (0%)	36	56

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	605	LEU
1	B	605	LEU
1	A	606	GLU
1	A	378	VAL
1	B	378	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	624/755 (83%)	599 (96%)	25 (4%)	28	50
1	B	625/755 (83%)	604 (97%)	21 (3%)	32	57
All	All	1249/1510 (83%)	1203 (96%)	46 (4%)	30	54

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	CYS
1	A	170	ILE
1	A	181	CYS
1	A	190	THR
1	A	248	LEU
1	A	261	PHE
1	A	287	THR
1	A	289	LEU
1	A	296	SER
1	A	403	ILE
1	A	406	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	420	LEU
1	A	516	SER
1	A	529	ILE
1	A	567	TYR
1	A	585	ASN
1	A	591	LEU
1	A	605	LEU
1	A	632	ILE
1	A	638	LEU
1	A	671	ILE
1	A	703	LEU
1	A	727	LEU
1	A	735	VAL
1	A	750	THR
1	B	49	CYS
1	B	170	ILE
1	B	181	CYS
1	B	190	THR
1	B	248	LEU
1	B	285	ASN
1	B	289	LEU
1	B	403	ILE
1	B	406	ILE
1	B	420	LEU
1	B	529	ILE
1	B	585	ASN
1	B	591	LEU
1	B	605	LEU
1	B	632	ILE
1	B	640	ASN
1	B	671	ILE
1	B	703	LEU
1	B	727	LEU
1	B	735	VAL
1	B	750	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	A	97	ASN
1	A	135	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	174	ASN
1	A	191	ASN
1	A	231	ASN
1	A	277	ASN
1	A	380	GLN
1	A	400	ASN
1	B	77	HIS
1	B	97	ASN
1	B	135	ASN
1	B	174	ASN
1	B	191	ASN
1	B	277	ASN
1	B	284	GLN
1	B	373	HIS
1	B	380	GLN
1	B	388	GLN
1	B	400	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

25 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.30	0	17,19,21	0.60	0
2	NAG	C	2	2	14,14,15	0.29	0	17,19,21	0.69	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	C	3	2	11,11,12	0.32	0	15,15,17	0.51	0
2	NAG	D	1	2,1	14,14,15	0.34	0	17,19,21	0.54	0
2	NAG	D	2	2	14,14,15	0.34	0	17,19,21	1.61	2 (11%)
2	BMA	D	3	2	11,11,12	0.37	0	15,15,17	0.39	0
3	NAG	E	1	1,3	14,14,15	0.30	0	17,19,21	0.68	1 (5%)
3	NAG	E	2	3	14,14,15	0.30	0	17,19,21	0.62	1 (5%)
3	BMA	E	3	3	11,11,12	0.29	0	15,15,17	0.55	0
3	MAN	E	4	3	11,11,12	0.36	0	15,15,17	0.80	1 (6%)
3	MAN	E	5	3	11,11,12	0.36	0	15,15,17	0.81	1 (6%)
3	NAG	F	1	1,3	14,14,15	0.32	0	17,19,21	0.62	0
3	NAG	F	2	3	14,14,15	0.34	0	17,19,21	0.60	0
3	BMA	F	3	3	11,11,12	0.32	0	15,15,17	0.63	0
3	MAN	F	4	3	11,11,12	0.41	0	15,15,17	0.82	1 (6%)
3	MAN	F	5	3	11,11,12	0.36	0	15,15,17	0.79	1 (6%)
4	NAG	G	1	4,1	14,14,15	0.30	0	17,19,21	0.58	0
4	NAG	G	2	4	14,14,15	0.30	0	17,19,21	0.59	0
4	BMA	G	3	4	11,11,12	0.39	0	15,15,17	0.74	0
4	MAN	G	4	4	11,11,12	0.38	0	15,15,17	1.00	1 (6%)
3	NAG	H	1	1,3	14,14,15	0.30	0	17,19,21	0.71	1 (5%)
3	NAG	H	2	3	14,14,15	0.28	0	17,19,21	0.58	0
3	BMA	H	3	3	11,11,12	0.30	0	15,15,17	0.55	0
3	MAN	H	4	3	11,11,12	0.37	0	15,15,17	0.88	1 (6%)
3	MAN	H	5	3	11,11,12	0.40	0	15,15,17	1.00	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
3	NAG	E	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1
3	MAN	E	4	3	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	E	5	3	-	0/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1
3	MAN	F	4	3	-	0/2/19/22	0/1/1/1
3	MAN	F	5	3	-	0/2/19/22	0/1/1/1
4	NAG	G	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	0/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	MAN	G	4	4	-	0/2/19/22	0/1/1/1
3	NAG	H	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	H	2	3	-	0/6/23/26	0/1/1/1
3	BMA	H	3	3	-	0/2/19/22	0/1/1/1
3	MAN	H	4	3	-	0/2/19/22	0/1/1/1
3	MAN	H	5	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	NAG	O5-C1-C2	-5.99	102.02	111.29
4	G	4	MAN	C1-O5-C5	3.54	116.94	112.19
3	H	5	MAN	C1-O5-C5	3.36	116.68	112.19
3	H	4	MAN	C1-O5-C5	3.00	116.21	112.19
3	E	5	MAN	C1-O5-C5	2.52	115.57	112.19
3	F	5	MAN	C1-O5-C5	2.52	115.56	112.19
3	E	4	MAN	C1-O5-C5	2.50	115.54	112.19
3	F	4	MAN	C1-O5-C5	2.29	115.25	112.19
3	H	1	NAG	C1-O5-C5	2.11	115.01	112.19
2	D	2	NAG	O4-C4-C3	-2.08	105.47	110.38
3	E	2	NAG	C1-O5-C5	2.07	114.97	112.19
3	E	1	NAG	C1-O5-C5	2.07	114.96	112.19

There are no chirality outliers.


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C4-C5-C6-O6


There are no ring outliers.



No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	B	902	1	14,14,15	0.28	0	17,19,21	0.69	1 (5%)
6	RQY	B	908	-	29,29,29	5.39	15 (51%)	39,42,42	1.26	5 (12%)
5	NAG	A	903	1	14,14,15	0.33	0	17,19,21	0.63	0
5	NAG	B	906	1	14,14,15	0.34	0	17,19,21	0.83	1 (5%)
5	NAG	A	904	1	14,14,15	0.32	0	17,19,21	0.75	1 (5%)
5	NAG	B	905	1	14,14,15	0.32	0	17,19,21	0.47	0
5	NAG	B	907	1	14,14,15	0.32	0	17,19,21	0.75	2 (11%)
5	NAG	A	902	1	14,14,15	0.31	0	17,19,21	0.59	0
5	NAG	B	901	1	14,14,15	0.30	0	17,19,21	0.52	0
5	NAG	A	905	1	14,14,15	0.30	0	17,19,21	0.53	0
5	NAG	A	907	1	14,14,15	0.31	0	17,19,21	0.76	2 (11%)
5	NAG	B	903	1	14,14,15	0.33	0	17,19,21	0.65	0
5	NAG	A	906	1	14,14,15	0.29	0	17,19,21	0.48	0
5	NAG	A	901	1	14,14,15	0.30	0	17,19,21	0.56	0
6	RQY	A	908	-	29,29,29	5.40	15 (51%)	39,42,42	1.31	6 (15%)
5	NAG	B	904	1	14,14,15	0.34	0	17,19,21	0.80	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	902	1	-	0/6/23/26	0/1/1/1
6	RQY	B	908	-	-	0/12/20/20	0/4/4/4
5	NAG	A	903	1	-	0/6/23/26	0/1/1/1
5	NAG	B	906	1	-	0/6/23/26	0/1/1/1
5	NAG	A	904	1	-	0/6/23/26	0/1/1/1
5	NAG	B	905	1	-	0/6/23/26	0/1/1/1
5	NAG	B	907	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	902	1	-	0/6/23/26	0/1/1/1
5	NAG	B	901	1	-	2/6/23/26	0/1/1/1
5	NAG	A	905	1	-	0/6/23/26	0/1/1/1
5	NAG	A	907	1	-	0/6/23/26	0/1/1/1
5	NAG	B	903	1	-	0/6/23/26	0/1/1/1
5	NAG	A	906	1	-	1/6/23/26	0/1/1/1
5	NAG	A	901	1	-	2/6/23/26	0/1/1/1
6	RQY	A	908	-	-	0/12/20/20	0/4/4/4
5	NAG	B	904	1	-	0/6/23/26	0/1/1/1

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	908	RQY	C06-C05	12.36	1.59	1.39
6	B	908	RQY	C06-C05	12.20	1.58	1.39
6	A	908	RQY	C03-C04	11.21	1.57	1.39
6	B	908	RQY	C03-C04	11.21	1.57	1.39
6	A	908	RQY	C06-C01	8.38	1.51	1.39
6	B	908	RQY	C06-C01	8.26	1.51	1.39
6	A	908	RQY	C20-N21	7.96	1.49	1.34
6	B	908	RQY	C19-C20	7.89	1.53	1.38
6	B	908	RQY	C20-N21	7.86	1.49	1.34
6	B	908	RQY	C22-N21	7.85	1.49	1.34
6	A	908	RQY	C22-N21	7.84	1.49	1.34
6	A	908	RQY	C23-C22	7.84	1.53	1.38
6	B	908	RQY	C23-C22	7.66	1.53	1.38
6	B	908	RQY	C03-C02	7.59	1.51	1.38
6	A	908	RQY	C03-C02	7.49	1.50	1.38
6	A	908	RQY	C19-C20	7.46	1.52	1.38
6	A	908	RQY	C23-C18	7.01	1.50	1.39
6	A	908	RQY	C19-C18	7.00	1.50	1.39
6	B	908	RQY	C19-C18	6.93	1.49	1.39
6	B	908	RQY	C23-C18	6.83	1.49	1.39
6	A	908	RQY	C02-C01	6.58	1.49	1.39
6	B	908	RQY	C02-C01	6.54	1.49	1.39
6	B	908	RQY	C05-C09	-3.92	1.38	1.44
6	A	908	RQY	C05-C09	-3.84	1.38	1.44
6	B	908	RQY	C18-C08	3.69	1.53	1.47
6	A	908	RQY	C01-C10	3.66	1.59	1.52
6	B	908	RQY	C01-C10	3.66	1.59	1.52
6	A	908	RQY	C18-C08	3.58	1.53	1.47
6	A	908	RQY	C08-C09	2.42	1.40	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	908	RQY	C08-C09	2.34	1.40	1.38

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	908	RQY	C26-C20-N21	2.90	120.95	116.56
6	B	908	RQY	C25-C22-N21	2.69	120.62	116.56
6	A	908	RQY	C17-C10-C13	2.68	115.51	109.68
5	B	906	NAG	C1-O5-C5	2.57	115.63	112.19
6	B	908	RQY	C18-C08-N07	2.56	122.97	119.74
6	B	908	RQY	C26-C20-N21	2.45	120.26	116.56
6	A	908	RQY	C24-C11-C09	-2.44	107.47	112.90
6	A	908	RQY	C18-C08-N07	2.41	122.77	119.74
5	B	902	NAG	O5-C1-C2	-2.36	107.63	111.29
6	A	908	RQY	C25-C22-N21	2.28	120.00	116.56
5	B	904	NAG	C1-C2-N2	2.19	113.89	110.43
6	B	908	RQY	C18-C08-C09	-2.16	128.57	131.29
5	A	904	NAG	C1-C2-N2	2.10	113.75	110.43
5	B	907	NAG	C1-O5-C5	2.08	114.98	112.19
5	A	907	NAG	O5-C1-C2	-2.05	108.11	111.29
6	B	908	RQY	C17-C10-C13	2.05	114.13	109.68
5	A	907	NAG	C1-O5-C5	2.04	114.92	112.19
6	A	908	RQY	C18-C08-C09	-2.04	128.73	131.29
5	B	907	NAG	O5-C1-C2	-2.02	108.17	111.29

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	901	NAG	O5-C5-C6-O6
5	A	901	NAG	O5-C5-C6-O6
5	A	906	NAG	O5-C5-C6-O6
5	A	901	NAG	C1-C2-N2-C7
5	B	901	NAG	C1-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	908	RQY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	737/811 (90%)	0.28	27 (3%) 45 42	39, 78, 122, 165	2 (0%)
1	B	737/811 (90%)	0.64	54 (7%) 21 21	41, 101, 149, 182	2 (0%)
All	All	1474/1622 (90%)	0.46	81 (5%) 30 30	39, 87, 140, 182	4 (0%)

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	92	ILE	5.6
1	A	606	GLU	4.2
1	B	236	TYR	3.8
1	B	417	PHE	3.8
1	B	605	LEU	3.7
1	B	65	VAL	3.7
1	A	818	GLU	3.7
1	B	89	LEU	3.6
1	B	470	PHE	3.5
1	A	605	LEU	3.4
1	B	732	LEU	3.4
1	A	236	TYR	3.4
1	B	392	GLN	3.2
1	B	83	PHE	3.1
1	A	703	LEU	3.1
1	B	151	LEU	3.1
1	B	751	ILE	3.1
1	B	121	PHE	3.0
1	B	702	PHE	3.0
1	B	679	PHE	3.0
1	A	783	PHE	3.0
1	B	678	PHE	3.0
1	B	46	ILE	2.9
1	A	762	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	94	LEU	2.8
1	A	607	SER	2.8
1	B	64	TYR	2.7
1	B	78	ILE	2.7
1	B	45	VAL	2.7
1	B	603	TYR	2.6
1	B	132	LEU	2.6
1	B	767	LEU	2.6
1	A	678	PHE	2.6
1	B	187	CYS	2.6
1	A	260	CYS	2.5
1	A	130	LEU	2.5
1	B	74	PHE	2.5
1	A	817	LEU	2.5
1	B	130	LEU	2.5
1	B	162	THR	2.5
1	A	707	LEU	2.4
1	B	61	VAL	2.4
1	A	604	ASN	2.4
1	A	756	LEU	2.4
1	B	122	LEU	2.4
1	B	606	GLU	2.3
1	A	786	TRP	2.3
1	B	127	LEU	2.3
1	A	750	THR	2.3
1	B	203	LEU	2.3
1	B	192	ILE	2.3
1	B	330	GLY	2.3
1	B	141	PRO	2.2
1	A	39	LYS	2.2
1	B	67	GLU	2.2
1	B	190	THR	2.2
1	A	732	LEU	2.2
1	B	116	ILE	2.2
1	B	414	PHE	2.2
1	B	637	GLY	2.2
1	A	261	PHE	2.2
1	A	34	TYR	2.2
1	B	75	ILE	2.2
1	B	653	HIS	2.2
1	A	136	GLN	2.1
1	A	769	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	57	VAL	2.1
1	B	795	ILE	2.1
1	A	435	LYS	2.1
1	B	604	ASN	2.1
1	B	96	HIS	2.1
1	B	762	THR	2.1
1	B	55	GLN	2.1
1	A	795	ILE	2.1
1	B	387	PHE	2.1
1	B	780	ILE	2.0
1	A	32	ARG	2.0
1	A	734	GLU	2.0
1	B	731	PHE	2.0
1	B	62	GLY	2.0
1	B	161	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MAN	G	4	11/12	0.32	0.12	119,123,124,126	0
3	MAN	F	5	11/12	0.41	0.13	123,126,132,133	0
2	BMA	D	3	11/12	0.47	0.12	92,95,97,97	0
3	MAN	F	4	11/12	0.47	0.12	115,118,121,122	0
3	MAN	E	5	11/12	0.65	0.10	98,100,104,106	0
4	BMA	G	3	11/12	0.67	0.10	102,106,110,115	0
3	MAN	H	4	11/12	0.67	0.09	109,112,117,119	0
3	MAN	E	4	11/12	0.68	0.10	92,97,98,100	0
3	BMA	F	3	11/12	0.69	0.10	110,116,122,123	0
3	MAN	H	5	11/12	0.73	0.07	117,120,123,125	0
3	BMA	H	3	11/12	0.76	0.08	97,105,109,113	0
2	BMA	C	3	11/12	0.79	0.09	78,80,84,85	0
3	BMA	E	3	11/12	0.81	0.10	81,88,92,95	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	G	2	14/15	0.81	0.11	81,85,90,97	0
2	NAG	D	2	14/15	0.87	0.10	73,79,84,87	0
3	NAG	F	2	14/15	0.91	0.12	93,97,100,103	0
3	NAG	H	1	14/15	0.92	0.08	64,67,73,74	0
3	NAG	E	2	14/15	0.92	0.08	60,66,69,75	0
4	NAG	G	1	14/15	0.92	0.09	63,67,69,74	0
2	NAG	C	2	14/15	0.93	0.11	63,67,69,73	0
3	NAG	H	2	14/15	0.93	0.06	73,78,84,90	0
2	NAG	D	1	14/15	0.93	0.10	52,55,59,66	0
2	NAG	C	1	14/15	0.94	0.09	51,56,59,60	0
3	NAG	F	1	14/15	0.95	0.10	81,85,91,91	0
3	NAG	E	1	14/15	0.96	0.07	49,53,57,58	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain H:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	B	901	14/15	0.37	0.15	181,186,196,197	0
5	NAG	A	903	14/15	0.38	0.14	120,124,128,129	0
5	NAG	A	906	14/15	0.48	0.14	103,119,127,132	0
5	NAG	B	906	14/15	0.55	0.13	132,137,141,142	0
5	NAG	B	907	14/15	0.56	0.13	141,144,149,152	0
5	NAG	B	905	14/15	0.59	0.13	99,102,106,109	0
5	NAG	B	904	14/15	0.60	0.20	147,152,158,158	0
5	NAG	A	905	14/15	0.61	0.15	112,113,121,121	0
5	NAG	B	902	14/15	0.63	0.12	159,164,170,174	0
5	NAG	B	903	14/15	0.64	0.18	142,148,154,158	0
5	NAG	A	904	14/15	0.69	0.14	126,132,139,141	0
5	NAG	A	902	14/15	0.69	0.10	120,123,129,130	0
5	NAG	A	901	14/15	0.72	0.10	125,129,138,139	0
5	NAG	A	907	14/15	0.88	0.08	101,107,112,115	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	RQY	A	908	26/26	0.91	0.10	46,50,55,56	0
6	RQY	B	908	26/26	0.93	0.10	43,51,58,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.