



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 28, 2026 – 01:36 AM UTC

PDB ID : 2RDI / pdb_00002rdi
Title : Snapshots of a Y-family DNA polymerase in replication: Dpo4 in apo and binary/ternary complex forms
Authors : Wong, J.H.Y.; Ling, H.
Deposited on : 2007-09-24
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	20% 78% 16% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	354	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2949 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	0	0
			2701	1735	464	495	7			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLU	deletion	UNP Q97W02
A	?	-	ALA	deletion	UNP Q97W02
A	?	-	ILE	deletion	UNP Q97W02
A	?	-	GLY	deletion	UNP Q97W02
A	?	-	LEU	deletion	UNP Q97W02
A	?	-	ASP	deletion	UNP Q97W02
A	?	-	LYS	deletion	UNP Q97W02
A	?	-	PHE	deletion	UNP Q97W02
A	?	-	PHE	deletion	UNP Q97W02
A	?	-	ASP	deletion	UNP Q97W02

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	235	Total 235	O 235	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA polymerase IV



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.93Å 51.07Å 86.03Å 90.00° 98.08° 90.00°	Depositor
Resolution (Å)	20.35 – 1.92 20.35 – 1.92	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.35-1.92) 95.3 (20.35-1.92)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.205 , 0.247 0.272 , 0.308	Depositor DCC
R_{free} test set	1217 reflections (4.15%)	wwPDB-VP
Wilson B-factor (Å ²)	36.8	Xtriage
Anisotropy	0.164	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2949	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/2738	0.93	0/3677

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	240	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2701	0	2855	56	0
2	A	7	0	10	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	6	0	8	9	0
4	A	235	0	0	9	0
All	All	2949	0	2873	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:LEU:HG	4:A:582:HOH:O	1.52	1.07
1:A:14:GLN:HE22	1:A:139:THR:H	1.17	0.93
1:A:252:LYS:HG2	1:A:264:TYR:OH	1.73	0.89
1:A:10:TYR:HE2	3:A:354:GOL:HO3	1.25	0.81
1:A:241:VAL:HG12	1:A:241:VAL:O	1.82	0.79
1:A:240:ARG:HA	1:A:241:VAL:HB	1.67	0.77
1:A:31:CYS:HB3	1:A:61:ILE:HD11	1.69	0.75
1:A:197:LEU:HD12	1:A:216:MET:HE1	1.75	0.68
3:A:354:GOL:H32	4:A:548:HOH:O	1.96	0.65
1:A:240:ARG:HA	1:A:241:VAL:CB	2.25	0.65
1:A:197:LEU:CD1	1:A:216:MET:HE1	2.28	0.63
1:A:10:TYR:HD2	3:A:354:GOL:HO1	1.48	0.61
1:A:12:TYR:HB2	1:A:45:THR:CG2	2.30	0.61
1:A:174:LEU:HB3	1:A:178:LEU:HD22	1.85	0.58
1:A:259:GLU:HA	1:A:259:GLU:OE1	2.02	0.58
1:A:259:GLU:OE1	1:A:262:LYS:HD3	2.03	0.58
1:A:76:MET:HE3	1:A:78:LYS:HB2	1.85	0.57
1:A:10:TYR:HD2	3:A:354:GOL:O1	1.88	0.57
1:A:257:ASN:HB3	1:A:260:GLU:HB3	1.87	0.56
1:A:285:HIS:HD2	4:A:415:HOH:O	1.87	0.56
1:A:10:TYR:HE2	3:A:354:GOL:O3	1.88	0.55
1:A:240:ARG:CA	1:A:241:VAL:HB	2.37	0.55
1:A:278:LYS:HE3	1:A:280:ILE:HD11	1.90	0.53
1:A:12:TYR:HB2	1:A:45:THR:HG21	1.90	0.52
1:A:278:LYS:HE2	1:A:278:LYS:O	2.10	0.52
1:A:57:ALA:HB2	3:A:354:GOL:H12	1.92	0.51
1:A:278:LYS:CE	1:A:280:ILE:HD11	2.41	0.50
1:A:252:LYS:HG2	1:A:264:TYR:HH	1.70	0.50
1:A:32:VAL:CG2	1:A:44:ALA:HB2	2.42	0.49
1:A:79:GLU:H	1:A:79:GLU:CD	2.19	0.49
1:A:241:VAL:O	1:A:241:VAL:CG1	2.55	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:ARG:HA	1:A:241:VAL:CG2	2.41	0.48
1:A:275:LYS:HG3	4:A:577:HOH:O	2.13	0.46
1:A:217:ILE:O	1:A:221:LYS:HE3	2.15	0.46
1:A:300:ARG:HD2	4:A:517:HOH:O	2.15	0.46
1:A:240:ARG:HA	1:A:241:VAL:HG23	1.99	0.45
1:A:12:TYR:HB2	1:A:45:THR:HG23	1.97	0.45
1:A:248:ILE:HD13	1:A:334:GLY:HA3	1.99	0.45
1:A:56:LYS:HG3	4:A:466:HOH:O	2.16	0.45
1:A:282:LYS:HD2	1:A:341:ILE:HG12	1.99	0.44
1:A:32:VAL:HG23	1:A:44:ALA:HB2	1.99	0.44
1:A:89:MET:HE3	1:A:101:ILE:HG23	2.00	0.44
3:A:354:GOL:C3	4:A:548:HOH:O	2.60	0.44
1:A:57:ALA:N	3:A:354:GOL:O2	2.51	0.43
1:A:106:GLU:OE2	1:A:152:LYS:NZ	2.49	0.43
1:A:99:ILE:HB	4:A:582:HOH:O	2.17	0.43
1:A:256:ARG:HG3	1:A:329:LYS:HA	2.01	0.43
1:A:31:CYS:HB3	1:A:61:ILE:CD1	2.45	0.42
1:A:68:LEU:HD13	1:A:71:ALA:HB2	2.01	0.42
1:A:40:SER:HA	1:A:41:GLY:HA3	1.84	0.41
1:A:174:LEU:O	1:A:178:LEU:HB2	2.20	0.41
1:A:256:ARG:NH2	1:A:323:LEU:O	2.50	0.41
1:A:188:ASN:HD22	1:A:188:ASN:HA	1.71	0.41
1:A:9:ASP:O	1:A:10:TYR:C	2.63	0.41
1:A:71:ALA:HB3	1:A:73:TYR:CE2	2.56	0.41
1:A:159:LYS:NZ	4:A:461:HOH:O	2.54	0.41
1:A:45:THR:HA	3:A:354:GOL:H12	2.02	0.41
1:A:284:ILE:HG12	1:A:285:HIS:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/342 (97%)	323 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/301 (98%)	272 (92%)	24 (8%)	11	2

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	THR
1	A	68	LEU
1	A	76	MET
1	A	79	GLU
1	A	90	ASN
1	A	138	ILE
1	A	178	LEU
1	A	188	ASN
1	A	194	LEU
1	A	196	LYS
1	A	202	LEU
1	A	234	ASN
1	A	238	ARG
1	A	244	SER
1	A	245	ILE
1	A	252	LYS
1	A	278	LYS
1	A	284	ILE
1	A	313	SER
1	A	321	LYS
1	A	323	LEU
1	A	335	VAL
1	A	341	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	352	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	GLN
1	A	82	GLN
1	A	90	ASN
1	A	188	ASN
1	A	234	ASN
1	A	254	ASN
1	A	320	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	354	-	5,5,5	0.66	0	5,5,5	1.03	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PEG	A	353	-	6,6,6	0.51	0	5,5,5	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	354	-	-	2/4/4/4	-
2	PEG	A	353	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	354	GOL	O1-C1-C2-C3
2	A	353	PEG	O1-C1-C2-O2
3	A	354	GOL	O1-C1-C2-O2
2	A	353	PEG	O2-C3-C4-O4

There are no ring outliers.

1 monomer is involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	354	GOL	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	336/342 (98%)	1.29	67 (19%) 3 4	20, 47, 57, 60	1 (0%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	352	THR	4.8
1	A	40	SER	3.4
1	A	316	VAL	3.3
1	A	194	LEU	3.3
1	A	309	GLU	3.2
1	A	41	GLY	3.2
1	A	253	ARG	3.2
1	A	317	LYS	3.1
1	A	83	GLN	3.0
1	A	116	ARG	3.0
1	A	198	GLY	3.0
1	A	62	VAL	3.0
1	A	341	ILE	3.0
1	A	239	THR	2.9
1	A	299	GLY	2.9
1	A	263	PRO	2.8
1	A	303	PRO	2.8
1	A	320	GLN	2.8
1	A	199	ILE	2.8
1	A	304	HIS	2.7
1	A	313	SER	2.7
1	A	30	VAL	2.7
1	A	216	MET	2.6
1	A	208	ILE	2.6
1	A	226	ILE	2.6
1	A	241	VAL	2.6
1	A	220	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	95	TYR	2.6
1	A	244	SER	2.6
1	A	257	ASN	2.5
1	A	311	ALA	2.5
1	A	46	ALA	2.5
1	A	94	GLU	2.5
1	A	289	VAL	2.5
1	A	248	ILE	2.5
1	A	33	PHE	2.4
1	A	114	LYS	2.4
1	A	28	VAL	2.4
1	A	312	TYR	2.4
1	A	310	THR	2.4
1	A	68	LEU	2.3
1	A	206	LEU	2.3
1	A	10	TYR	2.3
1	A	323	LEU	2.3
1	A	195	LYS	2.3
1	A	115	VAL	2.3
1	A	183	VAL	2.3
1	A	240	ARG	2.2
1	A	305	GLY	2.2
1	A	45	THR	2.2
1	A	69	PRO	2.2
1	A	197	LEU	2.2
1	A	25	GLY	2.2
1	A	210	PHE	2.2
1	A	72	VAL	2.1
1	A	209	GLU	2.1
1	A	67	ILE	2.1
1	A	117	ASP	2.1
1	A	321	LYS	2.1
1	A	20	ASN	2.1
1	A	228	LEU	2.1
1	A	261	ILE	2.1
1	A	74	LEU	2.1
1	A	245	ILE	2.0
1	A	252	LYS	2.0
1	A	120	GLU	2.0
1	A	215	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	354	6/6	0.75	0.21	59,62,65,65	0
2	PEG	A	353	7/7	0.76	0.14	47,48,49,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.