



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:26 PM UTC

PDB ID : 8RD2 / pdb_00008rd2
Title : Trypanosoma brucei Invariant Surface Glycoprotein 75 (ISG75)
Authors : Stodkilde-Jorgensen, K.; Mikkelsen, J.H.
Deposited on : 2023-12-07
Resolution : 2.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	 7% 74% 15% • 10%
1	B	440	 7% 72% 18% • 9%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6261 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Invariant surface glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	398	3121	1921	562	622	16	0	0	0
1	B	400	3140	1933	567	624	16	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	134	ALA	ASN	engineered mutation	UNP Q26769
B	134	ALA	ASN	engineered mutation	UNP Q26769


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Invariant surface glycoprotein



- Molecule 1: Invariant surface glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	120.01Å 120.01Å 107.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.67 – 2.69 53.67 – 2.69	Depositor EDS
% Data completeness (in resolution range)	52.0 (53.67-2.69) 52.1 (53.67-2.69)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 2.69Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.228 , 0.272 0.228 , 0.273	Depositor DCC
R_{free} test set	1223 reflections (2.88%)	wwPDB-VP
Wilson B-factor (Å ²)	82.7	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 62.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.041 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6261	wwPDB-VP
Average B, all atoms (Å ²)	94.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.11	0/3160	0.29	0/4240
1	B	0.11	0/3180	0.30	0/4268
All	All	0.11	0/6340	0.30	0/8508

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3121	0	3094	39	0
1	B	3140	0	3119	46	0
All	All	6261	0	6213	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:173:ARG:NH1	1:B:183:ASP:OD2	2.17	0.77
1:A:44:ARG:HD3	1:A:220:PRO:HA	1.69	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:289:GLU:HB3	1:A:293:GLN:H	1.55	0.71
1:A:216:ASN:HA	1:A:219:LYS:HD2	1.72	0.70
1:A:111:LYS:O	1:A:115:ASN:ND2	2.28	0.66
1:B:181:ARG:HH11	1:B:344:ARG:HH12	1.44	0.63
1:A:414:LYS:NZ	1:A:418:GLU:OE2	2.30	0.61
1:B:173:ARG:HH12	1:B:186:LYS:HA	1.67	0.59
1:A:96:ALA:HA	1:A:99:LEU:HD12	1.86	0.57
1:B:36:LYS:HD3	1:B:229:ILE:HD13	1.86	0.56
1:A:113:LEU:HD21	1:A:272:LEU:HD11	1.86	0.56
1:A:223:TYR:CZ	1:A:225:MET:HB2	2.41	0.55
1:A:350:ALA:HB1	1:A:360:VAL:HG21	1.89	0.55
1:B:302:GLU:HG2	1:B:414:LYS:HE3	1.90	0.54
1:B:60:LEU:HD13	1:B:208:MET:HA	1.90	0.53
1:B:57:MET:HA	1:B:60:LEU:HD12	1.90	0.53
1:A:251:ASN:OD1	1:A:251:ASN:N	2.34	0.52
1:B:345:LEU:HA	1:B:348:GLU:HB2	1.92	0.52
1:B:312:ARG:HD2	1:B:404:THR:HG23	1.91	0.52
1:B:429:GLU:O	1:B:433:GLU:HG2	2.10	0.52
1:B:308:GLU:O	1:B:312:ARG:N	2.43	0.51
1:A:312:ARG:HA	1:A:403:ALA:HB1	1.93	0.50
1:B:282:LYS:HB3	1:B:421:GLU:HG3	1.93	0.50
1:B:309:GLU:OE2	1:B:411:LYS:NZ	2.45	0.50
1:B:169:ASN:O	1:B:194:LYS:NZ	2.29	0.50
1:A:91:ASP:HB3	1:B:366:LYS:NZ	2.26	0.50
1:B:415:LYS:O	1:B:419:GLU:HG2	2.11	0.49
1:B:320:GLU:OE2	1:B:324:LYS:HE2	2.12	0.49
1:A:91:ASP:HB3	1:B:366:LYS:HZ1	1.77	0.49
1:A:177:TYR:CD2	1:A:180:SER:HB3	2.48	0.48
1:A:351:GLU:OE2	1:A:354:LYS:HD3	2.13	0.48
1:A:186:LYS:HG2	1:A:187:TYR:H	1.79	0.48
1:A:106:GLU:HA	1:A:109:TYR:HB3	1.94	0.48
1:A:178:GLU:O	1:A:179:HIS:HB3	2.13	0.48
1:B:274:LEU:HD21	1:B:412:GLU:OE2	2.14	0.47
1:A:99:LEU:HD13	1:A:104:LEU:HD12	1.96	0.47
1:B:329:ALA:HB1	1:B:386:ALA:HA	1.98	0.46
1:B:274:LEU:HD11	1:B:278:ARG:HE	1.80	0.46
1:A:210:ARG:HG3	1:A:211:ALA:N	2.30	0.46
1:B:124:THR:HG21	1:B:265:LYS:HD3	1.98	0.46
1:A:380:LYS:HA	1:A:383:GLU:OE2	2.16	0.45
1:B:89:ILE:HG23	1:B:95:MET:HG3	1.98	0.45
1:A:349:LYS:HD3	1:A:349:LYS:HA	1.79	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:ALA:HB1	1:A:414:LYS:HA	1.97	0.45
1:B:111:LYS:HE3	1:B:115:ASN:HD21	1.82	0.45
1:A:173:ARG:NH2	1:A:188:SER:O	2.49	0.45
1:B:341:GLU:HA	1:B:344:ARG:HD2	1.98	0.45
1:B:168:PRO:HA	1:B:171:TYR:CZ	2.52	0.45
1:B:365:LYS:HA	1:B:365:LYS:HD2	1.72	0.44
1:A:31:LEU:HG	1:A:38:TYR:CZ	2.52	0.44
1:B:251:ASN:OD1	1:B:251:ASN:N	2.51	0.44
1:B:64:VAL:HG13	1:B:263:ILE:HD11	1.99	0.44
1:B:89:ILE:HD11	1:B:284:LEU:HA	2.00	0.43
1:B:83:ASN:O	1:B:87:ARG:HG3	2.18	0.43
1:A:179:HIS:CG	1:A:179:HIS:O	2.71	0.43
1:B:274:LEU:O	1:B:278:ARG:HG2	2.18	0.43
1:A:31:LEU:HD11	1:A:227:VAL:HG13	2.00	0.43
1:A:357:GLY:O	1:A:359:PRO:HD3	2.19	0.43
1:B:304:ARG:O	1:B:308:GLU:HG2	2.19	0.43
1:B:401:VAL:O	1:B:405:GLU:HG2	2.19	0.42
1:A:162:PRO:HD3	1:A:187:TYR:CZ	2.55	0.42
1:A:199:THR:HA	1:A:400:ARG:HH22	1.85	0.42
1:B:138:ALA:O	1:B:142:VAL:HG23	2.20	0.42
1:A:340:ALA:O	1:A:344:ARG:HG2	2.20	0.42
1:B:330:GLU:OE1	1:B:333:ARG:NE	2.49	0.41
1:A:198:SER:N	1:A:201:THR:OG1	2.50	0.41
1:B:29:GLU:HG3	1:B:227:VAL:HG21	2.03	0.41
1:B:93:LYS:HA	1:B:96:ALA:HB3	2.03	0.41
1:B:382:ALA:HA	1:B:385:GLN:HB2	2.02	0.41
1:B:100:SER:HB2	1:B:103:ARG:HB3	2.03	0.41
1:B:275:ARG:HA	1:B:275:ARG:HD3	1.89	0.41
1:A:162:PRO:HB3	1:A:187:TYR:CD1	2.56	0.41
1:B:101:ALA:O	1:B:104:LEU:HB3	2.21	0.41
1:B:162:PRO:HD3	1:B:187:TYR:CZ	2.55	0.41
1:A:150:ARG:HE	1:A:150:ARG:HB3	1.64	0.40
1:A:225:MET:HE2	1:A:225:MET:HA	2.03	0.40
1:B:354:LYS:HA	1:B:359:PRO:O	2.22	0.40
1:A:38:TYR:HB2	1:A:222:PRO:HB3	2.02	0.40
1:A:58:LYS:HB2	1:A:139:LEU:HD23	2.03	0.40
1:A:219:LYS:NZ	1:A:250:GLU:OE2	2.53	0.40
1:A:285:GLU:C	1:A:287:ASP:H	2.30	0.40
1:B:111:LYS:HE2	1:B:111:LYS:HB3	1.78	0.40
1:B:106:GLU:O	1:B:110:MET:HG2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/440 (89%)	373 (95%)	18 (5%)	1 (0%)	36	60
1	B	396/440 (90%)	379 (96%)	16 (4%)	1 (0%)	36	60
All	All	788/880 (90%)	752 (95%)	34 (4%)	2 (0%)	36	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	359	PRO
1	A	236	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/354 (90%)	303 (95%)	16 (5%)	22	48
1	B	321/354 (91%)	307 (96%)	14 (4%)	25	53
All	All	640/708 (90%)	610 (95%)	30 (5%)	23	51

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	PHE
1	A	94	VAL
1	A	140	ASP
1	A	150	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	173	ARG
1	A	185	HIS
1	A	190	LEU
1	A	210	ARG
1	A	225	MET
1	A	251	ASN
1	A	275	ARG
1	A	291	LEU
1	A	295	GLU
1	A	343	ARG
1	A	360	VAL
1	A	408	GLU
1	B	112	SER
1	B	140	ASP
1	B	151	THR
1	B	173	ARG
1	B	195	VAL
1	B	237	HIS
1	B	277	MET
1	B	284	LEU
1	B	291	LEU
1	B	348	GLU
1	B	359	PRO
1	B	363	GLU
1	B	385	GLN
1	B	431	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	A	238	GLN
1	B	45	GLN
1	B	339	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	398/440 (90%)	0.55	32 (8%) 18 15	35, 77, 160, 236	0
1	B	400/440 (90%)	0.66	32 (8%) 18 15	39, 94, 165, 205	0
All	All	798/880 (90%)	0.60	64 (8%) 18 15	35, 86, 162, 236	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	290	SER	6.3
1	A	223	TYR	4.6
1	B	435	LEU	4.4
1	A	227	VAL	4.3
1	A	44	ARG	4.2
1	A	224	HIS	4.2
1	B	417	ALA	4.2
1	A	435	LEU	4.2
1	A	291	LEU	3.7
1	B	99	LEU	3.6
1	B	151	THR	3.5
1	B	120	TYR	3.4
1	B	236	ALA	3.4
1	B	182	LEU	3.2
1	B	171	TYR	3.2
1	A	232	ALA	3.1
1	B	232	ALA	3.1
1	A	225	MET	3.0
1	B	239	ALA	3.0
1	B	113	LEU	3.0
1	B	100	SER	3.0
1	B	294	HIS	2.8
1	B	365	LYS	2.8
1	A	159	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	95	MET	2.6
1	A	183	ASP	2.6
1	A	184	PRO	2.6
1	B	244	ARG	2.6
1	B	357	GLY	2.6
1	B	297	ARG	2.5
1	A	241	PRO	2.5
1	A	431	LEU	2.5
1	A	42	HIS	2.5
1	A	229	ILE	2.4
1	A	151	THR	2.4
1	A	222	PRO	2.4
1	A	228	MET	2.4
1	A	94	VAL	2.4
1	A	289	GLU	2.3
1	B	191	CYS	2.3
1	B	359	PRO	2.3
1	B	235	PRO	2.3
1	B	233	GLY	2.3
1	B	178	GLU	2.3
1	B	338	GLU	2.3
1	A	226	ARG	2.3
1	B	86	PHE	2.3
1	A	41	ALA	2.2
1	A	234	ALA	2.2
1	A	38	TYR	2.2
1	B	368	LEU	2.2
1	A	239	ALA	2.1
1	B	187	TYR	2.1
1	B	104	LEU	2.1
1	A	283	VAL	2.1
1	B	101	ALA	2.1
1	A	179	HIS	2.1
1	A	29	GLU	2.1
1	B	91	ASP	2.1
1	A	101	ALA	2.1
1	B	231	GLY	2.0
1	A	242	ALA	2.0
1	B	291	LEU	2.0
1	A	39	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.