



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:47 AM UTC

PDB ID : 2REE / pdb_00002ree
Title : Crystal structure of the loading GNATL domain of CurA from *Lyngbya majuscula*
Authors : Geders, T.W.; Smith, J.L.
Deposited on : 2007-09-26
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	8% 79% 9% 11%
1	B	224	7% 78% 10% 11%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CurA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	199	Total	C	N	O	S	0	17	0
			1731	1107	295	319	10			
1	B	199	Total	C	N	O	S	0	13	0
			1701	1085	290	317	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	216	SER	-	expression tag	UNP Q6DNF2
A	217	ASN	-	expression tag	UNP Q6DNF2
A	218	ALA	-	expression tag	UNP Q6DNF2
B	216	SER	-	expression tag	UNP Q6DNF2
B	217	ASN	-	expression tag	UNP Q6DNF2
B	218	ALA	-	expression tag	UNP Q6DNF2

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 10 8 2	0	1
2	B	1	Total O S 10 8 2	0	1

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	243	Total O 245 245	0	2
5	B	214	Total O 215 215	0	1

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CurA

Chain A: 



• Molecule 1: CurA

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	91.46Å 91.46Å 138.88Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.40 – 1.95 43.40 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.6 (43.40-1.95) 99.6 (43.40-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.54 (at 1.95Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.172 , 0.211 0.221 , 0.253	Depositor DCC
R_{free} test set	2501 reflections (4.29%)	wwPDB-VP
Wilson B-factor (Å ²)	31.4	Xtriage
Anisotropy	0.465	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3925	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/1795	0.76	0/2432
1	B	0.65	0/1756	0.80	0/2385
All	All	0.68	0/3551	0.78	0/4817

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1731	0	1762	16	0
1	B	1701	0	1715	22	0
2	A	10	0	0	0	0
2	B	10	0	0	1	0
3	A	1	0	0	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	A	245	0	0	6	0
5	B	215	0	0	3	0
All	All	3925	0	3493	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:376[B]:ASN:OD1	1:B:378[B]:SER:HB2	1.51	1.11
1:B:411:GLN:NE2	5:B:793:HOH:O	1.86	1.06
1:B:269[B]:GLN:HE21	1:B:269[B]:GLN:H	1.24	0.86
1:B:378[B]:SER:HB3	1:B:380:LEU:HD22	1.69	0.74
1:B:269[B]:GLN:H	1:B:269[B]:GLN:NE2	1.85	0.74
1:B:377[A]:GLU:H	1:B:377[A]:GLU:CD	1.97	0.70
1:B:376[B]:ASN:OD1	1:B:378[B]:SER:CB	2.38	0.67
1:A:399:LEU:HB3	1:A:413[B]:CYS:SG	2.37	0.65
1:A:375[A]:LYS:O	5:A:889:HOH:O	2.15	0.64
1:B:376[B]:ASN:CG	1:B:378[B]:SER:HB2	2.23	0.63
1:B:368:MET:HG3	1:B:399:LEU:HD21	1.84	0.59
1:A:308[A]:THR:HG21	5:A:791:HOH:O	2.02	0.59
1:B:326:GLN:HG2	2:B:502[A]:SO4:O3	2.03	0.59
1:B:324:GLU:HG3	1:B:325:LEU:HD12	1.86	0.57
1:B:378[B]:SER:CB	5:B:747:HOH:O	2.52	0.57
1:A:343:ALA:HA	1:A:421:ILE:HD12	1.87	0.56
1:A:358[A]:ARG:HH21	1:A:385[A]:LEU:CD2	2.18	0.56
1:A:308[A]:THR:HG22	5:A:916:HOH:O	2.07	0.54
1:B:376[B]:ASN:C	1:B:378[B]:SER:H	2.14	0.54
1:A:358[A]:ARG:HH21	1:A:385[A]:LEU:HD23	1.74	0.53
1:B:250:SER:OG	1:B:252[B]:ASN:ND2	2.43	0.51
1:B:378[B]:SER:HB3	5:B:747:HOH:O	2.11	0.51
1:A:350[B]:LYS:NZ	5:A:725:HOH:O	2.45	0.49
1:A:377[A]:GLU:CD	1:A:377[A]:GLU:H	2.21	0.49
1:B:376[B]:ASN:C	1:B:378[B]:SER:N	2.69	0.49
1:A:409:GLU:OE1	5:A:915:HOH:O	2.20	0.47
1:B:378[B]:SER:HB2	1:B:380:LEU:HB2	1.96	0.47
1:A:375[A]:LYS:NZ	5:A:774:HOH:O	2.48	0.46
1:A:368[A]:MET:HG2	1:A:399:LEU:HD21	1.98	0.45
1:B:252[B]:ASN:HD22	1:B:252[B]:ASN:H	1.64	0.44
1:A:378[B]:SER:HB3	1:A:380:LEU:HD22	1.99	0.44
1:B:230:LEU:HD11	1:B:272:PHE:HB3	2.00	0.43
1:A:388:PHE:O	1:A:392:HIS:HD2	2.03	0.42
1:A:329:GLY:O	1:A:333:ARG:HD3	2.19	0.42
1:A:399:LEU:HD22	1:A:413[B]:CYS:SG	2.59	0.41
1:B:376[B]:ASN:OD1	1:B:378[B]:SER:N	2.52	0.41
1:B:239:ARG:HD2	1:B:239:ARG:HA	1.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/224 (96%)	210 (98%)	4 (2%)	0	100	100
1	B	210/224 (94%)	204 (97%)	6 (3%)	0	100	100
All	All	424/448 (95%)	414 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/204 (97%)	191 (96%)	7 (4%)	32	22
1	B	194/204 (95%)	184 (95%)	10 (5%)	21	9
All	All	392/408 (96%)	375 (96%)	17 (4%)	29	15

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	235[A]	ILE
1	A	235[B]	ILE
1	A	269[A]	GLN
1	A	269[B]	GLN
1	A	305	LEU
1	A	339	LEU
1	A	380	LEU
1	B	236[A]	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	236[B]	GLU
1	B	267	ILE
1	B	273	ILE
1	B	316	LEU
1	B	322	LEU
1	B	327	ASN
1	B	339	LEU
1	B	351	VAL
1	B	380	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	328	GLN
1	A	411	GLN
1	B	226	ASN
1	B	411	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	601	-	5,5,5	0.36	0	5,5,5	0.91	0
2	SO4	A	501[A]	-	4,4,4	0.30	0	6,6,6	0.35	0
2	SO4	B	502[A]	-	4,4,4	0.30	0	6,6,6	0.13	0
2	SO4	A	501[B]	-	4,4,4	0.27	0	6,6,6	0.16	0
2	SO4	B	502[B]	-	4,4,4	0.20	0	6,6,6	0.20	0
4	GOL	B	602	-	5,5,5	0.26	0	5,5,5	1.04	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	601	-	-	0/4/4/4	-
4	GOL	B	602	-	-	3/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	602	GOL	C3-C2-C1	-2.01	104.41	111.80

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	602	GOL	C1-C2-C3-O3
4	B	602	GOL	O2-C2-C3-O3
4	B	602	GOL	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	502[A]	SO4	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	199/224 (88%)	0.55	18 (9%) 15 17	9, 24, 39, 51	17 (8%)
1	B	199/224 (88%)	0.60	16 (8%) 18 21	9, 26, 42, 57	13 (6%)
All	All	398/448 (88%)	0.58	34 (8%) 16 19	9, 25, 41, 57	30 (7%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	227	TYR	7.0
1	B	278	ASP	5.8
1	A	377[A]	GLU	4.2
1	B	277	GLU	4.0
1	A	277	GLU	3.9
1	A	276	LEU	3.8
1	B	276	LEU	3.5
1	B	227	TYR	3.4
1	A	226	ASN	3.3
1	A	324	GLU	3.2
1	B	322	LEU	3.1
1	B	327	ASN	3.0
1	A	327	ASN	2.9
1	B	423	HIS	2.9
1	A	230	LEU	2.8
1	A	344	GLN	2.8
1	A	322	LEU	2.8
1	A	423	HIS	2.7
1	A	278	ASP	2.7
1	B	281	VAL	2.7
1	B	324	GLU	2.7
1	B	279	LYS	2.6
1	B	343	ALA	2.6
1	A	325	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	341	TYR	2.5
1	B	346	SER	2.4
1	A	228	TYR	2.4
1	A	273	ILE	2.2
1	A	341	TYR	2.2
1	B	235	ILE	2.1
1	B	309	GLU	2.1
1	A	274	LEU	2.1
1	A	421	ILE	2.0
1	B	226	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	502[A]	5/5	0.80	0.20	43,46,51,51	5
2	SO4	B	502[B]	5/5	0.80	0.20	41,42,47,50	5
2	SO4	A	501[A]	5/5	0.85	0.19	28,34,49,51	5
2	SO4	A	501[B]	5/5	0.85	0.19	40,46,54,55	5
4	GOL	B	602	6/6	0.89	0.16	36,47,54,55	0
4	GOL	A	601	6/6	0.95	0.12	31,34,41,42	0
3	CL	A	701	1/1	0.99	0.06	29,29,29,29	1

6.5 Other polymers [i](#)

There are no such residues in this entry.