



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:39 PM UTC

PDB ID : 2REF / pdb_00002ref
Title : Crystal structure of the loading GNATL domain of CurA from *Lyngbya majuscula* soaked with malonyl-CoA
Authors : Geders, T.W.; Smith, J.L.
Deposited on : 2007-09-26
Resolution : 2.75 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	
1	B	224	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 3510 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CurA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	203	Total 1679	C 1073	N 283	O 312	S 11	0	5	0
1	B	203	Total 1674	C 1069	N 284	O 311	S 10	0	4	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	216	SER	-	expression tag	UNP Q6DNF2
A	217	ASN	-	expression tag	UNP Q6DNF2
A	218	ALA	-	expression tag	UNP Q6DNF2
B	216	SER	-	expression tag	UNP Q6DNF2
B	217	ASN	-	expression tag	UNP Q6DNF2
B	218	ALA	-	expression tag	UNP Q6DNF2

- Molecule 2 is ACETYL COENZYME *A (CCD ID: ACO) (formula: $C_{23}H_{38}N_7O_{17}P_3S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	1
			102	46	14	34	6	2		
2	B	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	O	0	0
			2	2		
3	B	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CurA



- Molecule 1: CurA



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	91.89Å 91.89Å 139.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.50 – 2.75 46.50 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.50-2.75) 99.8 (46.50-2.75)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.44 (at 2.77Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.207 , 0.251 0.237 , 0.268	Depositor DCC
R_{free} test set	915 reflections (4.23%)	wwPDB-VP
Wilson B-factor (Å ²)	47.2	Xtrriage
Anisotropy	0.138	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 7.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.032 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3510	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	1/1729 (0.1%)	0.81	4/2347 (0.2%)
1	B	0.52	0/1721	0.79	0/2338
All	All	0.52	1/3450 (0.0%)	0.80	4/4685 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	303	VAL	CA-CB	5.78	1.57	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	303	VAL	N-CA-CB	7.33	115.12	110.50
1	A	377	GLU	N-CA-C	5.95	123.47	110.80
1	A	377	GLU	CA-C-N	5.62	131.81	121.70
1	A	377	GLU	C-N-CA	5.62	131.81	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1679	0	1683	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1674	0	1676	8	0
2	A	102	0	68	8	0
2	B	51	0	34	4	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
All	All	3510	0	3461	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:318:ALA:HA	2:B:1:ACO:H22	1.53	0.91
1:B:316:LEU:HB3	2:B:1:ACO:HH32	1.55	0.88
2:B:1:ACO:H62	2:B:1:ACO:O9P	1.83	0.77
1:A:318:ALA:HA	2:A:1[A]:ACO:H21	1.74	0.69
1:A:318:ALA:HA	2:A:1[A]:ACO:C2P	2.26	0.65
1:A:316:LEU:CB	2:A:1[A]:ACO:HH32	2.32	0.59
1:A:300:CYS:O	1:A:303:VAL:HG22	2.04	0.57
1:A:399:LEU:HD22	1:A:413[B]:CYS:SG	2.45	0.57
2:A:1[A]:ACO:H131	2:A:1[A]:ACO:HN8	1.69	0.56
1:B:318:ALA:HA	2:B:1:ACO:C2P	2.33	0.55
2:A:1[B]:ACO:H32	2:A:1[B]:ACO:H141	1.90	0.53
1:A:226:ASN:HB3	1:A:276:LEU:HD11	1.89	0.53
1:A:337:PHE:CE1	1:B:222:CYS:HB3	2.46	0.50
1:A:399:LEU:HB3	1:A:413[B]:CYS:SG	2.52	0.50
1:A:332:ASP:OD1	1:A:392:HIS:NE2	2.35	0.48
1:A:316:LEU:HB2	2:A:1[A]:ACO:HH32	1.95	0.47
1:A:316:LEU:HB3	2:A:1[A]:ACO:HH32	1.96	0.47
1:B:394:ALA:HB2	1:B:419:TYR:CZ	2.49	0.47
1:A:284:ILE:HG12	1:A:338:MET:HE2	1.97	0.46
1:A:249:TRP:CZ2	2:A:1[B]:ACO:N4P	2.84	0.46
1:B:233:PRO:HG3	1:B:264:ILE:HG22	1.99	0.45
1:A:303:VAL:HG23	1:A:304:PRO:HD3	2.00	0.44
1:A:226:ASN:HB3	1:A:276:LEU:CD1	2.48	0.43
1:A:394:ALA:HB2	1:A:419:TYR:CZ	2.54	0.42
1:A:282:GLY:HA2	1:A:320:ASN:O	2.20	0.42
1:A:376:ASN:OD1	1:A:378:SER:HB3	2.20	0.42
1:A:284:ILE:HG23	1:A:338:MET:HE2	2.02	0.42
1:B:368:MET:HG3	1:B:399:LEU:HD21	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:303:VAL:N	1:B:304:PRO:CD	2.83	0.41
1:A:223:PHE:O	1:A:224:GLU:C	2.64	0.41
1:A:233:PRO:HB3	1:A:273:ILE:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/224 (92%)	197 (96%)	7 (3%)	2 (1%)	12	24
1	B	205/224 (92%)	199 (97%)	6 (3%)	0	100	100
All	All	411/448 (92%)	396 (96%)	13 (3%)	2 (0%)	24	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	GLU
1	A	378	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/204 (94%)	178 (93%)	13 (7%)	14	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	190/204 (93%)	181 (95%)	9 (5%)	23	45
All	All	381/408 (93%)	359 (94%)	22 (6%)	18	34

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	224	GLU
1	A	235	ILE
1	A	236	GLU
1	A	267	ILE
1	A	269	GLN
1	A	279	LYS
1	A	305	LEU
1	A	339	LEU
1	A	340	GLN
1	A	344	GLN
1	A	375	LYS
1	A	377	GLU
1	A	380	LEU
1	B	224	GLU
1	B	267	ILE
1	B	303	VAL
1	B	305	LEU
1	B	316	LEU
1	B	339	LEU
1	B	351	VAL
1	B	380	LEU
1	B	421	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	293	GLN
1	A	327	ASN
1	B	327	ASN
1	B	411	GLN
1	B	423	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACO	A	1[B]	-	51,53,53	1.20	5 (9%)	73,79,79	1.50	10 (13%)
2	ACO	B	1	-	51,53,53	1.19	3 (5%)	73,79,79	1.63	13 (17%)
2	ACO	A	1[A]	-	51,53,53	1.21	4 (7%)	73,79,79	1.50	9 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACO	A	1[B]	-	-	10/51/67/67	0/3/3/3
2	ACO	B	1	-	-	17/51/67/67	0/3/3/3
2	ACO	A	1[A]	-	-	22/51/67/67	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	ACO	C5A-C4A	4.80	1.47	1.39
2	A	1[A]	ACO	C5A-C4A	4.74	1.47	1.39
2	A	1[B]	ACO	C5A-C4A	4.74	1.47	1.39
2	B	1	ACO	C5A-C6A	3.14	1.49	1.41
2	A	1[B]	ACO	C5A-C6A	2.96	1.49	1.41
2	A	1[A]	ACO	C5A-C6A	2.93	1.49	1.41
2	B	1	ACO	C8A-N7A	2.59	1.36	1.31
2	A	1[A]	ACO	C8A-N7A	2.57	1.36	1.31
2	A	1[B]	ACO	C8A-N7A	2.56	1.36	1.31
2	A	1[A]	ACO	P1A-O3A	2.12	1.61	1.59
2	A	1[B]	ACO	P2A-O3A	2.07	1.61	1.59
2	A	1[B]	ACO	C5A-N7A	-2.02	1.35	1.39

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	ACO	C5A-C4A-N3A	-5.58	119.04	126.72
2	A	1[A]	ACO	C5A-C4A-N3A	-5.51	119.13	126.72
2	A	1[B]	ACO	C5A-C4A-N3A	-5.44	119.22	126.72
2	A	1[A]	ACO	N3A-C4A-N9A	4.33	134.54	127.17
2	B	1	ACO	N3A-C4A-N9A	4.31	134.50	127.17
2	A	1[B]	ACO	N3A-C4A-N9A	4.22	134.35	127.17
2	B	1	ACO	C4A-C5A-N7A	-3.90	106.12	110.58
2	B	1	ACO	C2A-N3A-C4A	3.84	121.22	111.83
2	A	1[B]	ACO	C2A-N3A-C4A	3.84	121.22	111.83
2	A	1[A]	ACO	C2A-N3A-C4A	3.82	121.15	111.83
2	A	1[B]	ACO	N3A-C2A-N1A	-3.77	122.87	128.58
2	A	1[A]	ACO	N3A-C2A-N1A	-3.67	123.03	128.58
2	B	1	ACO	N3A-C2A-N1A	-3.62	123.10	128.58
2	A	1[B]	ACO	C4A-C5A-N7A	-3.58	106.49	110.58
2	A	1[A]	ACO	C4A-C5A-N7A	-3.53	106.54	110.58
2	B	1	ACO	C4A-N9A-C8A	3.14	109.03	105.74
2	B	1	ACO	C5A-N7A-C8A	2.94	108.06	103.45
2	A	1[A]	ACO	C4A-N9A-C8A	2.87	108.75	105.74
2	A	1[B]	ACO	C4A-N9A-C8A	2.84	108.72	105.74
2	B	1	ACO	CEP-CBP-CAP	2.74	113.43	108.77
2	B	1	ACO	C6A-C5A-N7A	2.67	137.25	132.09
2	A	1[B]	ACO	C5A-N7A-C8A	2.66	107.64	103.45
2	A	1[A]	ACO	C5A-N7A-C8A	2.63	107.59	103.45
2	B	1	ACO	N9A-C8A-N7A	-2.55	110.32	113.94
2	A	1[B]	ACO	C6A-C5A-N7A	2.55	137.00	132.09
2	A	1[A]	ACO	C6A-C5A-N7A	2.44	136.79	132.09
2	B	1	ACO	O4B-C1B-N9A	2.37	112.63	108.09

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1[B]	ACO	N9A-C8A-N7A	-2.26	110.73	113.94
2	A	1[A]	ACO	N9A-C8A-N7A	-2.22	110.78	113.94
2	B	1	ACO	O2A-P1A-O1A	2.06	122.03	112.44
2	B	1	ACO	C6P-C7P-N8P	2.04	116.33	112.00
2	A	1[B]	ACO	C2A-N1A-C6A	2.02	122.05	118.73

There are no chirality outliers.

All (49) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1[A]	ACO	CEP-CBP-CCP-O6A
2	A	1[A]	ACO	CAP-CBP-CCP-O6A
2	A	1[A]	ACO	O9P-C9P-CAP-CBP
2	A	1[A]	ACO	N8P-C9P-CAP-CBP
2	A	1[A]	ACO	N8P-C9P-CAP-OAP
2	A	1[A]	ACO	CAP-C9P-N8P-C7P
2	A	1[A]	ACO	C6P-C5P-N4P-C3P
2	A	1[A]	ACO	S1P-C2P-C3P-N4P
2	A	1[A]	ACO	C3P-C2P-S1P-C
2	A	1[A]	ACO	O-C-S1P-C2P
2	A	1[A]	ACO	CH3-C-S1P-C2P
2	A	1[B]	ACO	C5P-C6P-C7P-N8P
2	A	1[B]	ACO	S1P-C2P-C3P-N4P
2	A	1[B]	ACO	O-C-S1P-C2P
2	B	1	ACO	OAP-CAP-CBP-CCP
2	B	1	ACO	OAP-CAP-CBP-CEP
2	B	1	ACO	O9P-C9P-CAP-CBP
2	B	1	ACO	N8P-C9P-CAP-CBP
2	B	1	ACO	C5P-C6P-C7P-N8P
2	B	1	ACO	O-C-S1P-C2P
2	B	1	ACO	CH3-C-S1P-C2P
2	B	1	ACO	C6P-C7P-N8P-C9P
2	A	1[A]	ACO	O5P-C5P-N4P-C3P
2	A	1[B]	ACO	CH3-C-S1P-C2P
2	B	1	ACO	OAP-CAP-CBP-CDP
2	A	1[A]	ACO	O9P-C9P-N8P-C7P
2	A	1[A]	ACO	C2B-C1B-N9A-C8A
2	B	1	ACO	N8P-C9P-CAP-OAP
2	A	1[B]	ACO	C2B-C1B-N9A-C8A
2	A	1[A]	ACO	N4P-C5P-C6P-C7P
2	B	1	ACO	C2B-C1B-N9A-C8A
2	B	1	ACO	P2A-O3A-P1A-O2A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	1[A]	ACO	O5P-C5P-C6P-C7P
2	A	1[A]	ACO	CDP-CBP-CCP-O6A
2	A	1[B]	ACO	C3P-C2P-S1P-C
2	A	1[A]	ACO	P2A-O3A-P1A-O2A
2	A	1[B]	ACO	P2A-O3A-P1A-O2A
2	B	1	ACO	P2A-O3A-P1A-O1A
2	A	1[A]	ACO	C2B-C1B-N9A-C4A
2	A	1[B]	ACO	C2B-C1B-N9A-C4A
2	B	1	ACO	C2B-C1B-N9A-C4A
2	A	1[A]	ACO	C2P-C3P-N4P-C5P
2	B	1	ACO	N4P-C5P-C6P-C7P
2	B	1	ACO	O4B-C1B-N9A-C8A
2	B	1	ACO	O5P-C5P-C6P-C7P
2	A	1[A]	ACO	O4B-C1B-N9A-C8A
2	A	1[B]	ACO	O4B-C1B-N9A-C8A
2	A	1[A]	ACO	P2A-O3A-P1A-O1A
2	A	1[B]	ACO	P2A-O3A-P1A-O1A

There are no ring outliers.

3 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1[B]	ACO	2	0
2	B	1	ACO	4	0
2	A	1[A]	ACO	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/224 (90%)	0.58	18 (8%) 15 15	22, 38, 42, 46	5 (2%)
1	B	203/224 (90%)	0.62	14 (6%) 23 22	23, 38, 42, 46	4 (1%)
All	All	406/448 (90%)	0.60	32 (7%) 18 19	22, 38, 42, 46	9 (2%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	278	ASP	4.8
1	A	278	ASP	4.5
1	A	227	TYR	4.4
1	B	423	HIS	4.1
1	A	222	CYS	3.9
1	A	223	PHE	3.6
1	A	277	GLU	3.6
1	A	423	HIS	3.5
1	B	225	ASN	3.4
1	A	225	ASN	3.4
1	B	422	GLN	3.2
1	B	377	GLU	3.1
1	B	223	PHE	3.1
1	B	227	TYR	3.1
1	A	344	GLN	3.0
1	A	422	GLN	3.0
1	B	324	GLU	2.9
1	B	222	CYS	2.8
1	B	224	GLU	2.8
1	A	377	GLU	2.7
1	B	309	GLU	2.7
1	A	324	GLU	2.6
1	A	224	GLU	2.4
1	A	226	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	347	GLY	2.4
1	A	340	GLN	2.3
1	B	332	ASP	2.3
1	A	342	CYS	2.2
1	A	424	ARG	2.1
1	A	332	ASP	2.1
1	B	349	GLU	2.1
1	B	277	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACO	A	1[A]	51/51	0.89	0.15	30,36,50,51	51
2	ACO	A	1[B]	51/51	0.89	0.15	57,60,68,68	51
2	ACO	B	1	51/51	0.91	0.12	43,51,60,61	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around ACO A 1 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ACO A 1 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.