



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:42 AM UTC

PDB ID : 3REK / pdb_00003rek
Title : 2.6 Angstrom Crystal Structure of the Nucleosome Core Particle Assembled with a 146 bp Alpha-Satellite DNA (NCP146b) Derivatized with Oxaliplatin
Authors : Wu, B.; Davey, C.A.
Deposited on : 2011-04-04
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	2% 61% 10% 27%
1	E	135	7% 64% 8% 28%
2	B	102	2% 64% 17% 19%
2	F	102	6% 71% 5% 24%
3	C	129	12% 59% 18% 20%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	129	
4	D	122	
4	H	122	
5	I	146	
5	J	146	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	PT	J	87	-	-	-	X
8	SO4	D	1101	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12052 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	98	Total	C	N	O	S	0	0	0
			808	509	156	140	3			
1	E	97	Total	C	N	O	S	0	0	0
			801	504	155	139	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	GLY	variant	UNP P84233
E	102	ALA	GLY	variant	UNP P84233

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	83	Total	C	N	O	S	0	0	0
			662	418	129	114	1			
2	F	78	Total	C	N	O	S	0	0	0
			619	391	120	107	1			

- Molecule 3 is a protein called Histone H2A type1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	103	Total	C	N	O	0	0	0
			795	501	155	139			
3	G	105	Total	C	N	O	0	0	0
			809	510	158	141			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	99	ARG	GLY	variant	UNP P06897
C	123	SER	ALA	variant	UNP P06897

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	99	ARG	GLY	variant	UNP P06897
G	123	SER	ALA	variant	UNP P06897

- Molecule 4 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	99	Total	C	N	O	S	0	0	0
			785	493	146	144	2			
4	H	94	Total	C	N	O	S	0	0	0
			736	463	132	139	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	29	THR	SER	variant	UNP P02281
H	29	THR	SER	variant	UNP P02281

- Molecule 5 is a DNA chain called DNA (146-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	146	Total	C	N	O	P	0	0	0
			2990	1430	541	874	145			
5	J	146	Total	C	N	O	P	0	0	0
			2990	1430	541	874	145			

- Molecule 6 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mn	0	0
			1	1		
6	D	1	Total	Mn	0	0
			1	1		

- Molecule 7 is PLATINUM (II) ION (CCD ID: PT) (formula: Pt).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Pt	0	0
			1	1		
7	B	2	Total	Pt	0	0
			2	2		

Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	E	1	Total	Pt	0	0
			1	1		
7	F	1	Total	Pt	0	0
			1	1		
7	I	18	Total	Pt	0	0
			18	18		
7	J	17	Total	Pt	0	0
			17	17		


- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	C	1	Total	O	S	0	0
			5	4	1		
8	D	1	Total	O	S	0	0
			5	4	1		
8	G	1	Total	O	S	0	0
			5	4	1		

- Molecule 1: Histone H3.2







• Molecule 3: Histone H2A type1



• Molecule 4: Histone H2B 1.1




• Molecule 4: Histone H2B 1.1





• Molecule 5: DNA (146-MER)





• Molecule 5: DNA (146-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	105.00Å 109.44Å 176.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.00 – 2.60 57.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	93.1 (57.00-2.60) 93.1 (57.00-2.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.260 , 0.281 0.249 , 0.270	Depositor DCC
R_{free} test set	1210 reflections (2.06%)	wwPDB-VP
Wilson B-factor (Å ²)	63.0	Xtriage
Anisotropy	0.485	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12052	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PT, SO4, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	0/820	0.93	1/1099 (0.1%)
1	E	0.59	0/812	0.82	0/1088
2	B	0.83	0/669	1.05	1/894 (0.1%)
2	F	0.65	0/626	0.88	0/837
3	C	0.62	0/805	0.91	0/1088
3	G	0.80	0/819	0.92	0/1106
4	D	0.70	0/796	0.95	0/1065
4	H	0.76	0/747	0.97	0/1004
5	I	0.34	0/3354	0.96	1/5175 (0.0%)
5	J	0.31	0/3354	0.95	0/5175
All	All	0.55	0/12802	0.94	3/18531 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	VAL	N-CA-C	-5.57	107.69	113.10
5	I	53	DC	P-O3'-C3'	5.36	128.23	120.20
2	B	30	THR	N-CA-C	5.06	117.38	110.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	808	0	846	13	0
1	E	801	0	838	6	0
2	B	662	0	708	17	0
2	F	619	0	659	7	0
3	C	795	0	846	23	0
3	G	809	0	864	36	0
4	D	785	0	825	32	0
4	H	736	0	760	23	0
5	I	2990	0	1651	29	0
5	J	2990	0	1651	39	0
6	A	1	0	0	0	0
6	D	1	0	0	0	0
7	A	1	0	0	0	0
7	B	2	0	0	0	0
7	E	1	0	0	0	0
7	F	1	0	0	0	0
7	I	18	0	0	0	0
7	J	17	0	0	1	0
8	C	5	0	0	0	0
8	D	5	0	0	2	0
8	G	5	0	0	0	0
All	All	12052	0	9648	170	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 170 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:75:HIS:CD2	4:H:93:THR:HG21	1.81	1.13
2:F:75:HIS:HD2	4:H:93:THR:HG21	1.15	1.08
4:D:79:HIS:CE1	3:G:38:ASN:HD22	1.80	1.00
4:D:79:HIS:HE1	3:G:38:ASN:HD22	0.93	0.92
5:J:67:DG:H2''	5:J:68:DG:H5''	1.50	0.90

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	96/135 (71%)	94 (98%)	2 (2%)	0	100	100
1	E	95/135 (70%)	93 (98%)	2 (2%)	0	100	100
2	B	81/102 (79%)	79 (98%)	2 (2%)	0	100	100
2	F	76/102 (74%)	75 (99%)	1 (1%)	0	100	100
3	C	101/129 (78%)	97 (96%)	4 (4%)	0	100	100
3	G	103/129 (80%)	99 (96%)	4 (4%)	0	100	100
4	D	97/122 (80%)	89 (92%)	5 (5%)	3 (3%)	3	5
4	H	92/122 (75%)	89 (97%)	2 (2%)	1 (1%)	11	25
All	All	741/976 (76%)	715 (96%)	22 (3%)	4 (0%)	24	46

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	101	GLY
4	H	101	GLY
4	D	30	ARG
4	D	44	GLN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/110 (77%)	79 (93%)	6 (7%)	13	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	84/110 (76%)	79 (94%)	5 (6%)	17	37
2	B	68/78 (87%)	67 (98%)	1 (2%)	57	80
2	F	63/78 (81%)	62 (98%)	1 (2%)	55	79
3	C	82/101 (81%)	75 (92%)	7 (8%)	10	22
3	G	83/101 (82%)	77 (93%)	6 (7%)	13	30
4	D	85/102 (83%)	79 (93%)	6 (7%)	13	31
4	H	80/102 (78%)	73 (91%)	7 (9%)	9	21
All	All	630/782 (81%)	591 (94%)	39 (6%)	16	36

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	76	THR
4	H	85	THR
3	G	81	ARG
4	H	33	SER
4	H	103	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
4	H	60	ASN
4	H	46	HIS
2	F	75	HIS
3	G	104	GLN
1	E	68	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 45 ligands modelled in this entry, 42 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	SO4	G	1103	-	4,4,4	0.31	0	6,6,6	0.57	0
8	SO4	D	1101	-	4,4,4	0.43	0	6,6,6	0.64	0
8	SO4	C	1102	-	4,4,4	0.34	0	6,6,6	0.46	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	D	1101	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	98/135 (72%)	0.19	3 (3%) 51 45	38, 52, 72, 90	0
1	E	97/135 (71%)	0.75	9 (9%) 14 11	50, 70, 91, 101	0
2	B	83/102 (81%)	0.28	2 (2%) 59 54	39, 49, 59, 62	0
2	F	78/102 (76%)	0.67	6 (7%) 19 15	52, 63, 75, 80	0
3	C	103/129 (79%)	0.81	15 (14%) 6 4	54, 68, 87, 93	0
3	G	105/129 (81%)	0.17	4 (3%) 44 38	36, 54, 74, 84	0
4	D	99/122 (81%)	0.83	14 (14%) 6 5	51, 66, 115, 130	0
4	H	94/122 (77%)	0.33	2 (2%) 63 58	43, 55, 81, 94	0
5	I	146/146 (100%)	0.63	3 (2%) 63 58	70, 124, 159, 167	0
5	J	146/146 (100%)	0.60	4 (2%) 56 50	64, 120, 148, 166	0
All	All	1049/1268 (82%)	0.54	62 (5%) 28 23	36, 67, 140, 167	0

The worst 5 of 62 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	67	PHE	5.7
2	F	29	ILE	5.0
2	B	102	GLY	4.6
1	E	135	ALA	4.6
2	F	102	GLY	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	PT	J	86	1/1	-0.32	0.24	338,338,338,338	1
7	PT	I	90	1/1	-0.05	0.26	339,339,339,339	1
7	PT	J	87	1/1	0.14	0.56	94,94,94,94	1
7	PT	F	103	1/1	0.19	0.29	152,152,152,152	1
7	PT	J	88	1/1	0.27	0.17	153,153,153,153	1
7	PT	J	82	1/1	0.28	0.24	162,162,162,162	1
7	PT	I	89	1/1	0.32	0.19	155,155,155,155	1
7	PT	I	78	1/1	0.33	0.33	168,168,168,168	1
7	PT	J	89	1/1	0.33	0.23	142,142,142,142	1
7	PT	I	80	1/1	0.34	0.18	202,202,202,202	1
7	PT	J	79	1/1	0.35	0.17	139,139,139,139	1
7	PT	J	85	1/1	0.37	0.23	99,99,99,99	1
7	PT	J	83	1/1	0.40	0.13	227,227,227,227	1
7	PT	I	81	1/1	0.40	0.15	136,136,136,136	1
7	PT	I	88	1/1	0.41	0.21	176,176,176,176	1
7	PT	J	80	1/1	0.42	0.18	297,297,297,297	1
7	PT	I	84	1/1	0.44	0.20	197,197,197,197	1
7	PT	J	81	1/1	0.47	0.18	195,195,195,195	1
7	PT	I	101	1/1	0.48	0.29	194,194,194,194	1
7	PT	B	104	1/1	0.50	0.22	256,256,256,256	1
7	PT	I	85	1/1	0.51	0.17	192,192,192,192	1
7	PT	I	87	1/1	0.57	0.11	122,122,122,122	1
7	PT	I	79	1/1	0.61	0.12	140,140,140,140	1
7	PT	I	83	1/1	0.66	0.11	163,163,163,163	1
7	PT	I	77	1/1	0.66	0.20	169,169,169,169	1
7	PT	I	82	1/1	0.66	0.11	137,137,137,137	1
7	PT	I	75	1/1	0.70	0.15	182,182,182,182	1
7	PT	J	77	1/1	0.70	0.10	187,187,187,187	1
7	PT	B	103	1/1	0.70	0.14	155,155,155,155	1
7	PT	I	86	1/1	0.74	0.09	255,255,255,255	1
8	SO4	D	1101	5/5	0.74	0.11	83,83,84,85	0
7	PT	A	136	1/1	0.76	0.24	384,384,384,384	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	PT	I	76	1/1	0.78	0.19	185,185,185,185	1
7	PT	E	136	1/1	0.80	0.11	122,122,122,122	1
7	PT	J	75	1/1	0.81	0.18	147,147,147,147	1
7	PT	J	76	1/1	0.81	0.19	129,129,129,129	1
7	PT	J	78	1/1	0.82	0.16	71,71,71,71	1
7	PT	J	84	1/1	0.83	0.11	85,85,85,85	1
6	MN	A	1001	1/1	0.88	0.18	53,53,53,53	0
6	MN	D	1007	1/1	0.92	0.23	110,110,110,110	0
7	PT	J	73	1/1	0.92	0.07	119,119,119,119	1
7	PT	I	74	1/1	0.93	0.08	149,149,149,149	0
7	PT	J	74	1/1	0.93	0.07	144,144,144,144	1
8	SO4	G	1103	5/5	0.93	0.11	69,69,70,71	0
8	SO4	C	1102	5/5	0.97	0.09	61,63,65,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.