



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:17 AM UTC

PDB ID : 2RFI / pdb_00002rfi
Title : Crystal structure of catalytic domain of human euchromatic histone methyltransferase 1 in complex with SAH and dimethylated H3K9 peptide
Authors : Min, J.; Wu, H.; Loppnau, P.; Weigelt, J.; Sundstrom, M.; Arrowsmith, C.H.; Edwards, A.M.; Bochkarev, A.; Plotnikov, A.N.; Structural Genomics Consortium (SGC)
Deposited on : 2007-09-30
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	4% 84% 7% 8%
1	B	285	18% 82% 7% 12%
2	P	11	9% 64% 9% 27%
2	Q	11	27% 27% 45%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase, H3 lysine-9 specific 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	261	Total	C	N	O	S	0	5	0
			2147	1334	390	397	26			
1	B	252	Total	C	N	O	S	0	3	0
			2063	1288	376	373	26			


- Molecule 2 is a protein called Histone H3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	P	8	Total	C	N	O	0	0	0
			65	39	14	12			
2	Q	6	Total	C	N	O	0	0	0
			47	28	10	9			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Zn	0	0
			4	4		
3	B	4	Total	Zn	0	0
			4	4		

- Molecule 4 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 26	C 14	N 6	O 5	S 1	0	0
4	A	1	Total 26	C 14	N 6	O 5	S 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	390	Total O 390 390	0	0
5	B	291	Total O 291 291	0	0
5	P	4	Total O 4 4	0	0
5	Q	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone-lysine N-methyltransferase, H3 lysine-9 specific 5




- Molecule 1: Histone-lysine N-methyltransferase, H3 lysine-9 specific 5




- Molecule 2: Histone H3



- Molecule 2: Histone H3





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	84.59Å 85.63Å 95.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.80 – 1.59 42.80 – 1.59	Depositor EDS
% Data completeness (in resolution range)	96.3 (42.80-1.59) 96.3 (42.80-1.59)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 1.59Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.193 , 0.220 0.192 , 0.218	Depositor DCC
R_{free} test set	4475 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5068	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MLY, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/2198	0.72	0/2973
1	B	0.51	0/2113	0.71	0/2857
2	P	2.21	1/53 (1.9%)	0.93	0/69
2	Q	1.00	0/35	1.32	2/46 (4.3%)
All	All	0.58	1/4399 (0.0%)	0.73	2/5945 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	P	7	MLY	C-N	15.79	1.55	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Q	8	SER	CA-C-N	5.15	130.97	121.70
2	Q	8	SER	C-N-CA	5.15	130.97	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2147	0	2031	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2063	0	1960	17	0
2	P	65	0	73	0	0
2	Q	47	0	52	4	0
3	A	4	0	0	0	0
3	B	4	0	0	0	0
4	A	52	0	38	3	0
5	A	390	0	0	1	0
5	B	291	0	0	6	0
5	P	4	0	0	0	0
5	Q	1	0	0	0	0
All	All	5068	0	4154	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1141:SER:O	2:Q:7:MLY:HH12	1.76	0.86
1:A:1170:HIS:HD2	1:A:1211:TYR:H	1.21	0.84
1:B:1171:HIS:HD2	1:B:1173:GLU:H	1.30	0.80
1:A:1169:ASN:HD21	4:A:101:SAH:HN1	1.32	0.77
1:B:1170:HIS:HD2	1:B:1211:TYR:H	1.31	0.76
4:A:102:SAH:HN1	1:B:1169:ASN:HD21	1.31	0.76
1:B:990:ILE:CD1	5:B:1332:HOH:O	2.48	0.62
1:A:985[A]:ARG:HG3	5:A:1372:HOH:O	2.00	0.61
1:A:1170:HIS:CD2	1:A:1211:TYR:H	2.12	0.61
1:B:1170:HIS:CD2	1:B:1211:TYR:H	2.15	0.61
5:B:1322:HOH:O	2:Q:9:THR:HG22	1.99	0.60
5:B:1470:HOH:O	2:Q:9:THR:HG23	2.04	0.57
1:B:1128:LEU:HD21	1:B:1190:PHE:CD1	2.40	0.56
1:B:1170:HIS:CD2	1:B:1216:TRP:HE1	2.23	0.56
1:B:990:ILE:HD11	5:B:1332:HOH:O	2.06	0.56
1:B:1142:TYR:HA	2:Q:7:MLY:HH12	1.90	0.53
1:A:1087:ARG:HH11	1:A:1090:GLN:HE22	1.57	0.52
1:A:1128:LEU:HD23	1:A:1156:ILE:HG12	1.92	0.51
1:B:1087:ARG:HH21	1:B:1090:GLN:HE22	1.61	0.49
1:A:990:ILE:HD11	1:A:1095:ALA:O	2.13	0.48
1:B:1171:HIS:CD2	1:B:1173:GLU:H	2.21	0.47
1:A:989:ARG:HH12	1:A:1094:ARG:HH21	1.61	0.47
1:A:1093:LEU:HD23	1:A:1121:VAL:O	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1170:HIS:HE1	1:A:1172:CYS:SG	2.39	0.45
1:A:1096:ARG:H	1:A:1113[B]:GLN:NE2	2.15	0.44
1:B:1093:LEU:HD23	1:B:1093:LEU:C	2.43	0.44
1:B:1128:LEU:HD21	1:B:1190:PHE:CE1	2.53	0.43
1:B:989:ARG:NH2	5:B:1440:HOH:O	2.52	0.43
1:B:1170:HIS:HE1	1:B:1172:CYS:SG	2.42	0.43
1:B:1214:ARG:NH2	5:B:1508:HOH:O	2.53	0.42
1:A:1214:ARG:HA	1:A:1214:ARG:HD3	1.84	0.41
1:A:1215:PHE:CZ	1:A:1219:LYS:HG3	2.56	0.41
1:A:1169:ASN:HD22	4:A:101:SAH:H8	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/285 (93%)	254 (96%)	10 (4%)	0	100	100
1	B	251/285 (88%)	240 (96%)	11 (4%)	0	100	100
2	P	5/11 (46%)	5 (100%)	0	0	100	100
2	Q	3/11 (27%)	3 (100%)	0	0	100	100
All	All	523/592 (88%)	502 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/257 (94%)	238 (99%)	3 (1%)	63	43
1	B	230/257 (90%)	228 (99%)	2 (1%)	70	55
2	P	6/7 (86%)	6 (100%)	0	100	100
2	Q	4/7 (57%)	4 (100%)	0	100	100
All	All	481/528 (91%)	476 (99%)	5 (1%)	65	50

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1065	MET
1	A	1093	LEU
1	A	1135	ASP
1	B	998	VAL
1	B	1055	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1046	GLN
1	A	1090	GLN
1	A	1169	ASN
1	A	1170	HIS
1	A	1206	GLN
1	B	1090	GLN
1	B	1169	ASN
1	B	1170	HIS
1	B	1171	HIS
1	B	1206	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLY	P	7	2	9,10,11	0.57	0	6,11,13	0.75	0
2	MLY	Q	7	2	9,10,11	0.71	0	6,11,13	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLY	P	7	2	-	0/8/9/11	-
2	MLY	Q	7	2	-	5/8/9/11	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Q	7	MLY	CD-CE-NZ-CH1
2	Q	7	MLY	CD-CE-NZ-CH2
2	Q	7	MLY	CA-CB-CG-CD
2	Q	7	MLY	CG-CD-CE-NZ
2	Q	7	MLY	CE-CD-CG-CB

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Q	7	MLY	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 8 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	SAH	A	101	-	27,28,28	1.06	4 (14%)	36,40,40	1.88	11 (30%)
4	SAH	A	102	-	27,28,28	1.12	4 (14%)	36,40,40	1.96	9 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAH	A	101	-	-	4/15/31/31	0/3/3/3
4	SAH	A	102	-	-	3/15/31/31	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	102	SAH	C2-N1	2.79	1.38	1.33
4	A	102	SAH	C2-N3	2.65	1.38	1.33
4	A	101	SAH	C2-N3	2.48	1.38	1.33
4	A	101	SAH	C2-N1	2.23	1.37	1.33
4	A	102	SAH	OXT-C	-2.18	1.23	1.30
4	A	102	SAH	C8-N7	2.12	1.35	1.31
4	A	101	SAH	C8-N7	2.06	1.35	1.31
4	A	101	SAH	OXT-C	-2.03	1.24	1.30

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	102	SAH	N3-C2-N1	-5.62	120.08	128.58
4	A	101	SAH	N3-C2-N1	-5.02	120.99	128.58
4	A	101	SAH	C5-C4-N3	-4.61	120.37	126.72
4	A	102	SAH	C5-C4-N3	-4.45	120.59	126.72
4	A	102	SAH	N9-C8-N7	-3.42	109.09	113.94
4	A	102	SAH	C2-N3-C4	3.40	120.13	111.83
4	A	102	SAH	C5-N7-C8	3.31	108.66	103.45
4	A	101	SAH	C2-N3-C4	3.22	119.69	111.83
4	A	101	SAH	C5-N7-C8	3.18	108.45	103.45
4	A	101	SAH	N9-C8-N7	-3.10	109.54	113.94
4	A	102	SAH	OXT-C-O	-2.99	117.30	124.08
4	A	102	SAH	C5'-SD-CG	-2.83	93.86	102.26
4	A	102	SAH	N3-C4-N9	2.78	131.89	127.17
4	A	101	SAH	N3-C4-N9	2.75	131.85	127.17
4	A	101	SAH	OXT-C-O	-2.73	117.89	124.08
4	A	102	SAH	C4-C5-N7	-2.52	107.70	110.58
4	A	101	SAH	C4-C5-N7	-2.50	107.72	110.58
4	A	101	SAH	O4'-C1'-N9	2.38	112.66	108.09
4	A	101	SAH	C5'-SD-CG	-2.17	95.82	102.26
4	A	101	SAH	C6-C5-C4	2.08	120.02	117.18

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	102	SAH	C3'-C4'-C5'-SD
4	A	101	SAH	C-CA-CB-CG
4	A	102	SAH	CB-CG-SD-C5'
4	A	101	SAH	O-C-CA-N
4	A	101	SAH	CB-CG-SD-C5'
4	A	101	SAH	O-C-CA-CB
4	A	102	SAH	O-C-CA-CB


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	101	SAH	2	0
4	A	102	SAH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/285 (91%)	0.26	12 (4%) 37 39	7, 20, 33, 39	5 (1%)
1	B	252/285 (88%)	0.95	51 (20%) 3 2	12, 26, 55, 57	3 (1%)
2	P	7/11 (63%)	0.51	1 (14%) 6 5	18, 21, 26, 28	0
2	Q	5/11 (45%)	1.99	3 (60%) 0 0	39, 39, 47, 47	0
All	All	525/592 (88%)	0.61	67 (12%) 7 7	7, 22, 44, 57	8 (1%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1223	PHE	5.9
1	B	1232	CYS	5.1
1	B	1224	SER	5.0
1	B	1222	LEU	4.7
1	B	998	VAL	4.6
1	B	1235	SER	4.3
1	B	1218	ILE	4.2
1	B	1225	CYS	4.2
1	B	1217	ASP	4.0
1	B	1216	TRP	4.0
1	B	978	ILE	3.9
1	B	1102	THR	3.4
1	B	1093	LEU	3.4
1	A	1150	ASP	3.4
1	A	1138	GLU	3.3
2	Q	9	THR	3.3
1	B	1228	GLY	3.3
1	B	995	VAL	3.3
1	B	1220	GLY	3.2
1	B	1227	CYS	3.1
1	B	1003	CYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1153	VAL	3.0
1	B	1173	GLU	3.0
1	B	1229	SER	3.0
1	B	1226	ARG	2.9
1	B	1002	PRO	2.9
1	B	1215	PHE	2.9
1	B	997	ALA	2.8
1	A	1222	LEU	2.8
1	A	975	VAL	2.8
1	B	1221	LYS	2.8
1	A	1104	ASP	2.7
1	A	1094	ARG	2.7
1	B	1234	HIS	2.7
1	B	1172	CYS	2.7
1	B	1100	TYR	2.6
1	B	1067	GLU	2.6
2	P	9	THR	2.6
1	B	1136	VAL	2.6
1	B	1104	ASP	2.6
1	B	1138	GLU	2.5
1	B	1139	GLU	2.5
1	A	1060	LEU	2.5
1	A	1214	ARG	2.5
1	B	979	VAL	2.4
1	B	1204	GLY	2.4
1	B	981	ARG	2.3
1	B	1094	ARG	2.3
1	B	1034	ILE	2.2
1	A	1103	ARG	2.2
1	B	996	ASN	2.2
1	B	1103	ARG	2.2
1	B	1006	ASN	2.2
1	B	1020	ASN	2.2
1	A	1046	GLN	2.1
1	B	1231	LYS	2.1
1	B	1107	TRP	2.1
1	B	1065	MET	2.1
2	Q	5	ALA	2.1
1	B	1214	ARG	2.1
1	A	1235	SER	2.1
1	B	1099	LEU	2.0
1	B	1219	LYS	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	989	ARG	2.0
1	B	1233	ARG	2.0
2	Q	4	THR	2.0
1	A	1043	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MLY	Q	7	11/12	0.81	0.19	20,34,36,37	0
2	MLY	P	7	11/12	0.96	0.06	13,14,17,17	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SAH	A	102	26/26	0.84	0.16	44,46,47,48	0
3	ZN	B	504	1/1	0.86	0.11	43,43,43,43	0
3	ZN	A	504	1/1	0.96	0.05	25,25,25,25	0
4	SAH	A	101	26/26	0.97	0.06	17,21,22,23	0
3	ZN	A	503	1/1	0.98	0.03	17,17,17,17	0
3	ZN	A	502	1/1	0.99	0.03	16,16,16,16	0
3	ZN	A	501	1/1	0.99	0.02	17,17,17,17	0
3	ZN	B	501	1/1	1.00	0.01	16,16,16,16	0
3	ZN	B	502	1/1	1.00	0.02	16,16,16,16	0
3	ZN	B	503	1/1	1.00	0.02	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.