



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:15 PM UTC

PDB ID : 2RFV / pdb\_00002rfv  
Title : High resolution structure of L-methionine gamma-lyase from *Citrobacter freundii*  
Authors : Nikulin, A.D.; Revtovich, S.V.; Morozova, E.A.; Nevskaya, N.A.; Nikonov, S.V.; Garber, M.B.; Demidkina, T.V.  
Deposited on : 2007-10-02  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1216 (1.36-1.36)
Clashscore	190562	1232 (1.36-1.36)
Ramachandran outliers	187476	1220 (1.36-1.36)
Sidechain outliers	187428	1220 (1.36-1.36)
RSRZ outliers	180081	1214 (1.36-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3552 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methionine gamma-lyase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	391	3169	2008	542	597	1	21	0	31	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	133	PHE	LEU	SEE REMARK 999	UNP Q84AR1
A	137	ALA	GLY	SEE REMARK 999	UNP Q84AR1

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	382	Total O 382 382	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Methionine gamma-lyase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.45Å 122.80Å 127.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.35 20.00 – 1.35	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-1.35) 98.7 (20.00-1.35)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.76 (at 1.35Å)	Xtrriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.151 , 0.177 0.146 , 0.172	Depositor DCC
$R_{free}$ test set	4777 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.4	Xtrriage
Anisotropy	0.495	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.46 , 75.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.011 for -h,-l,-k	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3552	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, CSO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.40	3/3263 (0.1%)	0.65	1/4422 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	116	THR	C-O	-6.91	1.16	1.24
1	A	117	HIS	C-O	-5.49	1.17	1.24
1	A	118	ALA	C-O	-5.35	1.17	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	35	SER	N-CA-C	-5.22	107.46	113.88

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3169	0	3208	73	0
2	A	1	0	0	0	0
3	A	382	0	0	16	0
All	All	3552	0	3208	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:HIS:NE2	1:A:132[B]:ARG:HD2	1.57	1.20
1:A:105:HIS:NE2	1:A:132[B]:ARG:CD	2.15	1.09
1:A:132[B]:ARG:HH21	1:A:132[B]:ARG:HG3	1.24	1.02
1:A:4:CSO:HB3	3:A:633:HOH:O	1.67	0.93
1:A:60:ARG:HH11	1:A:60:ARG:HG3	1.38	0.86
1:A:58:TYR:HE2	1:A:60:ARG:HD3	1.41	0.86
1:A:105:HIS:CE1	1:A:132[A]:ARG:HD2	2.13	0.84
1:A:4:CSO:HB2	3:A:756:HOH:O	1.77	0.83
1:A:374[B]:ARG:HD2	1:A:375:LEU:N	1.99	0.78
1:A:39:PHE:HZ	1:A:57:ILE:HD13	1.52	0.74
1:A:132[B]:ARG:HG3	1:A:132[B]:ARG:NH2	1.87	0.73
1:A:7[B]:TYR:CE1	3:A:756:HOH:O	2.44	0.70
1:A:105:HIS:NE2	1:A:132[B]:ARG:HD3	2.06	0.69
1:A:60:ARG:NE	3:A:777:HOH:O	2.25	0.69
1:A:60:ARG:HG3	1:A:60:ARG:NH1	2.01	0.69
1:A:58:TYR:CE2	1:A:60:ARG:HD3	2.28	0.68
1:A:385[B]:ILE:HD12	3:A:755:HOH:O	1.94	0.67
1:A:132[A]:ARG:HD3	1:A:145:ALA:HB1	1.74	0.67
1:A:312[B]:ILE:HD11	1:A:340:LEU:HD13	1.74	0.67
1:A:40:ASP:H	1:A:44:GLN:NE2	1.93	0.66
1:A:4:CSO:CB	3:A:756:HOH:O	2.40	0.66
1:A:7[B]:TYR:HE1	3:A:756:HOH:O	1.78	0.65
1:A:361:GLU:H	1:A:361:GLU:CD	2.04	0.64
1:A:60:ARG:CZ	3:A:777:HOH:O	2.45	0.64
1:A:206:HIS:HB2	1:A:222[A]:ILE:HG22	1.81	0.61
1:A:105:HIS:NE2	1:A:132[A]:ARG:HD2	2.16	0.60
1:A:189[A]:MET:HA	1:A:189[A]:MET:HE2	1.83	0.60
1:A:132[B]:ARG:HH21	1:A:132[B]:ARG:CG	2.06	0.60
1:A:58:TYR:CD2	1:A:60:ARG:HG2	2.36	0.59
1:A:19[B]:GLN:NE2	3:A:746:HOH:O	2.35	0.59
1:A:19[B]:GLN:NE2	3:A:763:HOH:O	2.18	0.57
1:A:325[B]:ARG:NH1	1:A:352:SER:HB2	2.22	0.55
1:A:328:ILE:HD13	1:A:349:HIS:HB3	1.91	0.51
1:A:132[B]:ARG:CZ	1:A:147:ARG:NH2	2.74	0.50
1:A:157:THR:HA	1:A:158:PRO:C	2.36	0.50
1:A:59:THR:HA	1:A:62:GLY:O	2.12	0.50
1:A:60:ARG:HD3	3:A:777:HOH:O	2.13	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132[A]:ARG:HE	1:A:147:ARG:HE	1.61	0.49
1:A:391:HIS:CE1	1:A:395:LYS:HE3	2.49	0.48
1:A:60:ARG:CD	3:A:777:HOH:O	2.62	0.48
1:A:105:HIS:NE2	1:A:132[A]:ARG:HG3	2.30	0.46
1:A:210:LLP:O3	1:A:210:LLP:NZ	2.47	0.46
1:A:385[B]:ILE:CD1	3:A:755:HOH:O	2.60	0.46
1:A:240:ASP:C	3:A:760:HOH:O	2.58	0.45
1:A:60:ARG:HA	1:A:245[A]:CYS:SG	2.57	0.45
1:A:40:ASP:H	1:A:44:GLN:HE21	1.64	0.45
1:A:60:ARG:HH22	1:A:239:LYS:HG3	1.82	0.45
1:A:289:TYR:HB2	1:A:314[B]:SER:OG	2.18	0.44
1:A:325[B]:ARG:HH11	1:A:352:SER:HB2	1.82	0.44
1:A:208:VAL:HG22	1:A:220:GLY:C	2.43	0.44
1:A:132[B]:ARG:NE	1:A:147:ARG:NH2	2.66	0.43
1:A:163[A]:LEU:HD12	1:A:163[A]:LEU:N	2.32	0.43
1:A:60:ARG:HH11	1:A:60:ARG:CG	2.16	0.43
1:A:58:TYR:CE2	1:A:60:ARG:HG2	2.54	0.43
1:A:124:MET:HB2	1:A:125:PRO:HD3	2.01	0.43
1:A:132[B]:ARG:CZ	1:A:147:ARG:HH22	2.32	0.43
1:A:387[B]:ASN:ND2	3:A:771:HOH:O	2.52	0.43
1:A:345:THR:OG1	1:A:385[B]:ILE:HD13	2.20	0.42
1:A:60:ARG:O	1:A:245[A]:CYS:SG	2.76	0.42
1:A:2:SER:OG	1:A:3:ASP:N	2.51	0.42
1:A:7[B]:TYR:HB3	1:A:11:THR:HB	2.01	0.42
1:A:364:LEU:HD23	1:A:364:LEU:HA	1.92	0.42
1:A:122[B]:HIS:C	1:A:125:PRO:HD2	2.45	0.41
1:A:306[B]:MET:HE2	3:A:413:HOH:O	2.20	0.41
1:A:359:ALA:HB1	1:A:361:GLU:OE1	2.20	0.41
1:A:327[A]:MET:HE3	1:A:393:ILE:HG13	2.03	0.41
1:A:340:LEU:HB3	1:A:374[B]:ARG:NH1	2.36	0.41
1:A:113:TYR:CE2	1:A:115[B]:CYS:HB3	2.56	0.41
1:A:356:SER:N	1:A:357:PRO:CD	2.84	0.40
1:A:39:PHE:CZ	1:A:57:ILE:HD13	2.42	0.40
1:A:186:ASN:HB3	1:A:206:HIS:CE1	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	416/398 (104%)	412 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	348/323 (108%)	343 (99%)	5 (1%)	59	28

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	GLN
1	A	57	ILE
1	A	60	ARG
1	A	61	LEU
1	A	361	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	44	GLN
1	A	130	ASN

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	A	348	GLN
1	A	391	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	LLP	A	210	1	23,24,25	1.69	4 (17%)	25,32,34	1.44	3 (12%)
1	CSO	A	4	1	3,6,7	0.50	0	1,6,8	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	210	1	-	6/16/17/19	0/1/1/1
1	CSO	A	4	1	-	0/1/5/7	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	210	LLP	O3-C3	-5.81	1.23	1.36
1	A	210	LLP	C2-N1	2.31	1.38	1.33
1	A	210	LLP	C4-C4'	2.19	1.51	1.46
1	A	210	LLP	C4'-NZ	2.07	1.34	1.27

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	210	LLP	OP4-C5'-C5	4.63	118.04	109.36
1	A	210	LLP	C4-C4'-NZ	-3.16	109.44	124.04
1	A	210	LLP	C5-C6-N1	-2.24	120.19	123.83

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	210	LLP	O-C-CA-CB
1	A	210	LLP	C4-C4'-NZ-CE
1	A	210	LLP	CD-CE-NZ-C4'
1	A	210	LLP	CG-CD-CE-NZ
1	A	210	LLP	CA-CB-CG-CD
1	A	210	LLP	C3-C4-C4'-NZ

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	210	LLP	1	0
1	A	4	CSO	3	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	389/398 (97%)	0.34	40 (10%) <b>12</b> <b>16</b>	8, 18, 46, 66	31 (7%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	47	ALA	7.2
1	A	61	LEU	6.5
1	A	57	ILE	6.0
1	A	366	ALA	5.4
1	A	358	VAL	5.3
1	A	58	TYR	4.9
1	A	62	GLY	4.8
1	A	2	SER	4.7
1	A	45	GLY	4.7
1	A	55	GLY	4.7
1	A	40	ASP	4.3
1	A	46	ALA	4.2
1	A	113	TYR	4.1
1	A	56	TYR	3.6
1	A	42	ALA	3.6
1	A	357	PRO	3.4
1	A	114	GLY	3.3
1	A	161	PRO	3.3
1	A	39	PHE	3.3
1	A	364	LEU	3.2
1	A	23	SER	3.2
1	A	361	GLU	3.1
1	A	59	THR	3.1
1	A	49	PHE	3.1
1	A	351	ALA	3.1
1	A	354	THR	3.1
1	A	7[A]	TYR	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	353	MET	2.8
1	A	356	SER	2.8
1	A	359	ALA	2.8
1	A	360	PRO	2.7
1	A	355	HIS	2.6
1	A	6	THR	2.4
1	A	160	ASN	2.3
1	A	122[A]	HIS	2.2
1	A	328	ILE	2.1
1	A	188	PHE	2.1
1	A	48	ARG	2.0
1	A	391	HIS	2.0
1	A	115[A]	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSO	A	4	7/8	0.77	0.15	27,29,37,38	0
1	LLP	A	210	24/25	0.94	0.12	12,23,30,31	0

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CL	A	399	1/1	0.99	0.21	27,27,27,27	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.