



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 09:42 AM UTC

PDB ID : 3RF3 / pdb\_00003rf3  
Title : Shigella IpaA-VBS3 in complex with human vinculin  
Authors : Park, H.; Sharff, A.; Izard, T.  
Deposited on : 2011-04-05  
Resolution : 1.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6728 (1.64-1.60)
Clashscore	190562	7023 (1.64-1.60)
Ramachandran outliers	187476	6898 (1.64-1.60)
Sidechain outliers	187428	6896 (1.64-1.60)
RSRZ outliers	180081	6727 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	
1	B	258	
2	C	29	
2	D	29	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9685 atoms, of which 4498 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vinculin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	257	4045	1260	2048	340	385	12	0	2	0
1	B	258	4077	1268	2063	342	391	13	0	0	0


- Molecule 2 is a protein called Invasin ipaA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	C	23	371	113	192	30	36	0	0	0
2	D	24	378	115	195	31	37	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	484	GLY	-	expression tag	UNP P18010
C	485	SER	-	expression tag	UNP P18010
C	486	HIS	-	expression tag	UNP P18010
C	487	MET	-	expression tag	UNP P18010
D	484	GLY	-	expression tag	UNP P18010
D	485	SER	-	expression tag	UNP P18010
D	486	HIS	-	expression tag	UNP P18010
D	487	MET	-	expression tag	UNP P18010

- Molecule 3 is CACODYLATE ION (CCD ID: CAC) (formula: C<sub>2</sub>H<sub>6</sub>AsO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	As	C	O		
3	A	1	5	1	2	2	0	0
3	B	1	5	1	2	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	358	Total 358	O 358	0	0
4	B	389	Total 389	O 389	0	0
4	C	31	Total 31	O 31	0	0
4	D	26	Total 26	O 26	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vinculin




- Molecule 1: Vinculin



- Molecule 2: Invasin ipaA



- Molecule 2: Invasin ipaA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.58Å 85.32Å 137.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.00 – 1.61 53.00 – 1.61	Depositor EDS
% Data completeness (in resolution range)	(Not available) (53.00-1.61) 99.9 (53.00-1.61)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.46 (at 1.61Å)	Xtrriage
Refinement program	BUSTER 2.13.0	Depositor
R, $R_{free}$	0.184 , 0.195 0.180 , 0.198	Depositor DCC
$R_{free}$ test set	4129 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.7	Xtrriage
Anisotropy	0.637	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 59.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9685	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 52.53 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.7989e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.83	4/2031 (0.2%)	1.03	7/2753 (0.3%)
1	B	0.83	3/2041 (0.1%)	0.99	4/2765 (0.1%)
2	C	0.73	0/179	0.75	0/240
2	D	0.73	0/183	0.76	0/245
All	All	0.82	7/4434 (0.2%)	0.99	11/6003 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	190	MET	SD-CE	-7.53	1.60	1.79
1	B	174	MET	SD-CE	-7.25	1.61	1.79
1	A	190	MET	SD-CE	-6.54	1.63	1.79
1	A	174	MET	SD-CE	-6.06	1.64	1.79
1	A	98	ASP	CA-C	5.50	1.59	1.52
1	A	106	ASP	C-N	5.45	1.41	1.33
1	B	223	ILE	CG1-CD1	-5.26	1.31	1.51

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	TYR	N-CA-C	-7.07	103.57	111.28
1	A	31	GLU	N-CA-C	6.01	119.72	111.54
1	B	75	PRO	N-CA-C	5.92	117.93	110.70
1	B	30	GLY	CA-C-N	5.76	130.54	122.08
1	B	30	GLY	C-N-CA	5.76	130.54	122.08
1	A	223	ILE	N-CA-C	5.59	115.78	110.53
1	B	31	GLU	N-CA-C	5.05	118.56	111.90
1	A	28	GLU	CA-C-N	5.04	131.17	121.54
1	A	28	GLU	C-N-CA	5.04	131.17	121.54
1	A	106	ASP	CA-C-N	5.01	126.99	120.28

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	106	ASP	C-N-CA	5.01	126.99	120.28

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1997	2048	2040	3	0
1	B	2014	2063	2063	5	0
2	C	179	192	192	0	0
2	D	183	195	195	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	358	0	0	0	0
4	B	389	0	0	0	0
4	C	31	0	0	0	0
4	D	26	0	0	0	0
All	All	5187	4498	4490	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:211:ILE:O	1:A:215:THR:HG23	2.10	0.52
1:B:31:GLU:HB2	1:B:102:VAL:HG13	1.92	0.50
1:B:154:MET:HE2	1:B:213:VAL:HG11	1.92	0.50
1:B:171:MET:HE1	1:B:241:ILE:HG12	1.94	0.48
1:B:70:LEU:HD12	1:B:125:THR:HG22	1.95	0.48
1:A:44:VAL:HG21	1:A:92:ALA:HB2	2.01	0.42
1:A:91:ALA:HB2	1:A:107:TYR:HB3	2.02	0.42
1:B:223:ILE:HG23	1:B:224:GLU:N	2.36	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/258 (100%)	253 (98%)	3 (1%)	1 (0%)	30	14
1	B	256/258 (99%)	254 (99%)	2 (1%)	0	100	100
2	C	21/29 (72%)	21 (100%)	0	0	100	100
2	D	22/29 (76%)	22 (100%)	0	0	100	100
All	All	556/574 (97%)	550 (99%)	5 (1%)	1 (0%)	43	25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/229 (98%)	223 (100%)	1 (0%)	84	75
1	B	227/229 (99%)	225 (99%)	2 (1%)	70	54
2	C	22/26 (85%)	22 (100%)	0	100	100
2	D	22/26 (85%)	22 (100%)	0	100	100
All	All	495/510 (97%)	492 (99%)	3 (1%)	78	66

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	ILE
1	B	23	LEU
1	B	109	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	GLN
1	A	68	GLN
1	A	185	GLN
1	A	249	GLN
1	B	68	GLN
1	B	90	GLN
1	B	185	GLN
1	B	249	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CAC	B	300	-	2,4,4	2.58	2 (100%)	4,6,6	1.04	0
3	CAC	A	300	-	2,4,4	2.39	2 (100%)	4,6,6	1.78	1 (25%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	300	CAC	AS-C2	2.74	1.96	1.90
3	A	300	CAC	AS-C1	2.55	1.96	1.90
3	B	300	CAC	AS-C1	2.40	1.96	1.90
3	A	300	CAC	AS-C2	2.21	1.95	1.90

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	300	CAC	O1-AS-C2	-2.80	108.02	111.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	257/258 (99%)	0.41	20 (7%) 19 20	18, 29, 53, 80	1 (0%)
1	B	258/258 (100%)	0.23	14 (5%) 31 33	19, 28, 52, 72	0
2	C	23/29 (79%)	0.14	2 (8%) 16 17	17, 24, 44, 63	0
2	D	24/29 (82%)	0.01	3 (12%) 8 8	18, 25, 56, 81	0
All	All	562/574 (97%)	0.30	39 (6%) 23 24	17, 28, 53, 81	1 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	32	VAL	6.5
2	C	510	ILE	6.3
1	A	30	GLY	5.5
1	B	25	ILE	5.0
1	B	26	MET	4.8
1	A	31	GLU	4.4
1	A	28	GLU	4.4
1	A	27	HIS	4.3
1	A	23	LEU	4.0
1	B	32	VAL	3.9
1	A	25	ILE	3.5
1	A	107	TYR	3.5
1	B	1	MET	3.5
1	B	30	GLY	3.4
1	A	257	ALA	3.4
2	D	511	GLY	3.4
1	B	258	TRP	3.4
2	D	510	ILE	3.3
2	D	488	THR	3.2
1	B	27	HIS	3.2
1	B	257	ALA	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	258	TRP	3.1
1	B	31	GLU	3.0
1	A	24	VAL	2.8
1	A	108	LEU	2.8
1	B	23	LEU	2.8
1	A	29	GLU	2.7
1	B	223	ILE	2.7
1	B	28	GLU	2.6
1	A	126	PHE	2.5
2	C	488	THR	2.4
1	A	26	MET	2.3
1	A	223	ILE	2.3
1	B	150	VAL	2.3
1	A	37	ILE	2.2
1	A	22	HIS	2.1
1	A	103	PRO	2.1
1	A	69	ILE	2.0
1	B	107	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CAC	A	300	5/5	0.69	0.29	158,158,158,159	0
3	CAC	B	300	5/5	0.73	0.29	159,159,159,160	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.