



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 03:38 PM EDT

PDB ID : 4RF7 / pdb_00004rf7
Title : Crystal structure of double-domain arginine kinase from *Anthopleura japonica* in complex with substrate L-arginine
Authors : Wang, Z.; Qiao, Z.; Ye, S.; Zhang, R.
Deposited on : 2014-09-25
Resolution : 2.10 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	718	
1	B	718	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	B	1003	-	-	X	-
3	ARG	A	807	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 12184 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Arginine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	692	5470	3455	964	1026	25	0	1	0
1	B	700	5538	3493	979	1042	24	0	1	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP O15992
A	-1	PRO	-	expression tag	UNP O15992
A	0	HIS	-	expression tag	UNP O15992
B	-2	GLY	-	expression tag	UNP O15992
B	-1	PRO	-	expression tag	UNP O15992
B	0	HIS	-	expression tag	UNP O15992

- Molecule 2 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is ARGinine (CCD ID: ARG) (formula: C₆H₁₅N₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 12 6 4 2	0	0
3	A	1	Total C N O 12 6 4 2	0	0
3	A	1	Total C N O 12 6 4 2	0	0
3	B	1	Total C N O 12 6 4 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			12	6	4	2		
3	B	1	Total	C	N	O	0	0
			12	6	4	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	545	Total	O	0	0
			545	545		
4	B	527	Total	O	0	0
			527	527		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Arginine kinase



- Molecule 1: Arginine kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.50Å 59.27Å 162.26Å 90.00° 91.31° 90.00°	Depositor
Resolution (Å)	47.85 – 2.10 47.85 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.85-2.10) 91.0 (47.85-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.10Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.180 , 0.217 0.183 , 0.222	Depositor DCC
R_{free} test set	4401 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtrriage
Anisotropy	0.345	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.028 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12184	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/5582	0.71	0/7527
1	B	0.32	0/5653	0.75	3/7622 (0.0%)
All	All	0.31	0/11235	0.73	3/15149 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	643	ILE	CA-C-N	6.73	128.26	119.84
1	B	643	ILE	C-N-CA	6.73	128.26	119.84
1	B	643	ILE	N-CA-C	5.43	113.41	107.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5470	0	5469	38	0
1	B	5538	0	5522	50	0
2	A	16	0	12	1	0
2	B	16	0	12	5	0
3	A	36	0	36	8	0
3	B	36	0	36	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	545	0	0	12	1
4	B	527	0	0	12	1
All	All	12184	0	11087	91	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:130:ARG:HE	2:B:1003:ACT:H1	1.22	1.01
1:B:177:MET:O	4:B:1580:HOH:O	1.98	0.81
1:B:590:GLU:HG3	1:B:601:ARG:HH22	1.47	0.78
1:B:128:ARG:HH22	2:B:1003:ACT:H3	1.52	0.73
1:B:374:LEU:O	1:B:384:ARG:NH1	2.23	0.69
1:A:1:MET:O	4:A:1192:HOH:O	2.12	0.68
1:B:594:ASP:OD1	4:B:1337:HOH:O	2.11	0.68
1:B:664:ARG:NH1	4:B:1516:HOH:O	2.25	0.68
1:A:175:SER:OG	4:A:1269:HOH:O	2.11	0.67
1:B:662:GLN:HB3	1:B:686:ARG:HD3	1.76	0.66
1:B:126:SER:HB2	3:B:1006:ARG:HD2	1.79	0.65
1:A:166:ASP:OD1	1:A:252:ARG:NH2	2.31	0.64
3:B:1007:ARG:N	4:B:1353:HOH:O	2.31	0.62
1:B:636:ARG:NH2	1:B:682:ASP:OD1	2.33	0.61
1:A:705:LYS:NZ	1:A:709:GLU:OE2	2.33	0.61
1:A:549:ARG:NH1	4:A:1334:HOH:O	2.34	0.61
1:B:72:TYR:HH	3:B:1005:ARG:N	2.00	0.60
1:B:642:LYS:HA	1:B:680:VAL:HB	1.84	0.60
1:A:615:PHE:H	3:A:807:ARG:HG3	1.67	0.59
1:B:577:TRP:HB2	1:B:585:ARG:HB3	1.85	0.59
3:B:1006:ARG:NH1	4:B:1151:HOH:O	2.36	0.58
1:B:166:ASP:OD1	1:B:252:ARG:NH2	2.37	0.58
1:A:448:PRO:O	4:A:1044:HOH:O	2.17	0.58
1:A:651:ASP:OD1	1:A:654:LYS:NZ	2.38	0.57
1:B:590:GLU:HG3	1:B:601:ARG:NH2	2.19	0.57
1:B:376:ASN:HB3	1:B:379:VAL:HG23	1.87	0.56
1:B:268:LYS:NZ	4:B:1268:HOH:O	2.37	0.55
1:B:336:ARG:NH1	4:B:1380:HOH:O	2.39	0.54
1:B:128:ARG:HH12	2:B:1003:ACT:H2	1.72	0.53
1:B:305:THR:OG1	4:B:1236:HOH:O	2.18	0.53
1:B:649:HIS:CG	1:B:650:PRO:HD2	2.45	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:LYS:HD3	1:B:261:LEU:HD22	1.90	0.52
1:A:308:HIS:ND1	4:A:1260:HOH:O	2.34	0.51
1:B:372:GLU:OE2	1:B:375:LYS:NZ	2.36	0.51
1:A:130:ARG:HD2	1:A:233:ARG:CZ	2.41	0.51
1:B:225:TRP:HB2	1:B:233[A]:ARG:HB3	1.93	0.51
1:A:153:GLU:HA	1:A:226:LEU:HD21	1.93	0.50
1:A:466:ALA:O	4:A:970:HOH:O	2.19	0.50
1:B:443:GLU:OE1	4:B:1442:HOH:O	2.20	0.50
1:B:4:PRO:HD2	4:B:1545:HOH:O	2.11	0.49
1:B:179:GLU:OE1	4:B:1584:HOH:O	2.19	0.49
1:B:534:ARG:O	1:B:538:VAL:HG23	2.13	0.48
2:A:803:ACT:O	3:A:807:ARG:NH1	2.42	0.48
1:B:522:GLN:OE1	1:B:524:TYR:OH	2.13	0.48
1:B:94:HIS:CG	1:B:337:LEU:HD12	2.49	0.48
1:B:479:THR:HG21	1:B:595:ILE:HA	1.95	0.48
1:A:99:LEU:HD22	1:A:268:LYS:HG2	1.97	0.47
1:A:126:SER:OG	3:A:806:ARG:NH1	2.39	0.47
1:A:166:ASP:CG	1:A:252:ARG:HH22	2.22	0.47
1:A:233:ARG:CZ	4:A:1307:HOH:O	2.63	0.47
1:B:237:MET:SD	1:B:237:MET:N	2.87	0.47
1:B:47:LYS:HG2	1:B:53:THR:HG22	1.97	0.47
1:A:610:ASP:OD1	3:A:807:ARG:HD3	2.15	0.47
1:A:610:ASP:OD2	3:A:807:ARG:NH2	2.48	0.47
1:B:126:SER:CB	3:B:1006:ARG:HD2	2.45	0.46
1:B:480:ARG:HH21	1:B:587:ILE:HD13	1.81	0.46
1:B:644:PRO:HD2	1:B:711:GLU:OE2	2.16	0.46
1:A:643:ILE:O	1:A:647:SER:HB3	2.16	0.46
1:B:523:TYR:CE1	1:B:578:ILE:HD12	2.51	0.46
1:B:314:ILE:HD11	1:B:325:ASP:CG	2.42	0.45
1:B:128:ARG:HH22	2:B:1003:ACT:CH3	2.26	0.45
1:A:331:ILE:HB	1:A:348:MET:HE1	1.98	0.45
1:B:317:GLU:OE1	1:B:318:HIS:NE2	2.50	0.45
1:A:233:ARG:NH2	4:A:1366:HOH:O	2.50	0.45
1:A:615:PHE:H	3:A:807:ARG:CD	2.30	0.45
1:A:119:PRO:HD2	4:A:945:HOH:O	2.17	0.44
1:A:646:ALA:HB2	1:A:710:ILE:HB	1.98	0.44
1:A:146:LYS:NZ	1:A:210:GLU:OE1	2.46	0.44
1:A:523:TYR:CE1	1:A:578:ILE:HD12	2.52	0.44
1:B:708:LEU:O	1:B:712:LYS:HG2	2.19	0.43
1:A:49:LYS:HB3	1:A:49:LYS:HE2	1.72	0.43
1:A:544:PHE:CD1	1:A:580:GLU:HG3	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:ASP:CG	1:B:252:ARG:HH22	2.26	0.43
1:A:615:PHE:H	3:A:807:ARG:CG	2.31	0.42
1:B:171:TYR:OH	1:B:213:GLY:HA3	2.19	0.42
1:A:2:ALA:HB2	1:A:197:ARG:HD2	2.02	0.42
1:A:488:LYS:NZ	3:A:807:ARG:O	2.30	0.42
1:B:233[B]:ARG:HH21	2:B:1003:ACT:H2	1.84	0.42
1:A:183:GLN:NE2	1:A:187:ASP:OD1	2.53	0.42
1:B:482:ARG:HD3	1:B:585:ARG:HD2	2.01	0.42
1:A:479:THR:HG21	1:A:595:ILE:HA	2.02	0.42
1:B:644:PRO:HD2	1:B:711:GLU:CD	2.44	0.42
1:A:233:ARG:HD2	4:A:1307:HOH:O	2.18	0.41
1:B:237:MET:HB2	3:B:1006:ARG:HG2	2.01	0.41
1:B:237:MET:HB2	3:B:1006:ARG:CG	2.51	0.41
1:A:585:ARG:NH2	4:A:994:HOH:O	2.30	0.41
1:B:47:LYS:HE2	4:B:1405:HOH:O	2.19	0.41
1:A:233:ARG:NH2	4:A:1307:HOH:O	2.54	0.41
1:A:708:LEU:HD23	1:A:708:LEU:HA	1.92	0.41
1:B:407:TYR:HA	1:B:410:ILE:HG12	2.02	0.41
1:A:237:MET:SD	1:A:237:MET:N	2.94	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1407:HOH:O	4:B:1617:HOH:O[2_546]	1.97	0.23

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	687/718 (96%)	666 (97%)	18 (3%)	3 (0%)	30 28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	697/718 (97%)	677 (97%)	18 (3%)	2 (0%)	36	36
All	All	1384/1436 (96%)	1343 (97%)	36 (3%)	5 (0%)	30	28

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	644	PRO
1	A	580	GLU
1	A	644	PRO
1	A	228	GLU
1	B	580	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	597/616 (97%)	589 (99%)	8 (1%)	61	69
1	B	604/616 (98%)	596 (99%)	8 (1%)	61	69
All	All	1201/1232 (98%)	1185 (99%)	16 (1%)	59	69

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	130	ARG
1	A	237	MET
1	A	314	ILE
1	A	357	GLU
1	A	477	ARG
1	A	654	LYS
1	A	666	ILE
1	B	369	VAL
1	B	384	ARG
1	B	585	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	589	MET
1	B	666	ILE
1	B	680	VAL
1	B	686	ARG
1	B	713	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	A	453	ASN
1	A	640	HIS
1	B	184	GLN
1	B	453	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACT	B	1002	-	3,3,3	0.83	0	3,3,3	1.31	0
3	ARG	A	805	-	10,11,11	0.76	1 (10%)	9,13,13	0.90	1 (11%)
2	ACT	A	804	-	3,3,3	0.81	0	3,3,3	1.29	0
3	ARG	B	1005	-	10,11,11	0.74	1 (10%)	9,13,13	0.87	1 (11%)
2	ACT	B	1001	-	3,3,3	0.81	0	3,3,3	1.34	0
3	ARG	B	1007	-	10,11,11	0.75	1 (10%)	9,13,13	0.94	1 (11%)
3	ARG	A	806	-	10,11,11	0.77	1 (10%)	9,13,13	0.96	1 (11%)
2	ACT	B	1004	-	3,3,3	0.81	0	3,3,3	1.33	0
3	ARG	B	1006	-	10,11,11	0.73	1 (10%)	9,13,13	0.92	1 (11%)
2	ACT	A	801	-	3,3,3	0.78	0	3,3,3	1.38	0
3	ARG	A	807	-	10,11,11	0.77	1 (10%)	9,13,13	0.99	1 (11%)
2	ACT	B	1003	-	3,3,3	0.79	0	3,3,3	1.11	0
2	ACT	A	802	-	3,3,3	0.85	0	3,3,3	1.31	0
2	ACT	A	803	-	3,3,3	0.83	0	3,3,3	1.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ARG	A	805	-	-	0/11/11/11	-
3	ARG	B	1005	-	-	1/11/11/11	-
3	ARG	B	1007	-	-	4/11/11/11	-
3	ARG	A	806	-	-	5/11/11/11	-
3	ARG	B	1006	-	-	1/11/11/11	-
3	ARG	A	807	-	-	1/11/11/11	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	807	ARG	OXT-C	-2.28	1.23	1.30
3	A	806	ARG	OXT-C	-2.22	1.23	1.30
3	A	805	ARG	OXT-C	-2.19	1.23	1.30
3	B	1007	ARG	OXT-C	-2.17	1.23	1.30
3	B	1006	ARG	OXT-C	-2.15	1.23	1.30
3	B	1005	ARG	OXT-C	-2.11	1.23	1.30

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	807	ARG	OXT-C-O	-2.86	117.60	124.08
3	A	806	ARG	OXT-C-O	-2.83	117.65	124.08
3	B	1007	ARG	OXT-C-O	-2.77	117.80	124.08
3	B	1006	ARG	OXT-C-O	-2.72	117.90	124.08
3	A	805	ARG	OXT-C-O	-2.59	118.21	124.08
3	B	1005	ARG	OXT-C-O	-2.50	118.40	124.08

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	806	ARG	O-C-CA-N
3	B	1007	ARG	N-CA-CB-CG
3	A	806	ARG	NE-CD-CG-CB
3	A	806	ARG	OXT-C-CA-N
3	B	1006	ARG	CA-CB-CG-CD
3	A	807	ARG	CA-CB-CG-CD
3	B	1007	ARG	O-C-CA-CB
3	B	1007	ARG	OXT-C-CA-CB
3	A	806	ARG	O-C-CA-CB
3	A	806	ARG	OXT-C-CA-CB
3	B	1007	ARG	C-CA-CB-CG
3	B	1005	ARG	NE-CD-CG-CB

There are no ring outliers.

7 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1005	ARG	1	0
3	B	1007	ARG	1	0
3	A	806	ARG	1	0
3	B	1006	ARG	5	0
3	A	807	ARG	7	0
2	B	1003	ACT	5	0
2	A	803	ACT	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	692/718 (96%)	-0.12	24 (3%) 47 49	16, 37, 78, 127	1 (0%)
1	B	700/718 (97%)	-0.17	13 (1%) 66 69	15, 37, 79, 118	1 (0%)
All	All	1392/1436 (96%)	-0.14	37 (2%) 56 59	15, 37, 79, 127	2 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	646	ALA	6.8
1	B	681	PHE	4.0
1	B	666	ILE	4.0
1	A	679	GLY	3.9
1	B	663	ALA	3.7
1	A	643	ILE	3.6
1	A	644	PRO	3.5
1	A	314	ILE	3.4
1	A	475	TYR	3.2
1	A	710	ILE	3.0
1	B	643	ILE	2.8
1	B	680	VAL	2.8
1	A	704	VAL	2.7
1	A	1	MET	2.6
1	A	714	THR	2.6
1	A	652	PHE	2.6
1	A	476	ILE	2.6
1	B	710	ILE	2.6
1	A	473	GLY	2.6
1	B	644	PRO	2.5
1	B	646	ALA	2.5
1	A	707	LEU	2.5
1	B	322	THR	2.4
1	B	314	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	315	HIS	2.4
1	A	666	ILE	2.4
1	B	316	GLY	2.3
1	A	680	VAL	2.3
1	B	321	SER	2.3
1	A	595	ILE	2.3
1	A	681	PHE	2.2
1	A	712	LYS	2.2
1	A	683	ILE	2.2
1	A	639	VAL	2.1
1	A	682	ASP	2.1
1	A	474	THR	2.1
1	A	663	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ACT	A	802	4/4	0.53	0.23	62,63,64,65	0
3	ARG	B	1007	12/12	0.63	0.22	78,88,94,97	0
2	ACT	A	801	4/4	0.65	0.23	54,54,56,59	0
2	ACT	A	803	4/4	0.71	0.19	60,63,64,67	0
2	ACT	B	1003	4/4	0.77	0.13	28,33,35,43	0
3	ARG	A	807	12/12	0.80	0.17	39,62,78,80	0
2	ACT	B	1004	4/4	0.80	0.16	83,85,85,85	0
3	ARG	B	1006	12/12	0.82	0.16	61,63,69,69	0
3	ARG	A	806	12/12	0.82	0.15	60,81,103,107	0
3	ARG	B	1005	12/12	0.84	0.13	50,54,59,59	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ACT	B	1001	4/4	0.85	0.14	48,53,53,56	0
3	ARG	A	805	12/12	0.86	0.12	53,56,58,59	0
2	ACT	A	804	4/4	0.87	0.14	71,75,75,77	0
2	ACT	B	1002	4/4	0.87	0.12	72,72,72,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.