



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:35 AM UTC

PDB ID : 2RG4 / pdb_00002rg4
Title : Crystal structure of the uncharacterized protein Q2CBJ1_9RHOB from Oceanicola granulosus HTCC2516
Authors : Malashkevich, V.N.; Toro, R.; Meyer, A.J.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2007-10-02
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	
1	B	216	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3621 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	206	Total	C	N	O	S	0	2	0
			1642	1050	276	309	7			
1	B	206	Total	C	N	O	S	0	0	0
			1630	1042	275	306	7			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q2CBJ1
A	2	SER	-	expression tag	UNP Q2CBJ1
A	3	LEU	-	expression tag	UNP Q2CBJ1
A	209	GLU	-	expression tag	UNP Q2CBJ1
A	210	GLY	-	expression tag	UNP Q2CBJ1
A	211	HIS	-	expression tag	UNP Q2CBJ1
A	212	HIS	-	expression tag	UNP Q2CBJ1
A	213	HIS	-	expression tag	UNP Q2CBJ1
A	214	HIS	-	expression tag	UNP Q2CBJ1
A	215	HIS	-	expression tag	UNP Q2CBJ1
A	216	HIS	-	expression tag	UNP Q2CBJ1
B	1	MET	-	expression tag	UNP Q2CBJ1
B	2	SER	-	expression tag	UNP Q2CBJ1
B	3	LEU	-	expression tag	UNP Q2CBJ1
B	209	GLU	-	expression tag	UNP Q2CBJ1
B	210	GLY	-	expression tag	UNP Q2CBJ1
B	211	HIS	-	expression tag	UNP Q2CBJ1
B	212	HIS	-	expression tag	UNP Q2CBJ1
B	213	HIS	-	expression tag	UNP Q2CBJ1
B	214	HIS	-	expression tag	UNP Q2CBJ1
B	215	HIS	-	expression tag	UNP Q2CBJ1
B	216	HIS	-	expression tag	UNP Q2CBJ1

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Fe 1	0	0
2	B	1	Total 1	Fe 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	197	Total 197	O 197	0	0
3	B	150	Total 150	O 150	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized protein



• Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	69.48Å 69.48Å 164.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.38 – 1.90 19.38 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.2 (19.38-1.90) 96.1 (19.38-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.78 (at 1.90Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.207 , 0.277 0.208 , 0.277	Depositor DCC
R_{free} test set	1584 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	20.5	Xtriage
Anisotropy	0.143	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3621	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.44	9/1697 (0.5%)	1.54	21/2311 (0.9%)
1	B	1.23	1/1679 (0.1%)	1.35	14/2287 (0.6%)
All	All	1.34	10/3376 (0.3%)	1.45	35/4598 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	5
All	All	0	9

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	105	ILE	CA-CB	8.02	1.64	1.55
1	A	23	GLY	N-CA	6.96	1.54	1.44
1	A	23	GLY	CA-C	-6.63	1.42	1.51
1	A	78	LYS	CA-C	6.59	1.61	1.52
1	B	80	LEU	N-CA	5.71	1.53	1.46
1	A	169	ARG	CA-C	5.70	1.59	1.52
1	A	175	VAL	C-O	5.43	1.29	1.24
1	A	167	THR	CA-CB	5.34	1.61	1.53
1	A	20	SER	C-O	-5.33	1.17	1.24
1	A	76	LEU	N-CA	-5.29	1.40	1.46

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	134	MET	CA-C-N	20.77	145.81	119.84
1	A	134	MET	C-N-CA	20.77	145.81	119.84
1	A	23	GLY	N-CA-C	-10.85	90.20	112.34
1	B	136	GLU	N-CA-C	-10.77	96.16	111.56
1	B	91	LEU	N-CA-C	9.72	124.94	113.18
1	A	135	PRO	N-CA-C	9.12	131.26	112.47
1	B	57	TYR	N-CA-CB	8.90	125.53	110.49
1	A	130	THR	CB-CA-C	-6.99	97.13	109.71
1	B	137	GLY	CA-C-N	6.97	134.25	121.70
1	B	137	GLY	C-N-CA	6.97	134.25	121.70
1	B	91	LEU	CA-C-N	6.93	134.18	121.70
1	B	91	LEU	C-N-CA	6.93	134.18	121.70
1	A	145	ASP	CA-C-N	6.83	126.52	119.56
1	A	145	ASP	C-N-CA	6.83	126.52	119.56
1	A	136	GLU	N-CA-C	-6.47	92.88	111.00
1	A	153	ALA	N-CA-C	-6.45	105.45	113.38
1	A	55	PRO	CA-C-N	-6.37	108.93	121.41
1	A	55	PRO	C-N-CA	-6.37	108.93	121.41
1	B	57	TYR	CA-CB-CG	-6.34	102.48	113.90
1	A	134	MET	C-N-CD	-6.22	99.50	125.00
1	A	135	PRO	CA-C-N	6.15	132.76	121.70
1	A	135	PRO	C-N-CA	6.15	132.76	121.70
1	B	148	LEU	CA-C-N	-5.95	113.68	119.87
1	B	148	LEU	C-N-CA	-5.95	113.68	119.87
1	A	59	SER	CA-C-N	5.87	128.15	120.28
1	A	59	SER	C-N-CA	5.87	128.15	120.28
1	A	83	HIS	N-CA-C	-5.78	104.90	111.14
1	B	69	ARG	N-CA-C	5.51	118.46	111.69
1	B	73	PHE	N-CA-C	5.43	117.28	111.36
1	A	30	GLU	N-CA-C	-5.25	105.55	111.28
1	A	121	HIS	CA-C-N	5.21	125.22	119.90
1	A	121	HIS	C-N-CA	5.21	125.22	119.90
1	B	92	GLU	N-CA-C	-5.20	96.44	111.00
1	A	188	GLU	N-CA-CB	-5.20	101.87	111.53
1	B	138	THR	N-CA-CB	5.20	120.33	111.50

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	134	MET	Peptide
1	A	136	GLU	Peptide
1	A	22	HIS	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	56	GLY	Peptide
1	B	135	PRO	Peptide
1	B	136	GLU	Peptide
1	B	137	GLY	Peptide
1	B	56	GLY	Peptide
1	B	91	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1642	0	1572	40	0
1	B	1630	0	1558	48	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	197	0	0	14	0
3	B	150	0	0	4	1
All	All	3621	0	3130	85	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:164:GLU:HB2	3:B:332:HOH:O	1.42	1.16
1:A:148:LEU:HA	1:A:151:MET:HE2	1.18	1.08
1:A:148:LEU:HA	1:A:151:MET:CE	1.93	0.99
1:A:119:HIS:CE1	3:A:410:HOH:O	2.13	0.98
1:B:148:LEU:HD13	1:B:167:THR:CG2	1.96	0.94
1:B:121:HIS:CE1	3:B:243:HOH:O	2.23	0.89
1:B:158:ARG:O	1:B:160:GLY:N	2.04	0.88
1:B:161:ALA:HB2	1:B:166:ARG:HH11	1.39	0.86
1:B:148:LEU:HD13	1:B:167:THR:HG21	1.56	0.86
1:B:41:ASP:HB2	3:B:338:HOH:O	1.76	0.85
1:A:163[A]:GLU:HG2	3:A:231:HOH:O	1.77	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:163:GLU:HG3	1:B:169:ARG:HD2	1.58	0.83
1:A:148:LEU:CA	1:A:151:MET:HE2	2.04	0.82
1:B:158:ARG:HB2	1:B:158:ARG:HH11	1.46	0.79
1:A:78:LYS:HA	1:A:78:LYS:HE3	1.65	0.78
1:B:13:ARG:HG3	1:B:13:ARG:HH11	1.50	0.77
1:B:166:ARG:O	1:B:169:ARG:NH1	2.22	0.72
1:A:118:SER:OG	1:A:186:ARG:HD3	1.93	0.69
1:B:148:LEU:HD13	1:B:167:THR:HG23	1.74	0.69
1:B:158:ARG:C	1:B:160:GLY:H	2.01	0.68
1:A:114:GLY:O	1:A:191:MET:HA	1.95	0.66
1:B:196:GLU:HG3	3:B:323:HOH:O	2.00	0.62
1:B:161:ALA:HB2	1:B:166:ARG:NH1	2.15	0.61
1:A:119:HIS:ND1	3:A:410:HOH:O	2.27	0.61
1:A:5[A]:GLN:HG3	3:A:262:HOH:O	1.99	0.61
1:B:55:PRO:HG2	1:B:115:VAL:O	2.01	0.59
1:B:161:ALA:HA	1:B:162:ARG:C	2.27	0.59
1:B:158:ARG:HB2	1:B:158:ARG:NH1	2.16	0.59
1:A:5[A]:GLN:NE2	3:A:262:HOH:O	2.35	0.59
1:A:39:ALA:O	1:A:46:GLN:NE2	2.36	0.58
1:A:5[A]:GLN:HG2	1:A:7:LYS:HD3	1.85	0.58
1:B:164:GLU:HA	1:B:169:ARG:NH2	2.19	0.57
1:A:198:ARG:HD2	3:A:240:HOH:O	2.04	0.56
1:B:148:LEU:CD1	1:B:167:THR:HG21	2.33	0.56
1:B:48:TRP:CE2	1:B:52:GLU:HG3	2.42	0.54
1:B:135:PRO:CB	1:B:136:GLU:HB3	2.37	0.54
1:B:59:SER:HB2	1:B:63:LEU:HD11	1.90	0.54
1:A:198:ARG:CD	3:A:240:HOH:O	2.54	0.54
1:B:100:LEU:HD23	1:B:207:TRP:HA	1.89	0.54
1:A:56:GLY:HA3	3:A:273:HOH:O	2.07	0.53
1:A:118:SER:OG	1:A:186:ARG:CG	2.57	0.52
1:A:125:VAL:HG12	1:A:126:ILE:HG12	1.91	0.52
3:A:354:HOH:O	1:B:123:HIS:CG	2.62	0.51
1:B:13:ARG:HG3	1:B:13:ARG:NH1	2.22	0.51
1:B:162:ARG:C	1:B:164:GLU:H	2.18	0.51
1:B:59:SER:HB2	1:B:63:LEU:CD1	2.41	0.50
1:B:135:PRO:HB3	1:B:136:GLU:HB3	1.94	0.50
1:A:43:ASP:OD2	3:A:267:HOH:O	2.20	0.50
1:A:189:VAL:HG13	1:A:198:ARG:NH1	2.28	0.49
1:B:38:ILE:HG12	1:B:70:PHE:CZ	2.46	0.49
1:B:118:SER:OG	1:B:186:ARG:HD3	2.12	0.49
1:A:105:ILE:HG13	1:A:203:PHE:HB3	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:LEU:HD12	1:A:179:LEU:HD13	1.94	0.49
1:B:13:ARG:NH1	1:B:90:ASP:OD2	2.46	0.49
1:A:134:MET:N	1:A:135:PRO:HD2	2.24	0.48
1:A:119:HIS:HE1	3:A:410:HOH:O	1.67	0.48
1:A:131:TYR:HD1	1:A:134:MET:HE3	1.78	0.48
3:A:354:HOH:O	1:B:123:HIS:CD2	2.66	0.48
1:A:118:SER:OG	1:A:186:ARG:CD	2.59	0.47
1:A:173:PRO:HG3	1:A:179:LEU:HD11	1.97	0.47
1:B:140:ALA:O	1:B:190:PRO:HD2	2.15	0.47
1:A:103:GLU:HG3	3:A:330:HOH:O	2.13	0.46
1:B:57:TYR:CD1	1:B:57:TYR:C	2.92	0.46
1:A:148:LEU:N	1:A:149:PRO:HD2	2.32	0.45
1:A:36:TYR:CZ	1:A:199:ILE:HD11	2.51	0.45
1:B:135:PRO:HG3	1:B:196:GLU:HB2	1.97	0.45
1:A:51:ARG:NH2	3:A:312:HOH:O	2.39	0.45
1:B:163:GLU:HG3	1:B:169:ARG:CD	2.38	0.45
1:A:131:TYR:HD1	1:A:134:MET:CE	2.30	0.45
1:B:171:VAL:O	1:B:173:PRO:HD3	2.18	0.44
1:A:116:HIS:HB2	1:A:189:VAL:HB	2.00	0.44
1:A:48:TRP:O	1:A:52:GLU:HB2	2.18	0.44
1:B:135:PRO:CA	1:B:136:GLU:HB3	2.48	0.43
1:B:143:LEU:O	1:B:168:PHE:HA	2.17	0.43
1:A:23:GLY:HA3	1:A:79:SER:HA	2.00	0.43
1:B:73:PHE:O	1:B:77:VAL:HG23	2.19	0.43
1:A:123:HIS:HD2	1:B:152:MET:O	2.03	0.42
1:B:136:GLU:O	1:B:137:GLY:C	2.63	0.41
1:B:164:GLU:HA	1:B:169:ARG:HH21	1.83	0.41
1:A:186:ARG:NH1	1:B:186:ARG:HD3	2.35	0.41
1:A:5[A]:GLN:HG2	1:A:7:LYS:CD	2.51	0.41
1:A:163[A]:GLU:HG2	1:A:163[A]:GLU:H	1.74	0.41
1:A:152:MET:O	1:B:123:HIS:HD2	2.03	0.40
1:B:58:THR:HA	1:B:107:ILE:O	2.21	0.40
1:B:134:MET:O	1:B:135:PRO:C	2.64	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:182:GLU:OE2	3:B:283:HOH:O[7_556]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/216 (95%)	199 (97%)	6 (3%)	1 (0%)	24	16
1	B	204/216 (94%)	189 (93%)	11 (5%)	4 (2%)	6	1
All	All	410/432 (95%)	388 (95%)	17 (4%)	5 (1%)	10	4

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	135	PRO
1	B	138	THR
1	B	159	LYS
1	B	137	GLY
1	B	160	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/178 (97%)	157 (91%)	15 (9%)	9	4
1	B	170/178 (96%)	154 (91%)	16 (9%)	8	3
All	All	342/356 (96%)	311 (91%)	31 (9%)	9	3

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	78	LYS
1	A	115	VAL
1	A	130	THR
1	A	134	MET
1	A	135	PRO
1	A	141	LEU
1	A	157	ARG
1	A	159	LYS
1	A	163[A]	GLU
1	A	163[B]	GLU
1	A	164	GLU
1	A	174	LYS
1	A	175	VAL
1	A	195	GLU
1	B	3	LEU
1	B	37	SER
1	B	50	GLU
1	B	63	LEU
1	B	92	GLU
1	B	94	GLU
1	B	95	LEU
1	B	98	LYS
1	B	138	THR
1	B	143	LEU
1	B	156	THR
1	B	158	ARG
1	B	159	LYS
1	B	162	ARG
1	B	174	LYS
1	B	193	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	123	HIS
1	B	116	HIS
1	B	123	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	206/216 (95%)	-0.03	8 (3%)	43	46	11, 22, 43, 71	2 (0%)
1	B	206/216 (95%)	0.26	9 (4%)	39	41	15, 28, 55, 77	0
All	All	412/432 (95%)	0.11	17 (4%)	41	44	11, 25, 51, 77	2 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	137	GLY	5.4
1	B	161	ALA	4.2
1	B	163	GLU	3.9
1	B	137	GLY	3.7
1	B	136	GLU	3.1
1	A	56	GLY	2.8
1	B	158	ARG	2.7
1	A	136	GLU	2.7
1	A	135	PRO	2.7
1	A	138	THR	2.6
1	B	160	GLY	2.4
1	A	23	GLY	2.3
1	B	165	LEU	2.3
1	A	68	TRP	2.2
1	B	63	LEU	2.2
1	A	97	GLY	2.1
1	B	169	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FE	A	217	1/1	0.99	0.05	32,32,32,32	0
2	FE	B	217	1/1	0.99	0.04	40,40,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.