



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:06 AM UTC

PDB ID : 2RGY / pdb_00002rgy
Title : Crystal structure of transcriptional regulator of LacI family from Burkholderia phymatum
Authors : Agarwal, R.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2007-10-05
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	3% 71% 22% • 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator, LacI family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	Se	0	0	0
			2122	1344	370	398	6	4			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	78	MSE	-	expression tag	UNP A0FW12
A	79	SER	-	expression tag	UNP A0FW12
A	80	LEU	-	expression tag	UNP A0FW12
A	360	GLU	-	expression tag	UNP A0FW12
A	361	GLY	-	expression tag	UNP A0FW12
A	362	HIS	-	expression tag	UNP A0FW12
A	363	HIS	-	expression tag	UNP A0FW12
A	364	HIS	-	expression tag	UNP A0FW12
A	365	HIS	-	expression tag	UNP A0FW12
A	366	HIS	-	expression tag	UNP A0FW12
A	367	HIS	-	expression tag	UNP A0FW12


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	174	Total	O	0	0
			174	174		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional regulator, LacI family



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	58.54Å 118.78Å 85.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.72 – 2.05 42.72 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.7 (42.72-2.05) 97.7 (42.72-2.05)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.41 (at 2.05Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.210 , 0.267 0.226 , 0.269	Depositor DCC
R_{free} test set	552 reflections (2.87%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2296	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	3/2168 (0.1%)	0.97	13/2941 (0.4%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	170	MSE	SE-CE	-9.44	1.67	1.95
1	A	353	MSE	SE-CE	-6.12	1.77	1.95
1	A	277	MSE	SE-CE	-5.18	1.79	1.95

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	306	SER	N-CA-C	-7.11	104.63	113.38
1	A	342	GLU	N-CA-C	-6.67	99.67	110.20
1	A	183	ALA	N-CA-C	-5.94	105.80	113.16
1	A	314	LEU	N-CA-C	5.65	117.90	109.25
1	A	150	VAL	N-CA-C	5.63	116.23	108.12
1	A	117	HIS	N-CA-C	5.46	118.17	110.14
1	A	337	TYR	N-CA-C	5.34	119.77	112.88
1	A	241	VAL	N-CA-C	5.29	113.69	107.77
1	A	229	ASP	N-CA-C	-5.28	105.44	111.14
1	A	174	ASN	N-CA-C	5.25	119.27	112.86
1	A	336	CYS	N-CA-C	5.19	117.73	111.40
1	A	305	TYR	N-CA-C	5.04	119.01	113.21
1	A	145	CYS	N-CA-C	-5.01	103.53	110.35

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2122	0	2062	35	0
2	A	174	0	0	5	0
All	All	2296	0	2062	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:ASP:HB2	1:A:345:ARG:NH2	1.93	0.84
1:A:182:ASP:HB3	2:A:374:HOH:O	1.89	0.73
1:A:131:GLU:O	1:A:135:GLU:HG3	1.91	0.71
1:A:210:VAL:HB	1:A:243:LEU:HD22	1.75	0.68
1:A:86:GLY:O	1:A:87:ILE:HD13	1.93	0.67
1:A:85:LEU:HG	1:A:87:ILE:HG12	1.76	0.66
1:A:213:GLY:H	1:A:220:ASN:ND2	1.96	0.64
1:A:87:ILE:HG21	1:A:119:VAL:HG23	1.81	0.63
1:A:165:ARG:HD3	2:A:427:HOH:O	1.99	0.61
1:A:98:SER:OG	1:A:304:ASP:HB3	2.01	0.59
1:A:88:ILE:HD12	1:A:111:LEU:HD21	1.83	0.59
1:A:87:ILE:HD12	1:A:117:HIS:HB2	1.85	0.58
1:A:331:TRP:O	1:A:335:GLN:HG2	2.06	0.56
1:A:213:GLY:H	1:A:220:ASN:HD21	1.53	0.55
1:A:162:GLU:HG3	2:A:427:HOH:O	2.06	0.55
1:A:100:TYR:O	1:A:104:LEU:HG	2.09	0.53
1:A:86:GLY:C	1:A:87:ILE:HD13	2.34	0.52
1:A:301:TYR:O	1:A:302:ASP:CB	2.56	0.52
1:A:247:ASP:O	1:A:248:PHE:HB2	2.10	0.52
1:A:334:ASN:HD21	1:A:341:TRP:H	1.56	0.52
1:A:111:LEU:HD12	1:A:118:VAL:HB	1.94	0.49
1:A:212:SER:HA	1:A:273:ALA:HB2	1.94	0.49
1:A:263:SER:O	1:A:264:LYS:HB2	2.13	0.49
1:A:301:TYR:O	1:A:302:ASP:HB2	2.12	0.48
1:A:283:ALA:O	1:A:287:GLN:HG3	2.14	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:ALA:O	1:A:202:GLU:HG3	2.15	0.46
1:A:116:ARG:HD3	1:A:337:TYR:CZ	2.51	0.46
1:A:199:THR:O	1:A:203:HIS:HD2	1.99	0.46
1:A:223:ARG:CZ	2:A:438:HOH:O	2.64	0.45
1:A:182:ASP:HB2	1:A:345:ARG:HH22	1.76	0.44
1:A:191:ARG:HG3	1:A:349:VAL:HG11	2.00	0.44
1:A:251:GLU:HG2	1:A:252:GLY:N	2.33	0.43
1:A:293:PRO:HB2	1:A:358:ALA:HB3	2.00	0.43
1:A:246:SER:HB3	1:A:277:MSE:SE	2.69	0.42
1:A:276:THR:HA	2:A:511:HOH:O	2.18	0.42

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	273/290 (94%)	261 (96%)	11 (4%)	1 (0%)	30 22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	302	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	224/232 (97%)	220 (98%)	4 (2%)	51 51

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	134	LEU
1	A	243	LEU
1	A	340	LYS
1	A	359	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	106	GLN
1	A	164	HIS
1	A	203	HIS
1	A	220	ASN
1	A	334	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	271/290 (93%)	0.22	8 (2%) 52 53	15, 23, 33, 48	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	305	TYR	6.9
1	A	304	ASP	4.6
1	A	306	SER	4.3
1	A	182	ASP	2.4
1	A	239	ASP	2.3
1	A	126	GLU	2.3
1	A	222	GLU	2.2
1	A	165	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.