



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 09:14 AM UTC

PDB ID : 3RGD / pdb_00003rgd
Title : Iron loaded frog M ferritin. Short soaking time
Authors : Bertini, I.; Lalli, D.; Mangani, S.; Pozzi, C.; Rosa, C.; Theil, E.C.; Turano, P.
Deposited on : 2011-04-08
Resolution : 2.89 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	 72% 23% . .
1	B	176	 73% 21% . .
1	C	176	 68% 26% . .
1	D	176	 66% 28% . .
1	E	176	 72% 23% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	176	 77% 18% . .
1	G	176	 74% 20% . .
1	H	176	 73% 19% 5% .
1	I	176	 66% 26% 5% .
1	J	176	 71% 23% . .
1	K	176	 65% 26% 6% . .
1	L	176	 68% 27% . .
1	M	176	 71% 24% . .
1	N	176	 71% 22% . .
1	O	176	 75% 19% . .
1	P	176	 67% 29% . .
1	Q	176	 64% 28% 6% .
1	R	176	 73% 22% . .
1	S	176	 71% 23% . .
1	T	176	 70% 24% . .
1	U	176	 % 77% 19% . .
1	V	176	 70% 24% . .
1	W	176	 74% 20% . .
1	X	176	 72% 22% . . .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 34941 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferritin, middle subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	B	173	Total	C	N	O	S	0	0	0
			1427	897	248	275	7			
1	C	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	D	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	E	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	F	171	Total	C	N	O	S	0	0	0
			1409	886	245	271	7			
1	G	171	Total	C	N	O	S	0	0	0
			1409	886	245	271	7			
1	H	171	Total	C	N	O	S	0	0	0
			1409	886	245	271	7			
1	I	171	Total	C	N	O	S	0	0	0
			1409	886	245	271	7			
1	J	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	K	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	L	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	M	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	N	171	Total	C	N	O	S	0	0	0
			1409	886	245	271	7			
1	O	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	P	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	R	171	Total	C	N	O	S	0	0	0
			1409	886	245	271	7			
1	S	171	Total	C	N	O	S	0	0	0
			1409	886	245	271	7			
1	T	171	Total	C	N	O	S	0	0	0
			1409	886	245	271	7			
1	U	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	V	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	W	172	Total	C	N	O	S	0	0	0
			1418	892	247	272	7			
1	X	171	Total	C	N	O	S	0	0	0
			1409	886	245	271	7			

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Fe	0	0
			2	2		
2	B	3	Total	Fe	0	0
			3	3		
2	C	4	Total	Fe	0	0
			4	4		
2	D	4	Total	Fe	0	0
			4	4		
2	E	3	Total	Fe	0	0
			3	3		
2	F	3	Total	Fe	0	0
			3	3		
2	G	2	Total	Fe	0	0
			2	2		
2	H	3	Total	Fe	0	0
			3	3		
2	I	3	Total	Fe	0	0
			3	3		
2	J	3	Total	Fe	0	0
			3	3		
2	K	3	Total	Fe	0	0
			3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	L	2	Total 2	Fe 2	0	0
2	M	2	Total 2	Fe 2	0	0
2	N	3	Total 3	Fe 3	0	0
2	O	3	Total 3	Fe 3	0	0
2	P	2	Total 2	Fe 2	0	0
2	Q	2	Total 2	Fe 2	0	0
2	R	2	Total 2	Fe 2	0	0
2	S	2	Total 2	Fe 2	0	0
2	T	3	Total 3	Fe 3	0	0
2	U	2	Total 2	Fe 2	0	0
2	V	2	Total 2	Fe 2	0	0
2	W	2	Total 2	Fe 2	0	0
2	X	2	Total 2	Fe 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	36	Total 36	O 36	0	0
3	B	50	Total 50	O 50	0	0
3	C	32	Total 32	O 32	0	0
3	D	54	Total 54	O 54	0	0
3	E	46	Total 46	O 46	0	0
3	F	32	Total 32	O 32	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	29	Total 29	O 29	0	0
3	H	43	Total 43	O 43	0	0
3	I	46	Total 46	O 46	0	0
3	J	36	Total 36	O 36	0	0
3	K	45	Total 45	O 45	0	0
3	L	49	Total 49	O 49	0	0
3	M	30	Total 30	O 30	0	0
3	N	22	Total 22	O 22	0	0
3	O	34	Total 34	O 34	0	0
3	P	48	Total 48	O 48	0	0
3	Q	24	Total 24	O 24	0	0
3	R	36	Total 36	O 36	0	0
3	S	30	Total 30	O 30	0	0
3	T	35	Total 35	O 35	0	0
3	U	32	Total 32	O 32	0	0
3	V	43	Total 43	O 43	0	0
3	W	48	Total 48	O 48	0	0
3	X	39	Total 39	O 39	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ferritin, middle subunit




- Molecule 1: Ferritin, middle subunit



- Molecule 1: Ferritin, middle subunit



- Molecule 1: Ferritin, middle subunit





- Molecule 1: Ferritin, middle subunit

Chain E: 72% 23% . .



- Molecule 1: Ferritin, middle subunit

Chain F: 77% 18% . .



- Molecule 1: Ferritin, middle subunit

Chain G: 74% 20% . .



- Molecule 1: Ferritin, middle subunit

Chain H: 73% 19% 5% .



- Molecule 1: Ferritin, middle subunit


Chain I: 66% 26% 5% .






- Molecule 1: Ferritin, middle subunit

Chain J: 71% 23% . .




- Molecule 1: Ferritin, middle subunit

Chain K: 65% 26% 6% . .




- Molecule 1: Ferritin, middle subunit

Chain L: 68% 27% . .



- Molecule 1: Ferritin, middle subunit


Chain M: 71% 24% . .



- Molecule 1: Ferritin, middle subunit

Chain N: 71% 22% . .






- Molecule 1: Ferritin, middle subunit

Chain O: 75% 19%




- Molecule 1: Ferritin, middle subunit

Chain P: 67% 29%




- Molecule 1: Ferritin, middle subunit

Chain Q: 64% 28% 6%




- Molecule 1: Ferritin, middle subunit


Chain R: 73% 22%



- Molecule 1: Ferritin, middle subunit

Chain S: 71% 23%






- Molecule 1: Ferritin, middle subunit




- Molecule 1: Ferritin, middle subunit



- Molecule 1: Ferritin, middle subunit




- Molecule 1: Ferritin, middle subunit



- Molecule 1: Ferritin, middle subunit





4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	210.74Å 210.74Å 322.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.54 – 2.89 48.54 – 2.89	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.54-2.89) 99.7 (48.54-2.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.45 (at 2.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.172 , 0.229 0.172 , 0.227	Depositor DCC
R_{free} test set	9240 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	33.2	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.046 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	34941	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/1446	0.83	0/1944
1	B	0.55	0/1455	0.79	0/1956
1	C	0.50	0/1446	0.82	1/1944 (0.1%)
1	D	0.54	0/1446	0.85	4/1944 (0.2%)
1	E	0.60	0/1446	0.85	4/1944 (0.2%)
1	F	0.53	0/1437	0.78	0/1933
1	G	0.52	0/1437	0.84	2/1933 (0.1%)
1	H	0.55	0/1437	0.83	2/1933 (0.1%)
1	I	0.54	0/1437	0.83	0/1933
1	J	0.49	0/1446	0.79	0/1944
1	K	0.53	0/1446	0.82	1/1944 (0.1%)
1	L	0.64	0/1446	0.88	2/1944 (0.1%)
1	M	0.50	0/1446	0.77	0/1944
1	N	0.52	0/1437	0.78	0/1933
1	O	0.54	0/1446	0.82	0/1944
1	P	0.53	0/1446	0.76	0/1944
1	Q	0.51	0/1446	0.77	1/1944 (0.1%)
1	R	0.49	0/1437	0.76	0/1933
1	S	0.49	0/1437	0.80	1/1933 (0.1%)
1	T	0.52	0/1437	0.82	1/1933 (0.1%)
1	U	0.53	0/1446	0.79	1/1944 (0.1%)
1	V	0.56	0/1446	0.85	0/1944
1	W	0.52	0/1446	0.81	0/1944
1	X	0.71	4/1437 (0.3%)	0.87	2/1933 (0.1%)
All	All	0.54	4/34632 (0.0%)	0.81	22/46569 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	44	ALA	CA-CB	-6.13	1.43	1.53
1	X	45	LEU	C-O	-5.95	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	43	VAL	CA-CB	-5.57	1.49	1.55
1	X	134	TYR	C-O	-5.03	1.17	1.24

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	135	LEU	N-CA-C	7.67	119.72	111.36
1	X	44	ALA	N-CA-C	7.14	121.25	111.39
1	C	43	VAL	N-CA-C	-6.17	106.94	112.12
1	D	84	LYS	CA-C-N	6.17	126.18	119.89
1	D	84	LYS	C-N-CA	6.17	126.18	119.89

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1418	0	1370	31	0
1	B	1427	0	1376	29	0
1	C	1418	0	1370	35	0
1	D	1418	0	1370	37	0
1	E	1418	0	1370	28	0
1	F	1409	0	1357	25	0
1	G	1409	0	1357	21	0
1	H	1409	0	1357	31	0
1	I	1409	0	1357	33	0
1	J	1418	0	1370	41	0
1	K	1418	0	1370	46	0
1	L	1418	0	1370	32	0
1	M	1418	0	1370	34	0
1	N	1409	0	1357	37	0
1	O	1418	0	1370	27	0
1	P	1418	0	1370	35	0
1	Q	1418	0	1370	50	0
1	R	1409	0	1357	28	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	1409	0	1357	28	0
1	T	1409	0	1357	32	0
1	U	1418	0	1370	25	0
1	V	1418	0	1370	39	0
1	W	1418	0	1370	30	0
1	X	1409	0	1357	39	0
2	A	2	0	0	0	0
2	B	3	0	0	0	0
2	C	4	0	0	0	0
2	D	4	0	0	0	0
2	E	3	0	0	0	0
2	F	3	0	0	0	0
2	G	2	0	0	0	0
2	H	3	0	0	0	0
2	I	3	0	0	0	0
2	J	3	0	0	0	0
2	K	3	0	0	0	0
2	L	2	0	0	0	0
2	M	2	0	0	0	0
2	N	3	0	0	0	0
2	O	3	0	0	0	0
2	P	2	0	0	0	0
2	Q	2	0	0	0	0
2	R	2	0	0	0	0
2	S	2	0	0	0	0
2	T	3	0	0	0	0
2	U	2	0	0	0	0
2	V	2	0	0	0	0
2	W	2	0	0	0	0
2	X	2	0	0	0	0
3	A	36	0	0	4	0
3	B	50	0	0	3	0
3	C	32	0	0	1	0
3	D	54	0	0	5	0
3	E	46	0	0	3	0
3	F	32	0	0	1	0
3	G	29	0	0	1	0
3	H	43	0	0	1	0
3	I	46	0	0	1	0
3	J	36	0	0	6	0
3	K	45	0	0	2	0
3	L	49	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	M	30	0	0	3	0
3	N	22	0	0	2	0
3	O	34	0	0	2	0
3	P	48	0	0	1	0
3	Q	24	0	0	1	0
3	R	36	0	0	2	0
3	S	30	0	0	4	0
3	T	35	0	0	3	0
3	U	32	0	0	1	0
3	V	43	0	0	1	0
3	W	48	0	0	4	0
3	X	39	0	0	5	0
All	All	34941	0	32769	667	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 667 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:20:MET:HE1	1:D:110:ALA:HB1	1.32	1.06
1:A:142:ILE:HG22	1:X:5:VAL:HG22	1.44	1.00
1:B:5:VAL:HG22	1:V:142:ILE:HG22	1.39	0.99
1:C:142:ILE:HG22	1:S:5:VAL:HG22	1.40	0.99
1:V:20:MET:HE1	1:V:110:ALA:HB1	1.45	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	170/176 (97%)	167 (98%)	3 (2%)	0	100	100
1	B	171/176 (97%)	166 (97%)	4 (2%)	1 (1%)	21	51
1	C	170/176 (97%)	165 (97%)	5 (3%)	0	100	100
1	D	170/176 (97%)	165 (97%)	5 (3%)	0	100	100
1	E	170/176 (97%)	166 (98%)	4 (2%)	0	100	100
1	F	169/176 (96%)	163 (96%)	6 (4%)	0	100	100
1	G	169/176 (96%)	162 (96%)	7 (4%)	0	100	100
1	H	169/176 (96%)	166 (98%)	3 (2%)	0	100	100
1	I	169/176 (96%)	162 (96%)	7 (4%)	0	100	100
1	J	170/176 (97%)	166 (98%)	4 (2%)	0	100	100
1	K	170/176 (97%)	164 (96%)	6 (4%)	0	100	100
1	L	170/176 (97%)	164 (96%)	6 (4%)	0	100	100
1	M	170/176 (97%)	162 (95%)	8 (5%)	0	100	100
1	N	169/176 (96%)	163 (96%)	6 (4%)	0	100	100
1	O	170/176 (97%)	161 (95%)	9 (5%)	0	100	100
1	P	170/176 (97%)	161 (95%)	9 (5%)	0	100	100
1	Q	170/176 (97%)	160 (94%)	10 (6%)	0	100	100
1	R	169/176 (96%)	163 (96%)	6 (4%)	0	100	100
1	S	169/176 (96%)	162 (96%)	7 (4%)	0	100	100
1	T	169/176 (96%)	165 (98%)	4 (2%)	0	100	100
1	U	170/176 (97%)	164 (96%)	6 (4%)	0	100	100
1	V	170/176 (97%)	166 (98%)	4 (2%)	0	100	100
1	W	170/176 (97%)	165 (97%)	5 (3%)	0	100	100
1	X	169/176 (96%)	164 (97%)	5 (3%)	0	100	100
All	All	4072/4224 (96%)	3932 (97%)	139 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	173	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/157 (98%)	138 (90%)	15 (10%)	7	25
1	B	154/157 (98%)	137 (89%)	17 (11%)	6	20
1	C	153/157 (98%)	135 (88%)	18 (12%)	5	17
1	D	153/157 (98%)	137 (90%)	16 (10%)	6	22
1	E	153/157 (98%)	138 (90%)	15 (10%)	7	25
1	F	152/157 (97%)	134 (88%)	18 (12%)	5	17
1	G	152/157 (97%)	133 (88%)	19 (12%)	4	15
1	H	152/157 (97%)	138 (91%)	14 (9%)	8	27
1	I	152/157 (97%)	129 (85%)	23 (15%)	3	10
1	J	153/157 (98%)	137 (90%)	16 (10%)	6	22
1	K	153/157 (98%)	136 (89%)	17 (11%)	6	20
1	L	153/157 (98%)	137 (90%)	16 (10%)	6	22
1	M	153/157 (98%)	137 (90%)	16 (10%)	6	22
1	N	152/157 (97%)	136 (90%)	16 (10%)	6	22
1	O	153/157 (98%)	139 (91%)	14 (9%)	8	27
1	P	153/157 (98%)	141 (92%)	12 (8%)	11	35
1	Q	153/157 (98%)	140 (92%)	13 (8%)	10	31
1	R	152/157 (97%)	137 (90%)	15 (10%)	7	24
1	S	152/157 (97%)	135 (89%)	17 (11%)	6	19
1	T	152/157 (97%)	138 (91%)	14 (9%)	8	27
1	U	153/157 (98%)	141 (92%)	12 (8%)	11	35
1	V	153/157 (98%)	135 (88%)	18 (12%)	5	17
1	W	153/157 (98%)	138 (90%)	15 (10%)	7	25
1	X	152/157 (97%)	134 (88%)	18 (12%)	5	17
All	All	3664/3768 (97%)	3280 (90%)	384 (10%)	6	22

5 of 384 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	N	122	VAL
1	R	116	LYS
1	O	23	LEU
1	P	152	LEU
1	S	135	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 121 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	55	HIS
1	W	22	ASN
1	O	102	GLN
1	V	138	GLN
1	X	80	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 62 ligands modelled in this entry, 62 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	172/176 (97%)	-0.77	0	100	100	19, 27, 40, 52	0
1	B	173/176 (98%)	-0.80	0	100	100	20, 26, 40, 62	0
1	C	172/176 (97%)	-0.68	0	100	100	24, 31, 42, 54	0
1	D	172/176 (97%)	-0.83	0	100	100	17, 25, 39, 46	0
1	E	172/176 (97%)	-0.86	0	100	100	19, 25, 37, 44	0
1	F	171/176 (97%)	-0.82	0	100	100	20, 27, 39, 47	0
1	G	171/176 (97%)	-0.83	0	100	100	19, 25, 37, 47	0
1	H	171/176 (97%)	-0.88	0	100	100	19, 25, 35, 46	0
1	I	171/176 (97%)	-0.88	0	100	100	18, 24, 35, 46	0
1	J	172/176 (97%)	-0.64	0	100	100	24, 32, 47, 54	0
1	K	172/176 (97%)	-0.77	0	100	100	18, 26, 40, 50	0
1	L	172/176 (97%)	-0.81	0	100	100	16, 25, 39, 47	0
1	M	172/176 (97%)	-0.74	0	100	100	24, 30, 43, 53	0
1	N	171/176 (97%)	-0.67	0	100	100	24, 31, 43, 53	0
1	O	172/176 (97%)	-0.72	0	100	100	22, 29, 43, 55	0
1	P	172/176 (97%)	-0.88	0	100	100	19, 24, 38, 50	0
1	Q	172/176 (97%)	-0.70	0	100	100	22, 30, 42, 49	0
1	R	171/176 (97%)	-0.64	0	100	100	26, 32, 44, 53	0
1	S	171/176 (97%)	-0.66	0	100	100	23, 32, 44, 54	0
1	T	171/176 (97%)	-0.79	0	100	100	17, 26, 40, 43	0
1	U	172/176 (97%)	-0.72	1 (0%)	85	81	20, 28, 40, 51	0
1	V	172/176 (97%)	-0.82	0	100	100	17, 25, 40, 47	0
1	W	172/176 (97%)	-0.85	0	100	100	19, 26, 39, 48	0
1	X	171/176 (97%)	-0.70	0	100	100	23, 29, 42, 51	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	4120/4224 (97%)	-0.77	1 (0%) 100 100	16, 28, 41, 62	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	U	173	LYS	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FE	D	183	1/1	0.76	0.09	54,54,54,54	0
2	FE	K	183	1/1	0.82	0.08	55,55,55,55	0
2	FE	F	183	1/1	0.84	0.13	52,52,52,52	0
2	FE	H	183	1/1	0.89	0.12	56,56,56,56	0
2	FE	C	183	1/1	0.89	0.14	61,61,61,61	0
2	FE	D	181	1/1	0.90	0.20	63,63,63,63	0
2	FE	B	183	1/1	0.91	0.08	59,59,59,59	0
2	FE	O	181	1/1	0.91	0.16	72,72,72,72	0
2	FE	Q	181	1/1	0.91	0.13	69,69,69,69	0
2	FE	J	181	1/1	0.92	0.14	67,67,67,67	0
2	FE	S	181	1/1	0.92	0.10	64,64,64,64	0
2	FE	W	181	1/1	0.92	0.12	55,55,55,55	0
2	FE	E	181	1/1	0.93	0.15	60,60,60,60	0
2	FE	R	181	1/1	0.93	0.14	71,71,71,71	0
2	FE	J	183	1/1	0.93	0.10	60,60,60,60	0
2	FE	P	181	1/1	0.93	0.15	68,68,68,68	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FE	R	180	1/1	0.94	0.20	49,49,49,49	0
2	FE	G	180	1/1	0.94	0.18	52,52,52,52	0
2	FE	F	181	1/1	0.94	0.19	59,59,59,59	0
2	FE	T	183	1/1	0.94	0.12	56,56,56,56	0
2	FE	H	181	1/1	0.94	0.15	56,56,56,56	0
2	FE	B	180	1/1	0.95	0.15	56,56,56,56	0
2	FE	M	181	1/1	0.95	0.18	66,66,66,66	0
2	FE	N	181	1/1	0.95	0.14	73,73,73,73	0
2	FE	G	181	1/1	0.95	0.15	64,64,64,64	0
2	FE	P	180	1/1	0.95	0.19	50,50,50,50	0
2	FE	T	180	1/1	0.95	0.17	39,39,39,39	0
2	FE	U	181	1/1	0.95	0.17	61,61,61,61	0
2	FE	A	181	1/1	0.95	0.16	56,56,56,56	0
2	FE	X	180	1/1	0.95	0.19	51,51,51,51	0
2	FE	C	184	1/1	0.96	0.17	50,50,50,50	0
2	FE	I	181	1/1	0.96	0.19	49,49,49,49	0
2	FE	T	181	1/1	0.96	0.16	59,59,59,59	0
2	FE	C	180	1/1	0.96	0.15	54,54,54,54	0
2	FE	A	180	1/1	0.96	0.18	43,43,43,43	0
2	FE	K	181	1/1	0.96	0.16	53,53,53,53	0
2	FE	N	180	1/1	0.97	0.15	60,60,60,60	0
2	FE	S	180	1/1	0.97	0.11	53,53,53,53	0
2	FE	I	180	1/1	0.97	0.15	42,42,42,42	0
2	FE	O	180	1/1	0.97	0.12	47,47,47,47	0
2	FE	E	180	1/1	0.97	0.14	38,38,38,38	0
2	FE	J	180	1/1	0.97	0.13	54,54,54,54	0
2	FE	U	180	1/1	0.97	0.14	52,52,52,52	0
2	FE	L	181	1/1	0.97	0.15	59,59,59,59	0
2	FE	V	181	1/1	0.97	0.13	62,62,62,62	0
2	FE	W	180	1/1	0.97	0.14	40,40,40,40	0
2	FE	M	180	1/1	0.97	0.14	48,48,48,48	0
2	FE	C	181	1/1	0.97	0.19	56,56,56,56	0
2	FE	X	181	1/1	0.97	0.17	63,63,63,63	0
2	FE	I	184	1/1	0.98	0.14	39,39,39,39	0
2	FE	E	184	1/1	0.98	0.14	41,41,41,41	0
2	FE	L	180	1/1	0.98	0.13	35,35,35,35	0
2	FE	F	180	1/1	0.98	0.15	48,48,48,48	0
2	FE	Q	180	1/1	0.98	0.13	47,47,47,47	0
2	FE	V	180	1/1	0.98	0.10	49,49,49,49	0
2	FE	B	181	1/1	0.98	0.13	62,62,62,62	0
2	FE	H	180	1/1	0.98	0.12	40,40,40,40	0
2	FE	D	180	1/1	0.98	0.11	37,37,37,37	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FE	K	180	1/1	0.98	0.12	48,48,48,48	0
2	FE	N	184	1/1	0.98	0.09	42,42,42,42	0
2	FE	O	184	1/1	0.99	0.14	46,46,46,46	0
2	FE	D	184	1/1	0.99	0.14	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.