



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:48 PM UTC

PDB ID : 3RHK / pdb_00003rhk
Title : Crystal structure of the catalytic domain of c-Met kinase in complex with ARQ 197
Authors : Eathiraj, S.; Palma, R.; Volckova, E.; Hirschi, M.; France, D.S.; Ashwell, M.A.; Chan, T.C.
Deposited on : 2011-04-11
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	 4% 79% 13% • 6%
1	B	318	 2% 78% 14% • 6%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5303 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hepatocyte growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	0	0
			2317	1495	400	408	14			
1	B	299	Total	C	N	O	S	0	0	0
			2332	1506	395	417	14			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1029	MET	-	expression tag	UNP P08581
A	1030	GLY	-	expression tag	UNP P08581
A	1031	HIS	-	expression tag	UNP P08581
A	1032	HIS	-	expression tag	UNP P08581
A	1033	HIS	-	expression tag	UNP P08581
A	1034	HIS	-	expression tag	UNP P08581
A	1035	HIS	-	expression tag	UNP P08581
A	1036	HIS	-	expression tag	UNP P08581
A	1037	MET	-	expression tag	UNP P08581
B	1029	MET	-	expression tag	UNP P08581
B	1030	GLY	-	expression tag	UNP P08581
B	1031	HIS	-	expression tag	UNP P08581
B	1032	HIS	-	expression tag	UNP P08581
B	1033	HIS	-	expression tag	UNP P08581
B	1034	HIS	-	expression tag	UNP P08581
B	1035	HIS	-	expression tag	UNP P08581
B	1036	HIS	-	expression tag	UNP P08581
B	1037	MET	-	expression tag	UNP P08581

- Molecule 2 is 1-[(3R,4R)-4-(1H-indol-3-yl)-2,5-dioxopyrrolidin-3-yl]pyrrolo[3,2,1-ij]quinolinium (CCD ID: M97) (formula: C₂₃H₁₆N₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			28	23	3	2		
2	B	1	Total	C	N	O	0	0
			28	23	3	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	287	Total	O	0	0
			287	287		
3	B	311	Total	O	0	0
			311	311		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hepatocyte growth factor receptor



- Molecule 1: Hepatocyte growth factor receptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	53.47Å 58.67Å 64.96Å 88.41° 68.10° 85.52°	Depositor
Resolution (Å)	10.00 – 1.94 10.00 – 1.94	Depositor EDS
% Data completeness (in resolution range)	93.6 (10.00-1.94) 93.1 (10.00-1.94)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.93Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.200 , 0.254 0.202 , 0.256	Depositor DCC
R_{free} test set	2580 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	15.0	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.024 for -h,-k,-h+l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5303	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M97

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	13/2376 (0.5%)	0.90	7/3225 (0.2%)
1	B	0.99	17/2391 (0.7%)	0.87	0/3252
All	All	0.94	30/4767 (0.6%)	0.89	7/6477 (0.1%)

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1202	HIS	C-O	-8.70	1.13	1.24
1	B	1201	VAL	C-O	-8.33	1.15	1.24
1	A	1132	LYS	CA-CB	-7.87	1.40	1.53
1	A	1333	LEU	C-O	-6.90	1.16	1.24
1	B	1130	ILE	C-O	-6.86	1.16	1.24
1	A	1334	VAL	C-O	-6.70	1.16	1.24
1	A	1058	LEU	C-O	-6.32	1.16	1.23
1	A	1205	LEU	C-O	-6.26	1.16	1.24
1	B	1223	PHE	N-CA	-6.24	1.40	1.46
1	B	1224	GLY	C-O	-6.11	1.15	1.23
1	A	1204	ASP	C-O	-6.07	1.17	1.24
1	A	1135	SER	C-O	-5.98	1.16	1.24
1	B	1227	ARG	C-O	-5.91	1.16	1.24
1	A	1125	LEU	C-O	-5.89	1.17	1.24
1	A	1134	PHE	C-O	-5.78	1.17	1.23
1	A	1202	HIS	C-O	-5.66	1.17	1.24
1	B	1209	ASN	C-O	-5.65	1.16	1.24
1	B	1205	LEU	C-O	-5.62	1.17	1.24
1	B	1070	VAL	C-O	-5.57	1.17	1.24
1	B	1210	CYS	C-O	-5.54	1.17	1.24
1	B	1061	GLU	C-O	-5.51	1.17	1.24
1	A	1335	SER	C-O	-5.47	1.17	1.24
1	B	1073	PRO	C-O	-5.24	1.17	1.24
1	B	1207	ALA	CA-CB	-5.24	1.44	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1226	ALA	CA-CB	-5.23	1.44	1.53
1	B	1306	GLU	C-O	-5.16	1.17	1.23
1	A	1336	ARG	C-O	-5.15	1.18	1.24
1	A	1332	GLU	C-O	-5.10	1.18	1.24
1	B	1057	ALA	C-O	-5.08	1.17	1.24
1	B	1129	ILE	C-O	-5.04	1.18	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1130	ILE	N-CA-C	-9.72	94.13	109.20
1	A	1247	VAL	N-CA-C	5.71	116.35	110.36
1	A	1287	VAL	N-CA-C	5.67	116.04	108.11
1	A	1133	ASP	CB-CA-C	-5.61	103.38	111.91
1	A	1056	SER	CA-C-N	-5.34	114.11	122.65
1	A	1056	SER	C-N-CA	-5.34	114.11	122.65
1	A	1118	ILE	CB-CA-C	-5.27	105.03	112.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2317	0	2297	24	0
1	B	2332	0	2305	31	0
2	A	28	0	16	0	0
2	B	28	0	16	3	0
3	A	287	0	0	3	0
3	B	311	0	0	3	0
All	All	5303	0	4634	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1059:ASN:HD21	1:B:1061:GLU:HG2	1.19	1.03
1:A:1241:THR:HG21	3:A:401:HOH:O	1.59	1.01
1:B:1110:LYS:HG2	2:B:1:M97:H23	1.50	0.92
1:B:1059:ASN:HD21	1:B:1061:GLU:CG	1.92	0.81
1:B:1059:ASN:HD22	1:B:1062:LEU:H	1.26	0.81
1:A:1142:LEU:HD11	1:A:1155:VAL:HG23	1.63	0.81
1:B:1059:ASN:ND2	1:B:1061:GLU:HG2	1.96	0.81
1:A:1053:ILE:HD13	1:A:1121:VAL:HG21	1.73	0.71
1:A:1318:LYS:CD	1:A:1336:ARG:HH12	2.04	0.71
1:B:1110:LYS:HG2	2:B:1:M97:C23	2.23	0.69
1:A:1318:LYS:HD3	1:A:1336:ARG:HH12	1.58	0.68
1:B:1127:GLU:HA	1:B:1131:MET:HE2	1.78	0.65
1:B:1158:PRO:HB2	1:B:1160:MET:HE2	1.80	0.64
1:A:1241:THR:HG23	1:A:1243:ALA:H	1.63	0.64
1:B:1284:TYR:HB3	1:B:1287:VAL:HG22	1.81	0.61
1:B:1222:ASP:O	1:B:1222:ASP:CG	2.44	0.60
1:B:1059:ASN:ND2	1:B:1062:LEU:H	1.96	0.60
1:A:1129:ILE:O	3:A:383:HOH:O	2.17	0.59
1:B:1074:SER:HB3	3:B:435:HOH:O	2.03	0.58
1:B:1253:GLU:HA	1:B:1256:GLN:HE21	1.69	0.57
1:B:1247:VAL:HG21	1:B:1289:THR:HG22	1.88	0.55
1:A:1248:LYS:HG2	1:A:1292:ILE:HD13	1.90	0.52
1:A:1060:PRO:HA	1:A:1063:VAL:HG22	1.92	0.52
1:A:1171:ASN:C	1:A:1171:ASN:HD22	2.18	0.51
1:A:1230:TYR:CG	1:A:1244:LYS:HG2	2.46	0.51
1:A:1318:LYS:HZ2	1:A:1336:ARG:HH22	1.59	0.50
1:A:1318:LYS:NZ	1:A:1336:ARG:HH22	2.11	0.48
1:B:1212:LEU:HD13	1:B:1218:VAL:HG22	1.97	0.47
1:B:1171:ASN:HD22	1:B:1171:ASN:C	2.23	0.47
1:A:1251:ALA:HA	1:A:1267:TRP:CD2	2.50	0.47
1:B:1171:ASN:HD22	1:B:1173:THR:H	1.64	0.46
1:A:1230:TYR:CD1	1:A:1244:LYS:HG2	2.51	0.46
1:B:1212:LEU:CD1	1:B:1218:VAL:HG22	2.45	0.46
1:A:1130:ILE:O	1:A:1131:MET:CB	2.60	0.45
1:A:1131:MET:O	1:A:1132:LYS:CB	2.61	0.45
1:B:1219:LYS:HE2	3:B:74:HOH:O	2.16	0.45
1:A:1142:LEU:HD11	1:A:1155:VAL:CG2	2.39	0.44
1:B:1110:LYS:HB2	1:B:1110:LYS:HE2	1.48	0.44
1:B:1284:TYR:CB	1:B:1287:VAL:HG22	2.48	0.44
1:B:1134:PHE:HB3	1:B:1194:TYR:HE2	1.82	0.44
1:B:1158:PRO:HB2	1:B:1160:MET:CE	2.47	0.44
1:B:1046:LEU:HG	1:B:1048:GLN:HG2	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1110:LYS:HD2	1:B:1226:ALA:HB1	2.01	0.43
1:B:1208:ARG:HG2	3:B:565:HOH:O	2.18	0.42
1:B:1251:ALA:HA	1:B:1267:TRP:CD2	2.54	0.42
1:A:1211:MET:HE3	1:A:1223:PHE:CE2	2.55	0.42
1:A:1244:LYS:HE3	3:A:410:HOH:O	2.19	0.42
1:B:1282:PRO:HA	1:B:1283:PRO:HD3	1.96	0.42
1:B:1171:ASN:ND2	1:B:1173:THR:H	2.18	0.41
1:B:1085:GLY:HA3	2:B:1:M97:C13	2.50	0.41
1:A:1223:PHE:HB3	1:A:1226:ALA:HB3	2.02	0.41
1:A:1142:LEU:CD1	1:A:1155:VAL:HG23	2.41	0.41
1:A:1267:TRP:CD1	1:A:1267:TRP:C	2.98	0.41
1:A:1171:ASN:ND2	1:A:1173:THR:H	2.19	0.40
1:B:1171:ASN:HB3	1:B:1174:HIS:ND1	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/318 (93%)	287 (97%)	7 (2%)	2 (1%)	18	9
1	B	297/318 (93%)	289 (97%)	8 (3%)	0	100	100
All	All	593/636 (93%)	576 (97%)	15 (2%)	2 (0%)	36	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1132	LYS
1	A	1224	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/282 (88%)	244 (98%)	5 (2%)	48	39
1	B	254/282 (90%)	249 (98%)	5 (2%)	48	39
All	All	503/564 (89%)	493 (98%)	10 (2%)	48	39

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1049	ASN
1	A	1069	VAL
1	A	1118	ILE
1	A	1127	GLU
1	A	1140	LEU
1	B	1046	LEU
1	B	1059	ASN
1	B	1110	LYS
1	B	1150	GLU
1	B	1154	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1052	HIS
1	A	1100	ASN
1	A	1171	ASN
1	A	1239	ASN
1	B	1059	ASN
1	B	1068	HIS
1	B	1088	HIS
1	B	1167	ASN
1	B	1171	ASN
1	B	1256	GLN
1	B	1304	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	M97	B	1	-	32,33,33	2.91	9 (28%)	42,50,50	2.79	18 (42%)
2	M97	A	1	-	32,33,33	2.92	10 (31%)	42,50,50	2.71	17 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	M97	B	1	-	-	0/8/32/32	0/6/6/6
2	M97	A	1	-	-	0/8/32/32	0/6/6/6

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	M97	C17-C16	8.45	1.54	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	M97	C17-C16	8.35	1.54	1.36
2	B	1	M97	C18-N3	7.52	1.47	1.35
2	B	1	M97	C18-C17	7.43	1.54	1.38
2	A	1	M97	C18-N3	7.19	1.47	1.35
2	A	1	M97	C18-C17	7.11	1.53	1.38
2	B	1	M97	C16-C15	4.68	1.52	1.42
2	A	1	M97	C16-C15	4.65	1.52	1.42
2	A	1	M97	C19-N3	-4.02	1.33	1.39
2	A	1	M97	C10-C4	-3.56	1.37	1.44
2	B	1	M97	C19-N3	-3.34	1.34	1.39
2	B	1	M97	C8-N2	-3.19	1.33	1.37
2	A	1	M97	C7-N2	-2.92	1.33	1.37
2	B	1	M97	C20-C2	-2.83	1.38	1.44
2	B	1	M97	C10-C4	-2.67	1.39	1.44
2	A	1	M97	C20-C2	-2.65	1.39	1.44
2	A	1	M97	C8-N2	-2.59	1.34	1.37
2	B	1	M97	C7-N2	-2.52	1.34	1.37
2	A	1	M97	C15-C19	-2.26	1.38	1.42

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	M97	C10-C4-C5	7.08	111.66	106.25
2	A	1	M97	C4-C5-N1	-6.75	104.84	110.31
2	B	1	M97	C4-C5-N1	-6.71	104.87	110.31
2	B	1	M97	C10-C4-C5	6.56	111.26	106.25
2	A	1	M97	C18-N3-C19	5.49	125.10	118.26
2	B	1	M97	C17-C18-N3	-4.88	112.54	119.13
2	B	1	M97	C20-C2-C3	4.85	111.17	106.41
2	B	1	M97	C17-C16-C15	-4.74	113.89	120.48
2	B	1	M97	C18-N3-C19	4.58	123.97	118.26
2	B	1	M97	C6-C1-C7	-4.53	101.19	104.30
2	B	1	M97	C8-N2-C7	-4.49	110.72	113.87
2	B	1	M97	C6-C8-N2	4.39	110.79	107.69
2	A	1	M97	C1-C6-C8	-4.17	101.44	104.30
2	A	1	M97	C6-C8-N2	4.15	110.62	107.69
2	A	1	M97	C11-C10-C9	4.05	123.04	118.84
2	B	1	M97	C1-C7-N2	3.97	110.49	107.69
2	A	1	M97	C20-C2-C3	3.96	110.30	106.41
2	A	1	M97	C17-C16-C15	-3.88	115.09	120.48
2	A	1	M97	C17-C18-N3	-3.77	114.04	119.13
2	A	1	M97	C14-C9-C10	-3.65	118.65	122.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	M97	C1-C6-C8	-3.54	101.88	104.30
2	B	1	M97	C11-C10-C9	2.82	121.76	118.84
2	A	1	M97	C8-N2-C7	-2.80	111.91	113.87
2	B	1	M97	C14-C9-C10	-2.66	119.61	122.19
2	A	1	M97	C1-C7-N2	2.56	109.50	107.69
2	A	1	M97	C9-C10-C4	-2.53	104.62	106.98
2	A	1	M97	C6-C1-C7	-2.50	102.59	104.30
2	B	1	M97	O2-C8-C6	-2.49	123.27	126.77
2	A	1	M97	O2-C8-C6	-2.44	123.35	126.77
2	B	1	M97	C19-C20-C2	-2.41	102.44	105.90
2	A	1	M97	C10-C4-C1	-2.32	123.59	127.46
2	A	1	M97	C19-C20-C2	-2.19	102.75	105.90
2	B	1	M97	C9-C10-C4	-2.18	104.95	106.98
2	B	1	M97	C9-N1-C5	2.14	110.98	109.08
2	B	1	M97	C21-C20-C2	2.10	137.25	133.67

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	M97	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/318 (93%)	0.00	12 (4%) 42 47	4, 14, 30, 41	0
1	B	299/318 (94%)	-0.20	5 (1%) 69 75	4, 14, 22, 29	0
All	All	597/636 (93%)	-0.10	17 (2%) 55 62	4, 14, 27, 41	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1130	ILE	9.6
1	A	1346	GLY	6.5
1	A	1133	ASP	3.9
1	B	1048	GLN	3.3
1	A	1286	ASP	3.2
1	B	1286	ASP	2.9
1	A	1129	ILE	2.9
1	A	1127	GLU	2.4
1	A	1131	MET	2.4
1	A	1128	GLY	2.4
1	B	1291	ASP	2.2
1	A	1099	ASP	2.1
1	A	1101	ASP	2.1
1	A	1222	ASP	2.1
1	B	1231	ASP	2.1
1	A	1345	ILE	2.0
1	B	1287	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	M97	B	1	28/28	0.96	0.05	7,10,11,12	0
2	M97	A	1	28/28	0.97	0.04	7,9,10,11	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around M97 B 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.