



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:36 PM UTC

PDB ID : 3RHX / pdb_00003rhx
Title : Crystal structure of the catalytic domain of FGFR1 kinase in complex with ARQ 069
Authors : Eathiraj, S.; Palma, R.; Hirschi, M.; Volckova, E.; Nakuci, E.; Castro, J.; Chen, C.R.; Chan, T.C.; France, D.S.; Ashwell, M.A.
Deposited on : 2011-04-12
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	
1	B	306	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	1	-	-	X	-
4	EDO	A	767	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4994 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Basic fibroblast growth factor receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	275	Total	C	N	O	S	0	0	0
			2181	1391	373	400	17			
1	A	289	Total	C	N	O	S	0	0	0
			2258	1438	394	409	17			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	460	GLY	-	expression tag	UNP P11362
B	488	ALA	CYS	engineered mutation	UNP P11362
B	584	SER	CYS	engineered mutation	UNP P11362
A	460	GLY	-	expression tag	UNP P11362
A	488	ALA	CYS	engineered mutation	UNP P11362
A	584	SER	CYS	engineered mutation	UNP P11362

- Molecule 2 is (6S)-6-phenyl-5,6-dihydrobenzo[h]quinazolin-2-amine (CCD ID: 3RH) (formula: C₁₈H₁₅N₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	N	0	0
			21	18	3		
2	A	1	Total	C	N	0	0
			21	18	3		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	226	Total O 226 226	0	0
5	A	262	Total O 262 262	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	208.43Å 57.87Å 65.69Å 90.00° 107.22° 90.00°	Depositor
Resolution (Å)	8.00 – 2.01 8.00 – 2.01	Depositor EDS
% Data completeness (in resolution range)	97.3 (8.00-2.01) 95.7 (8.00-2.01)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.21 (at 2.02Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.201 , 0.262 0.213 , 0.266	Depositor DCC
R_{free} test set	2441 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 59.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4994	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 3RH, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.34	6/2306 (0.3%)	1.15	3/3123 (0.1%)
1	B	1.28	3/2224 (0.1%)	1.17	11/3003 (0.4%)
All	All	1.31	9/4530 (0.2%)	1.16	14/6126 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	536	LYS	N-CA	6.13	1.53	1.46
1	A	569	LEU	C-O	-6.11	1.17	1.24
1	A	671	ALA	N-CA	6.08	1.53	1.46
1	A	629	VAL	C-O	-5.55	1.18	1.24
1	B	623	ASP	C-O	-5.51	1.18	1.23
1	A	621	HIS	C-O	-5.36	1.17	1.24
1	B	701	TYR	CA-CB	5.27	1.56	1.52
1	B	570	ARG	C-O	-5.20	1.18	1.24
1	A	620	ILE	C-O	-5.17	1.18	1.24

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	744	ARG	CA-C-N	-8.91	111.66	120.21
1	B	744	ARG	C-N-CA	-8.91	111.66	120.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	675	ARG	CB-CA-C	7.32	121.58	111.63
1	B	610	GLY	N-CA-C	-6.57	104.55	112.49
1	B	506	ASN	N-CA-C	-6.40	103.92	112.94
1	A	593	GLU	N-CA-C	-5.72	99.68	109.24
1	A	752	GLU	N-CA-C	5.44	116.89	111.07
1	B	725	CYS	N-CA-C	-5.29	99.20	107.93
1	B	562	GLU	N-CA-C	5.26	117.66	110.24
1	B	486	GLU	N-CA-C	-5.24	102.14	109.69
1	A	725	CYS	N-CA-C	-5.24	99.73	108.32
1	B	659	ASN	N-CA-C	-5.09	100.78	108.67
1	B	701	TYR	CA-C-N	-5.02	114.63	119.90
1	B	701	TYR	C-N-CA	-5.02	114.63	119.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	503	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2243	38	0
1	B	2181	0	2191	40	0
2	A	21	0	15	1	0
2	B	21	0	15	1	0
3	B	5	0	0	1	0
4	A	16	0	24	12	0
4	B	4	0	6	0	0
5	A	262	0	0	8	0
5	B	226	0	0	9	0
All	All	4994	0	4494	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:609:ARG:HH11	1:B:609:ARG:CG	1.59	1.16
1:B:609:ARG:HG2	1:B:609:ARG:NH1	1.56	1.09
1:A:702:PRO:HG3	4:A:767:EDO:H11	1.39	1.03
1:A:533:GLU:OE1	4:A:1:EDO:H11	1.65	0.97
1:B:609:ARG:HH11	1:B:609:ARG:HG2	0.77	0.94
1:A:470:ARG:HH22	4:A:1:EDO:H22	1.33	0.92
1:A:749:GLN:HG2	5:A:796:HOH:O	1.76	0.84
1:A:702:PRO:HG3	4:A:767:EDO:C1	2.14	0.78
1:A:542:LYS:HE3	5:A:313:HOH:O	1.86	0.75
1:B:461:SER:N	5:B:285:HOH:O	2.21	0.74
1:A:731:MET:HB3	5:A:793:HOH:O	1.88	0.72
1:A:737:TRP:CD2	4:A:768:EDO:H21	2.28	0.68
1:A:542:LYS:HE2	5:A:287:HOH:O	1.93	0.68
1:A:702:PRO:HD3	4:A:767:EDO:H12	1.77	0.67
1:B:461:SER:N	5:B:2:HOH:O	2.29	0.66
1:A:533:GLU:OE1	4:A:1:EDO:C1	2.43	0.66
1:B:540:LYS:NZ	5:B:182:HOH:O	2.30	0.65
1:B:477:ARG:NH2	1:B:499:GLY:O	2.32	0.63
1:B:601:VAL:HG22	1:B:758:VAL:HG22	1.81	0.62
1:B:747:PHE:O	1:B:751:VAL:HG23	2.00	0.61
1:A:570:ARG:HE	1:A:661:ARG:NH1	1.98	0.61
1:A:737:TRP:CE3	4:A:768:EDO:H21	2.34	0.61
1:A:536:LYS:HB2	4:A:1:EDO:H12	1.83	0.61
1:A:592:GLU:N	5:A:325:HOH:O	2.33	0.61
1:B:570:ARG:NE	3:B:766:SO4:O2	2.34	0.61
1:A:606:GLN:NE2	1:A:637:MET:H	1.99	0.60
1:B:605:TYR:O	1:B:609:ARG:HG3	2.05	0.57
1:A:702:PRO:CG	4:A:767:EDO:H11	2.24	0.57
1:B:659:ASN:O	1:B:661:ARG:N	2.35	0.56
1:B:621:HIS:O	1:B:622:ARG:HB2	2.06	0.55
1:A:711:LYS:HE3	1:A:715:GLU:OE2	2.08	0.54
1:A:672:LEU:HD21	1:A:710:PHE:CE1	2.43	0.54
1:B:606:GLN:NE2	1:B:637:MET:H	2.07	0.53
1:B:609:ARG:CG	1:B:609:ARG:NH1	2.34	0.49
1:B:657:THR:OG1	1:B:659:ASN:O	2.29	0.49
1:B:708:GLU:OE2	5:B:253:HOH:O	2.20	0.49
1:A:570:ARG:NE	1:A:661:ARG:NH1	2.61	0.48
1:B:754:LEU:O	1:B:758:VAL:HG23	2.14	0.48
1:B:507:ARG:HG2	5:B:409:HOH:O	2.13	0.47
1:A:475:ARG:HG3	5:A:769:HOH:O	2.15	0.47
1:B:461:SER:CA	5:B:285:HOH:O	2.63	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:609:ARG:HD2	5:A:288:HOH:O	2.15	0.47
1:B:601:VAL:HG22	1:B:758:VAL:CG2	2.45	0.47
1:A:536:LYS:HD2	4:A:1:EDO:H12	1.95	0.46
1:A:672:LEU:HD21	1:A:710:PHE:HE1	1.79	0.46
1:A:491:GLN:HE21	1:A:491:GLN:HB3	1.52	0.46
1:B:622:ARG:HG2	1:B:677:TYR:CD1	2.51	0.46
1:B:606:GLN:HE22	1:B:636:VAL:HA	1.81	0.46
1:B:543:ASN:OD1	1:B:609:ARG:HD3	2.14	0.46
1:A:534:MET:HE2	1:A:538:ILE:HD11	1.98	0.46
1:A:572:TYR:O	1:A:576:ARG:HD2	2.16	0.45
1:B:491:GLN:HB2	5:B:323:HOH:O	2.17	0.44
1:B:669:PRO:HD3	1:B:684:TRP:CE3	2.53	0.44
1:A:611:MET:SD	1:A:639:ILE:HD13	2.58	0.43
1:B:675:ARG:O	1:B:675:ARG:HG2	2.16	0.43
1:B:748:LYS:HD3	5:B:193:HOH:O	2.18	0.43
1:A:731:MET:CB	5:A:793:HOH:O	2.58	0.43
1:A:731:MET:HA	1:A:731:MET:HE2	2.01	0.43
1:B:562:GLU:O	2:B:1:3RH:H7	2.18	0.43
1:A:494:LEU:HD13	1:A:563:TYR:CE1	2.54	0.43
1:A:667:MET:HE2	1:A:667:MET:HB3	1.84	0.43
1:A:568:ASN:OD1	1:A:568:ASN:C	2.62	0.42
1:B:670:GLU:OE1	1:B:670:GLU:N	2.44	0.42
1:B:665:LYS:NZ	5:B:406:HOH:O	2.52	0.42
1:A:718:ARG:HA	4:A:768:EDO:O2	2.20	0.42
1:A:684:TRP:CD1	1:A:684:TRP:C	2.97	0.41
1:A:490:GLY:HA2	1:A:517:LYS:CG	2.49	0.41
1:B:723:SER:HB2	1:A:721:LYS:NZ	2.35	0.41
1:B:570:ARG:HD3	1:B:661:ARG:NH2	2.35	0.41
1:B:718:ARG:NH2	1:B:737:TRP:O	2.53	0.41
1:B:540:LYS:HA	1:B:546:ASN:OD1	2.20	0.41
1:B:617:LYS:O	1:B:618:LYS:HB2	2.19	0.41
1:B:714:LYS:HE3	1:B:714:LYS:HB3	1.89	0.41
1:B:470:ARG:HE	1:B:470:ARG:HB2	1.52	0.40
1:A:640:ALA:HB1	2:A:2:3RH:H17	2.03	0.40
1:B:740:VAL:HA	1:B:741:PRO:HD3	1.84	0.40
1:B:667:MET:HE3	1:B:667:MET:HB3	1.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/306 (93%)	280 (98%)	5 (2%)	0	100	100
1	B	267/306 (87%)	260 (97%)	6 (2%)	1 (0%)	30	27
All	All	552/612 (90%)	540 (98%)	11 (2%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	660	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/268 (89%)	235 (98%)	4 (2%)	53	60
1	B	236/268 (88%)	226 (96%)	10 (4%)	26	25
All	All	475/536 (89%)	461 (97%)	14 (3%)	37	40

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	470	ARG
1	B	507	ARG
1	B	570	ARG
1	B	573	LEU
1	B	609	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	644	LEU
1	B	675	ARG
1	B	676	ILE
1	B	707	GLU
1	B	724	ASN
1	A	465	LEU
1	A	491	GLN
1	A	675	ARG
1	A	760	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	606	GLN
1	B	738	HIS
1	B	743	GLN
1	A	491	GLN
1	A	606	GLN
1	A	628	ASN
1	A	717	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	767	-	3,3,3	0.28	0	2,2,2	0.28	0
4	EDO	A	768	-	3,3,3	0.37	0	2,2,2	0.46	0
4	EDO	A	766	-	3,3,3	0.25	0	2,2,2	0.42	0
4	EDO	B	767	-	3,3,3	0.46	0	2,2,2	0.33	0
4	EDO	A	1	-	3,3,3	0.48	0	2,2,2	0.37	0
2	3RH	A	2	-	24,24,24	1.42	4 (16%)	29,34,34	2.48	12 (41%)
2	3RH	B	1	-	24,24,24	1.23	1 (4%)	29,34,34	2.16	11 (37%)
3	SO4	B	766	-	4,4,4	1.07	0	6,6,6	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	767	-	-	1/1/1/1	-
4	EDO	A	768	-	-	0/1/1/1	-
4	EDO	A	766	-	-	1/1/1/1	-
4	EDO	B	767	-	-	0/1/1/1	-
4	EDO	A	1	-	-	0/1/1/1	-
2	3RH	A	2	-	-	0/4/16/16	0/4/4/4
2	3RH	B	1	-	-	0/4/16/16	0/4/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2	3RH	C17-C10	3.30	1.44	1.39
2	B	1	3RH	C8-C9	3.02	1.56	1.53
2	A	2	3RH	C15-C14	2.74	1.44	1.38
2	A	2	3RH	C4-N5	2.17	1.38	1.33
2	A	2	3RH	C10-C9	-2.13	1.47	1.53

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	3RH	N5-C4-N6	5.34	122.92	117.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2	3RH	C15-C16-C17	-5.26	113.75	120.24
2	A	2	3RH	N6-C4-N3	-5.17	119.94	125.70
2	A	2	3RH	C1-N3-C4	5.04	121.99	117.03
2	A	2	3RH	C2-C1-N3	-4.41	117.90	122.70
2	B	1	3RH	N6-C4-N3	-4.01	121.24	125.70
2	B	1	3RH	C15-C16-C17	-3.75	115.62	120.24
2	A	2	3RH	C14-C13-C10	-3.68	116.24	120.65
2	A	2	3RH	N5-C4-N6	3.50	121.02	117.40
2	A	2	3RH	C12-C1-N3	3.23	122.24	117.46
2	B	1	3RH	C2-C7-N6	-3.21	118.61	123.83
2	B	1	3RH	C7-N6-C4	2.84	121.28	116.29
2	B	1	3RH	C1-N3-C4	2.70	119.68	117.03
2	A	2	3RH	C16-C17-C10	2.61	123.77	120.65
2	B	1	3RH	C14-C13-C10	-2.60	117.54	120.65
2	B	1	3RH	C21-C12-C11	2.59	121.31	118.50
2	B	1	3RH	C2-C1-N3	-2.56	119.92	122.70
2	A	2	3RH	C17-C10-C13	2.52	121.43	118.30
2	B	1	3RH	C10-C9-C11	2.38	116.52	113.34
2	A	2	3RH	C16-C15-C14	2.33	123.06	119.87
2	A	2	3RH	C7-N6-C4	2.26	120.27	116.29
2	B	1	3RH	C12-C1-C2	2.18	121.12	118.64
2	A	2	3RH	C2-C7-N6	-2.06	120.47	123.83

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	767	EDO	O1-C1-C2-O2
4	A	766	EDO	O1-C1-C2-O2


There are no ring outliers.

6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	767	EDO	4	0
4	A	768	EDO	3	0
4	A	1	EDO	5	0
2	A	2	3RH	1	0
2	B	1	3RH	1	0
3	B	766	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/306 (94%)	-0.44	3 (1%) 79 79	18, 33, 66, 76	0
1	B	275/306 (89%)	-0.27	7 (2%) 58 58	24, 36, 55, 67	0
All	All	564/612 (92%)	-0.36	10 (1%) 67 67	18, 34, 60, 76	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	659	ASN	6.1
1	A	519	ASP	5.5
1	B	653	TYR	5.3
1	B	654	TYR	4.5
1	B	657	THR	4.0
1	B	658	THR	3.5
1	B	675	ARG	3.1
1	A	464	GLU	2.2
1	B	731	MET	2.2
1	A	501	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	1	4/4	0.82	0.23	20,20,20,20	0
4	EDO	B	767	4/4	0.83	0.24	20,20,20,20	0
2	3RH	A	2	21/21	0.84	0.07	39,42,47,48	0
4	EDO	A	766	4/4	0.88	0.23	20,20,20,20	0
2	3RH	B	1	21/21	0.90	0.06	34,38,41,42	0
4	EDO	A	768	4/4	0.92	0.18	20,20,20,20	0
4	EDO	A	767	4/4	0.93	0.26	20,20,20,20	0
3	SO4	B	766	5/5	0.96	0.22	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.