



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:40 PM UTC

PDB ID : 4RH8 / pdb_00004rh8
Title : Crystal structure of the outer membrane lipopolysaccharide transport protein LptE (RlpB) from Escherichia coli in the tetragonal crystal form
Authors : Malojcic, G.; Kahne, D.
Deposited on : 2014-10-01
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	168	6% 66% 7% 5% . 21%
1	B	168	8% 64% 11% . . 21%
1	C	168	19% 67% 10% . . 20%
1	D	168	15% 66% 11% . 21%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4376 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LPS-assembly lipoprotein LptE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	133	Total	C	N	O	S	0	0	0
			1045	653	188	198	6			
1	B	133	Total	C	N	O	S	0	0	0
			1045	653	188	198	6			
1	C	135	Total	C	N	O	S	0	0	0
			1060	662	191	201	6			
1	D	133	Total	C	N	O	S	0	0	0
			1045	653	188	198	6			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	MET	CYS	conflict	UNP P0ADC1
A	183	LEU	THR	engineered mutation	UNP P0ADC1
A	184	VAL	PRO	engineered mutation	UNP P0ADC1
A	185	PRO	ALA	engineered mutation	UNP P0ADC1
B	19	MET	CYS	conflict	UNP P0ADC1
B	183	LEU	THR	engineered mutation	UNP P0ADC1
B	184	VAL	PRO	engineered mutation	UNP P0ADC1
B	185	PRO	ALA	engineered mutation	UNP P0ADC1
C	19	MET	CYS	conflict	UNP P0ADC1
C	183	LEU	THR	engineered mutation	UNP P0ADC1
C	184	VAL	PRO	engineered mutation	UNP P0ADC1
C	185	PRO	ALA	engineered mutation	UNP P0ADC1
D	19	MET	CYS	conflict	UNP P0ADC1
D	183	LEU	THR	engineered mutation	UNP P0ADC1
D	184	VAL	PRO	engineered mutation	UNP P0ADC1
D	185	PRO	ALA	engineered mutation	UNP P0ADC1

- Molecule 2 is FORMIC ACID (CCD ID: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			3	1	2		
2	B	1	Total	C	O	0	0
			3	1	2		
2	C	1	Total	C	O	0	0
			3	1	2		
2	D	1	Total	C	O	0	0
			3	1	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	57	Total	O	0	0
			57	57		
3	B	34	Total	O	0	0
			34	34		
3	C	32	Total	O	0	0
			32	32		
3	D	46	Total	O	0	0
			46	46		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LPS-assembly lipoprotein LptE



- Molecule 1: LPS-assembly lipoprotein LptE




- Molecule 1: LPS-assembly lipoprotein LptE



- Molecule 1: LPS-assembly lipoprotein LptE





4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	102.46Å 102.46Å 166.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	87.31 – 2.20 87.31 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (87.31-2.20) 100.0 (87.31-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.225 , 0.262 0.229 , 0.266	Depositor DCC
R_{free} test set	2260 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	46.6	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4376	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.36	5/1057 (0.5%)	1.26	12/1423 (0.8%)
1	B	1.17	0/1057	1.12	6/1423 (0.4%)
1	C	1.19	2/1073 (0.2%)	1.13	4/1446 (0.3%)
1	D	1.15	1/1057 (0.1%)	1.07	3/1423 (0.2%)
All	All	1.22	8/4244 (0.2%)	1.15	25/5715 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
1	C	0	4
1	D	0	3
All	All	0	12

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	126	PHE	CA-C	6.29	1.60	1.52
1	C	121	LYS	C-O	-6.20	1.17	1.23
1	A	33	MET	CA-C	-5.83	1.45	1.52
1	C	92	ASN	CA-CB	-5.70	1.43	1.53
1	A	127	PHE	CA-C	5.60	1.60	1.52
1	A	127	PHE	N-CA	5.36	1.53	1.46
1	A	107	THR	CB-CG2	-5.19	1.35	1.52
1	D	128	ASP	N-CA	5.07	1.55	1.46

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	91	ARG	N-CA-C	9.55	125.39	112.68
1	A	91	ARG	N-CA-C	9.28	125.02	112.68
1	A	107	THR	N-CA-CB	-8.01	96.80	111.13
1	A	36	MET	CG-SD-CE	-7.28	84.89	100.90
1	A	127	PHE	N-CA-C	7.15	126.02	110.80
1	B	69	ARG	CG-CD-NE	7.13	127.69	112.00
1	C	144	VAL	N-CA-CB	7.08	118.84	110.55
1	C	114	ASP	N-CA-C	6.99	119.54	110.53
1	A	150	ARG	CG-CD-NE	-6.87	96.90	112.00
1	A	126	PHE	CA-C-N	6.69	134.31	121.54
1	A	126	PHE	C-N-CA	6.69	134.31	121.54
1	A	126	PHE	CA-C-O	-6.68	114.31	121.45
1	A	36	MET	N-CA-CB	-6.53	101.44	111.56
1	B	150	ARG	CG-CD-NE	-6.47	97.76	112.00
1	C	68	THR	N-CA-C	-6.24	103.12	112.54
1	A	126	PHE	O-C-N	-5.90	116.32	123.10
1	A	91	ARG	CA-C-N	5.71	131.99	121.70
1	A	91	ARG	C-N-CA	5.71	131.99	121.70
1	B	91	ARG	CA-C-N	5.62	131.81	121.70
1	B	91	ARG	C-N-CA	5.62	131.81	121.70
1	D	79	LYS	CG-CD-CE	5.31	123.51	111.30
1	D	67	THR	CA-C-N	5.23	127.81	120.28
1	D	67	THR	C-N-CA	5.23	127.81	120.28
1	B	36	MET	CA-CB-CG	5.18	124.46	114.10
1	C	121	LYS	CB-CG-CD	5.12	123.09	111.30

There are no chirality outliers.

All (12) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	126	PHE	Peptide
1	A	91	ARG	Peptide
1	A	92	ASN	Peptide
1	B	126	PHE	Peptide
1	B	92	ASN	Peptide
1	C	66	GLU	Peptide
1	C	68	THR	Peptide
1	C	91	ARG	Peptide
1	C	92	ASN	Peptide
1	D	127	PHE	Peptide
1	D	91	ARG	Peptide
1	D	92	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1045	0	1079	15	0
1	B	1045	0	1079	19	0
1	C	1060	0	1092	17	0
1	D	1045	0	1079	11	0
2	A	3	0	2	0	0
2	B	3	0	2	0	0
2	C	3	0	2	1	0
2	D	3	0	1	0	0
3	A	57	0	0	2	0
3	B	34	0	0	2	0
3	C	32	0	0	3	0
3	D	46	0	0	3	0
All	All	4376	0	4336	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123:PHE:CE1	1:C:121:LYS:HD3	2.03	0.93
1:A:91:ARG:HH21	1:A:95:THR:HG21	1.52	0.74
1:B:91:ARG:HH21	1:B:95:THR:HG21	1.51	0.74
1:D:91:ARG:HH21	1:D:95:THR:HG21	1.53	0.73
1:C:91:ARG:HH21	1:C:95:THR:HG21	1.54	0.72
1:A:76:ARG:HB3	1:A:107:THR:HG22	1.73	0.71
1:B:92:ASN:HD22	1:B:92:ASN:N	1.89	0.69
1:A:92:ASN:HD22	1:A:92:ASN:N	1.88	0.69
1:D:144:VAL:HA	1:D:147:MET:HE3	1.74	0.67
1:A:123:PHE:HB2	3:D:341:HOH:O	1.95	0.66
1:D:121:LYS:HD2	3:D:341:HOH:O	1.96	0.65
1:C:95:THR:CG2	3:C:306:HOH:O	2.46	0.63
1:D:143:ILE:HG22	1:D:147:MET:HE2	1.83	0.61
1:C:98:TYR:CZ	1:C:129:ASN:HB3	2.36	0.61
1:A:28:GLN:HG3	3:A:356:HOH:O	2.01	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123:PHE:CE1	1:C:121:LYS:CD	2.80	0.60
1:B:95:THR:CG2	3:B:328:HOH:O	2.51	0.58
1:D:142:MET:HE1	1:D:145:LYS:HE3	1.87	0.56
1:B:150:ARG:NH1	1:B:153:GLU:OE2	2.40	0.55
1:A:150:ARG:NH1	1:A:153:GLU:OE2	2.40	0.55
1:B:92:ASN:N	1:B:92:ASN:ND2	2.56	0.54
1:A:92:ASN:N	1:A:92:ASN:ND2	2.55	0.54
1:B:150:ARG:HH11	1:B:153:GLU:CD	2.16	0.53
1:C:95:THR:HG23	3:C:306:HOH:O	2.09	0.53
1:A:150:ARG:HH11	1:A:153:GLU:CD	2.15	0.53
1:B:91:ARG:C	1:B:92:ASN:HD22	2.18	0.52
1:B:123:PHE:CZ	1:C:121:LYS:HD3	2.44	0.51
1:D:100:MET:HE2	1:D:147:MET:HE1	1.92	0.51
1:A:128:ASP:CG	1:A:128:ASP:O	2.54	0.50
1:A:91:ARG:C	1:A:92:ASN:HD22	2.20	0.49
1:C:43:PRO:O	1:C:52:ARG:NH2	2.36	0.49
1:B:43:PRO:O	1:B:52:ARG:NH2	2.35	0.49
1:A:124:ARG:HH21	1:A:146:GLU:CD	2.21	0.48
1:B:128:ASP:CG	1:B:128:ASP:O	2.56	0.48
1:B:36:MET:HG2	1:B:60:VAL:HG11	1.95	0.47
1:B:124:ARG:HH21	1:B:146:GLU:CD	2.21	0.47
1:B:87:ALA:HB1	1:D:79:LYS:NZ	2.30	0.46
1:C:98:TYR:CE1	1:C:129:ASN:HB3	2.51	0.46
1:D:142:MET:CE	1:D:145:LYS:HE3	2.45	0.46
1:B:36:MET:HE2	1:B:75:LEU:HB2	1.98	0.45
1:A:36:MET:HE2	1:A:75:LEU:HB2	1.98	0.45
1:C:36:MET:HG2	1:C:60:VAL:HG11	1.98	0.44
1:D:70:LYS:HD3	1:D:70:LYS:HA	1.86	0.44
1:C:139:GLU:CG	1:C:140:GLN:N	2.80	0.44
1:A:150:ARG:NH2	3:A:322:HOH:O	2.51	0.44
1:C:98:TYR:CE2	1:C:129:ASN:HB3	2.53	0.43
1:C:36:MET:HE2	1:C:75:LEU:HB2	2.00	0.43
1:A:65:LYS:O	1:A:65:LYS:HG3	2.17	0.43
1:C:95:THR:HG22	3:C:306:HOH:O	2.14	0.43
1:C:129:ASN:HA	1:C:130:PRO:HD3	1.80	0.42
1:A:98:TYR:CD1	1:A:98:TYR:N	2.88	0.42
1:B:95:THR:HG22	3:B:328:HOH:O	2.16	0.41
1:C:77:LEU:O	2:C:201:FMT:O1	2.38	0.41
1:C:139:GLU:HG2	1:C:140:GLN:N	2.35	0.41
1:D:103:THR:OG1	3:D:306:HOH:O	2.22	0.41
1:D:144:VAL:HA	1:D:147:MET:CE	2.48	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:84:LYS:HE2	1:B:140:GLN:NE2	2.36	0.40
1:B:91:ARG:NH2	1:B:95:THR:HG21	2.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/168 (77%)	127 (98%)	1 (1%)	1 (1%)	16	16
1	B	129/168 (77%)	128 (99%)	1 (1%)	0	100	100
1	C	131/168 (78%)	128 (98%)	2 (2%)	1 (1%)	16	16
1	D	129/168 (77%)	125 (97%)	4 (3%)	0	100	100
All	All	518/672 (77%)	508 (98%)	8 (2%)	2 (0%)	30	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	127	PHE
1	C	92	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/148 (79%)	108 (92%)	9 (8%)	12	13
1	B	117/148 (79%)	107 (92%)	10 (8%)	10	11
1	C	119/148 (80%)	114 (96%)	5 (4%)	26	36
1	D	117/148 (79%)	109 (93%)	8 (7%)	14	17
All	All	470/592 (79%)	438 (93%)	32 (7%)	14	17

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	65	LYS
1	A	69	ARG
1	A	81	SER
1	A	92	ASN
1	A	95	THR
1	A	107	THR
1	A	114	ASP
1	A	128	ASP
1	B	28	GLN
1	B	57	LEU
1	B	69	ARG
1	B	70	LYS
1	B	81	SER
1	B	92	ASN
1	B	95	THR
1	B	114	ASP
1	B	127	PHE
1	B	128	ASP
1	C	28	GLN
1	C	70	LYS
1	C	81	SER
1	C	95	THR
1	C	129	ASN
1	D	28	GLN
1	D	52	ARG
1	D	68	THR
1	D	81	SER
1	D	95	THR
1	D	114	ASP
1	D	139	GLU
1	D	145	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	B	92	ASN
1	B	94	GLN
1	B	140	GLN
1	C	44	ASN
1	C	99	GLN
1	D	92	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FMT	C	201	-	2,2,2	1.56	0	1,1,1	0.31	0
2	FMT	D	201	-	2,2,2	0.60	0	1,1,1	0.56	0
2	FMT	B	201	-	2,2,2	1.23	0	1,1,1	0.75	0
2	FMT	A	201	-	2,2,2	2.17	1 (50%)	1,1,1	0.16	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	FMT	O2-C	2.36	1.40	1.28

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	201	FMT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	133/168 (79%)	0.33	10 (7%)	20 17	27, 42, 79, 135	0
1	B	133/168 (79%)	0.60	13 (9%)	13 10	30, 49, 94, 134	0
1	C	135/168 (80%)	1.17	32 (23%)	2 1	31, 58, 120, 148	0
1	D	133/168 (79%)	1.03	25 (18%)	3 2	35, 56, 116, 133	0
All	All	534/672 (79%)	0.79	80 (14%)	5 4	27, 51, 111, 148	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	127	PHE	10.2
1	C	28	GLN	5.7
1	C	68	THR	5.5
1	B	127	PHE	5.5
1	B	123	PHE	4.9
1	D	70	LYS	4.9
1	B	126	PHE	4.8
1	C	67	THR	4.6
1	D	128	ASP	4.2
1	B	142	MET	4.2
1	C	65	LYS	4.1
1	B	28	GLN	4.1
1	C	167	ILE	4.0
1	C	162	ILE	4.0
1	C	165	ALA	3.8
1	B	128	ASP	3.8
1	C	130	PRO	3.7
1	C	70	LYS	3.6
1	A	170	ASP	3.5
1	C	170	ASP	3.5
1	D	69	ARG	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	112	GLY	3.4
1	A	142	MET	3.3
1	A	95	THR	3.3
1	C	63	LEU	3.2
1	D	167	ILE	3.2
1	C	164	ALA	3.2
1	B	98	TYR	3.2
1	D	168	ARG	3.1
1	D	65	LYS	3.1
1	D	63	LEU	3.1
1	A	128	ASP	3.1
1	A	28	GLN	3.0
1	C	116	TYR	3.0
1	C	139	GLU	2.9
1	D	170	ASP	2.9
1	D	68	THR	2.9
1	D	111	PRO	2.8
1	D	116	TYR	2.8
1	D	33	MET	2.8
1	C	33	MET	2.8
1	C	71	ASP	2.8
1	C	72	VAL	2.8
1	A	139	GLU	2.7
1	D	110	ILE	2.7
1	C	110	ILE	2.7
1	C	161	SER	2.6
1	B	139	GLU	2.6
1	B	125	SER	2.6
1	C	168	ARG	2.6
1	C	109	LEU	2.6
1	B	170	ASP	2.6
1	C	169	SER	2.5
1	D	109	LEU	2.5
1	D	115	ILE	2.5
1	C	34	LYS	2.5
1	D	28	GLN	2.4
1	D	164	ALA	2.4
1	C	32	THR	2.4
1	B	99	GLN	2.4
1	D	35	VAL	2.4
1	D	57	LEU	2.4
1	C	69	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	70	LYS	2.4
1	C	111	PRO	2.3
1	D	112	GLY	2.3
1	C	31	SER	2.3
1	A	140	GLN	2.3
1	A	93	GLY	2.2
1	A	126	PHE	2.2
1	D	114	ASP	2.2
1	C	140	GLN	2.2
1	C	113	ARG	2.1
1	D	67	THR	2.1
1	D	37	ILE	2.1
1	C	163	ARG	2.1
1	D	163	ARG	2.1
1	D	169	SER	2.1
1	B	68	THR	2.0
1	C	37	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FMT	C	201	3/3	0.76	0.25	48,48,63,76	0
2	FMT	B	201	3/3	0.83	0.16	42,42,55,61	0
2	FMT	A	201	3/3	0.84	0.19	40,40,44,60	0
2	FMT	D	201	3/3	0.95	0.14	59,59,67,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.