



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:09 AM UTC

PDB ID : 4RHZ / pdb_00004rhz
Title : Crystal structure of Cry23Aa1 and Cry37Aa1 binary protein complex
Authors : Rydel, T.J.; Williams, J.M.; Brown, G.R.; Guzov, V.M.; Sturman, E.J.; Evdokimov, A.
Deposited on : 2014-10-03
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	5% 72% 19% 6% .
2	B	126	10% 74% 20% 5% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3260 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cry23AA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	7	0
			2031	1284	326	417	4			

- Molecule 2 is a protein called Cry37AA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	125	Total	C	N	O	S	0	6	0
			1018	650	155	210	3			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	125	Total	O	0	0
			125	125		
5	B	84	Total	O	0	0
			84	84		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cry23AA1



• Molecule 2: Cry37AA1



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	126.39Å 126.39Å 79.23Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.50 – 2.35 49.50 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.50-2.35) 99.9 (49.50-2.35)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.89 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.192 , 0.245 0.186 , 0.238	Depositor DCC
R_{free} test set	1532 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	45.7	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.033 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3260	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.36	6/2096 (0.3%)	1.27	11/2869 (0.4%)
2	B	1.39	4/1065 (0.4%)	1.32	9/1455 (0.6%)
All	All	1.37	10/3161 (0.3%)	1.29	20/4324 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	81	GLU	C-O	7.96	1.34	1.24
2	B	106	VAL	CA-CB	7.09	1.64	1.54
2	B	10	ILE	N-CA	-6.13	1.39	1.46
2	B	98	VAL	CA-CB	5.58	1.61	1.54
1	A	79	HIS	CA-C	5.56	1.59	1.52
1	A	50	VAL	CA-C	5.35	1.59	1.52
1	A	93	THR	CA-C	5.16	1.58	1.52
1	A	106	VAL	CA-CB	5.14	1.59	1.54
1	A	87	GLU	CA-C	-5.09	1.46	1.52
1	A	24	LYS	C-O	5.00	1.29	1.24

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	241	GLY	N-CA-C	-10.31	99.57	112.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	108	SER	N-CA-C	8.34	118.95	108.19
2	B	81	GLU	CA-C-N	8.04	132.24	120.90
2	B	81	GLU	C-N-CA	8.04	132.24	120.90
2	B	65	ASP	CB-CA-C	-7.82	97.81	110.79
1	A	79	HIS	N-CA-C	7.51	120.58	109.24
2	B	18	TRP	N-CA-C	7.07	121.29	107.57
1	A	55	THR	CB-CA-C	-7.04	95.11	111.09
2	B	80	LYS	N-CA-C	6.64	119.82	109.52
2	B	17	GLU	N-CA-C	6.62	124.91	110.80
1	A	39	THR	CA-C-N	6.21	125.97	119.76
1	A	39	THR	C-N-CA	6.21	125.97	119.76
2	B	81	GLU	CA-C-O	5.39	126.52	120.70
2	B	83	ASP	N-CA-CB	-5.24	101.72	110.42
1	A	142	ILE	N-CA-C	5.23	112.73	107.55
1	A	244	THR	CA-C-N	5.22	125.14	119.76
1	A	244	THR	C-N-CA	5.22	125.14	119.76
1	A	32	LYS	N-CA-C	5.12	117.76	109.06
1	A	148	VAL	CB-CA-C	5.03	118.24	110.50
2	B	65	ASP	N-CA-CB	-5.00	102.76	110.12

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	108	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2031	0	1981	46	0
2	B	1018	0	960	30	0
3	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	125	0	0	4	0
5	B	84	0	0	1	0
All	All	3260	0	2941	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:ASN:ND2	1:A:242:ARG:CA	1.79	1.46
1:A:239:ASN:ND2	1:A:242:ARG:HA	0.96	1.25
1:A:239:ASN:HD21	1:A:243:SER:N	1.35	1.22
1:A:239:ASN:HD21	1:A:242:ARG:C	1.58	1.10
1:A:239:ASN:HD22	1:A:242:ARG:CA	1.46	1.08
2:B:81:GLU:HG2	2:B:82:TYR:CD1	1.89	1.07
1:A:239:ASN:ND2	1:A:243:SER:H	1.55	1.03
1:A:239:ASN:ND2	1:A:243:SER:N	2.04	0.97
1:A:54:LEU:CD1	1:A:154[B]:ILE:HD12	1.99	0.93
2:B:59:THR:O	2:B:60[B]:ILE:HD13	1.71	0.91
1:A:54:LEU:HD11	1:A:154[B]:ILE:HD12	1.53	0.88
2:B:22:GLU:HB2	2:B:82:TYR:CE1	2.10	0.86
1:A:239:ASN:ND2	1:A:242:ARG:C	2.23	0.85
2:B:81:GLU:CD	2:B:82:TYR:CE1	2.61	0.77
2:B:81:GLU:HG2	2:B:82:TYR:CE1	2.21	0.75
1:A:132[B]:HIS:NE2	5:A:659:HOH:O	2.19	0.73
2:B:56:TYR:CE2	2:B:58:GLN:HG3	2.25	0.71
2:B:11:ASN:OD1	2:B:55[B]:THR:HG22	1.91	0.70
2:B:22:GLU:HB2	2:B:82:TYR:HE1	1.56	0.70
1:A:54:LEU:HD11	1:A:154[B]:ILE:CD1	2.22	0.70
1:A:75:TRP:CZ3	1:A:77:GLU:HB3	2.26	0.70
2:B:81:GLU:OE2	2:B:82:TYR:CZ	2.45	0.69
1:A:145:LYS:HD2	1:A:238:ASP:OD2	1.93	0.68
2:B:81:GLU:CG	2:B:82:TYR:CE1	2.75	0.68
1:A:76:THR:HG22	1:A:133:THR:OG1	1.93	0.67
1:A:125:THR:HG23	5:A:653:HOH:O	1.94	0.66
1:A:54:LEU:HD12	1:A:154[B]:ILE:HD12	1.76	0.66
1:A:235:PRO:HG3	1:A:243:SER:O	1.95	0.66
2:B:81:GLU:OE2	2:B:82:TYR:CE1	2.49	0.65
2:B:73:MET:HG2	2:B:75:PHE:CZ	2.32	0.65
1:A:190:SER:HB2	1:A:193:ASP:CG	2.23	0.63
2:B:56:TYR:HE2	2:B:58:GLN:HG3	1.62	0.63
2:B:83:ASP:O	2:B:84:VAL:C	2.41	0.62
1:A:239:ASN:ND2	1:A:242:ARG:N	2.48	0.60
1:A:191:VAL:HG13	5:A:720:HOH:O	2.02	0.59
1:A:78:THR:OG1	1:A:131:THR:HG22	2.03	0.59
2:B:73:MET:HE3	2:B:121:TYR:CE1	2.37	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:82:TYR:OH	5:B:680:HOH:O	2.14	0.57
2:B:59:THR:C	2:B:60[B]:ILE:HD13	2.29	0.57
1:A:112:LEU:C	1:A:112:LEU:HD12	2.31	0.56
2:B:81:GLU:CG	2:B:82:TYR:CD1	2.76	0.55
1:A:154[A]:ILE:HD13	1:A:155:GLN:N	2.23	0.54
2:B:83:ASP:OD2	2:B:83:ASP:N	2.39	0.54
1:A:260:LEU:N	1:A:260:LEU:HD22	2.22	0.53
2:B:73:MET:HE3	2:B:121:TYR:CD1	2.44	0.53
1:A:102:LYS:HD3	1:A:113:THR:OG1	2.09	0.51
1:A:145:LYS:HD2	1:A:238:ASP:CG	2.36	0.51
1:A:190:SER:HB2	1:A:193:ASP:OD2	2.11	0.50
1:A:97:THR:HG23	5:A:706:HOH:O	2.11	0.50
2:B:24:TYR:CD1	2:B:45:ASP:HB2	2.47	0.49
2:B:17:GLU:O	2:B:17:GLU:HG3	2.12	0.49
1:A:166:GLU:HG2	1:A:214:LYS:HA	1.95	0.48
1:A:234[A]:TYR:CD1	1:A:245:PRO:HB3	2.48	0.48
2:B:81:GLU:O	2:B:89:ASP:N	2.47	0.47
1:A:239:ASN:HD21	1:A:242:ARG:CA	1.71	0.47
1:A:54:LEU:CD1	1:A:154[B]:ILE:CD1	2.81	0.46
2:B:83:ASP:HB3	2:B:86:ASN:O	2.17	0.45
1:A:238:ASP:O	1:A:240:LYS:HD3	2.17	0.45
1:A:245:PRO:HB2	2:B:34:PRO:HD2	1.99	0.45
2:B:81:GLU:O	2:B:88:ASP:HB3	2.16	0.45
1:A:190:SER:HB3	1:A:193:ASP:H	1.82	0.44
1:A:75:TRP:HZ3	1:A:77:GLU:HB3	1.80	0.44
1:A:45:ILE:HG22	1:A:47:THR:H	1.83	0.44
2:B:60[A]:ILE:HD13	2:B:60[A]:ILE:HG21	1.75	0.43
1:A:236:LEU:O	1:A:239:ASN:HB2	2.18	0.43
1:A:81:GLU:O	1:A:127:THR:HA	2.20	0.42
1:A:75:TRP:CE3	1:A:154[B]:ILE:HD13	2.55	0.41
1:A:32:LYS:O	1:A:169:MET:HA	2.20	0.41
1:A:166:GLU:HA	1:A:213:PHE:O	2.20	0.41
2:B:29:ALA:O	2:B:37:ASP:HA	2.20	0.41
2:B:81:GLU:HG2	2:B:82:TYR:H	1.86	0.41
1:A:44:GLU:OE1	1:A:106:VAL:HG12	2.21	0.41
2:B:81:GLU:HB3	2:B:89:ASP:HB2	2.01	0.41
1:A:10:ILE:HD13	1:A:10:ILE:HA	1.83	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/267 (99%)	257 (97%)	6 (2%)	1 (0%)	30	34
2	B	129/126 (102%)	120 (93%)	9 (7%)	0	100	100
All	All	393/393 (100%)	377 (96%)	15 (4%)	1 (0%)	36	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	242	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/235 (100%)	205 (88%)	29 (12%)	4	4
2	B	115/110 (104%)	107 (93%)	8 (7%)	14	15
All	All	349/345 (101%)	312 (89%)	37 (11%)	6	6

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	THR
1	A	24	LYS
1	A	44	GLU
1	A	55	THR
1	A	79	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	94	SER
1	A	98	LYS
1	A	99	GLN
1	A	100	SER
1	A	109	ASP
1	A	110	VAL
1	A	111	THR
1	A	112	LEU
1	A	148	VAL
1	A	153	ILE
1	A	154[A]	ILE
1	A	154[B]	ILE
1	A	156	ASN
1	A	175	CYS
1	A	190	SER
1	A	191	VAL
1	A	194	LEU
1	A	206	LYS
1	A	226	ARG
1	A	237	ASP
1	A	240	LYS
1	A	242	ARG
1	A	243	SER
1	A	260	LEU
2	B	22	GLU
2	B	39	GLU
2	B	48	LYS
2	B	58	GLN
2	B	61	LYS
2	B	65	ASP
2	B	73	MET
2	B	83	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	155	GLN
1	A	156	ASN
1	A	239	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	259/267 (97%)	0.15	14 (5%)	31 37	19, 41, 60, 92	7 (2%)
2	B	125/126 (99%)	0.16	13 (10%)	11 13	25, 35, 75, 82	6 (4%)
All	All	384/393 (97%)	0.15	27 (7%)	22 25	19, 40, 67, 92	13 (3%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	84	VAL	8.1
2	B	82	TYR	6.1
2	B	18	TRP	5.9
1	A	238	ASP	5.6
1	A	240	LYS	5.1
2	B	83	ASP	4.9
1	A	239	ASN	4.0
2	B	17	GLU	3.9
1	A	241	GLY	3.8
1	A	260	LEU	3.7
2	B	86	ASN	3.4
1	A	243	SER	3.2
2	B	85	GLY	2.9
2	B	87	ALA	2.9
2	B	48	LYS	2.8
1	A	106	VAL	2.6
2	B	16	GLY	2.6
1	A	93	THR	2.6
1	A	110	VAL	2.4
2	B	81	GLU	2.3
1	A	237	ASP	2.3
2	B	47	GLY	2.3
1	A	65	TYR	2.2
1	A	78	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	242	ARG	2.2
1	A	107	ASN	2.1
2	B	88	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	A	301	1/1	0.90	0.13	82,82,82,82	0
4	CA	B	501	1/1	0.98	0.08	62,62,62,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.