



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 06:05 PM UTC

PDB ID : 4RJ2 / pdb_00004rj2
Title : Crystal structure of E.coli purine nucleoside phosphorylase at 0.99 Å resolution
Authors : Timofeev, V.I.; Abramchik, Y.A.; Esipov, R.S.; Kuranova, I.P.
Deposited on : 2014-10-08
Resolution : 0.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 0.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1358 (1.04-0.96)
Clashscore	190562	1411 (1.04-0.96)
Ramachandran outliers	187476	1347 (1.04-0.96)
Sidechain outliers	187428	1348 (1.04-0.96)
RSRZ outliers	180081	1355 (1.04-0.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	6% 89% 9%
1	B	237	26% 89% 10%
1	C	237	9% 91% 8%
1	D	237	3% 89% 11%
1	E	237	8% 90% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	237	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	C	301[A]	-	-	X	-
2	GOL	D	301[A]	-	-	X	-
2	GOL	D	301[B]	-	-	X	-
2	GOL	D	301[C]	-	-	X	-
2	GOL	D	303	-	X	X	-
2	GOL	F	301	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12253 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Purine nucleoside phosphorylase DeoD-type.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	237	Total	C	N	O	S	0	20	0
			1960	1237	340	366	17			
1	B	237	Total	C	N	O	S	0	8	0
			1859	1176	318	347	18			
1	C	237	Total	C	N	O	S	0	8	0
			1861	1173	322	349	17			
1	D	237	Total	C	N	O	S	0	9	0
			1866	1180	320	349	17			
1	E	237	Total	C	N	O	S	0	4	0
			1829	1157	312	344	16			
1	F	237	Total	C	N	O	S	0	7	0
			1853	1170	319	346	18			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 18 9 9	0	1
2	C	1	Total C O 12 6 6	0	1
2	D	1	Total C O 18 9 9	0	1
2	D	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	F	1	Total C O 6 3 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	176	Total O 178 178	0	2
3	B	124	Total O 125 125	0	1
3	C	180	Total O 183 183	0	3
3	D	194	Total O 195 195	0	1
3	E	171	Total O 174 174	0	3
3	F	104	Total O 104 104	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Purine nucleoside phosphorylase DeoD-type




- Molecule 1: Purine nucleoside phosphorylase DeoD-type



- Molecule 1: Purine nucleoside phosphorylase DeoD-type




- Molecule 1: Purine nucleoside phosphorylase DeoD-type



K237

- Molecule 1: Purine nucleoside phosphorylase DeoD-type




Chain E:  8% 90% 9%



K237

- Molecule 1: Purine nucleoside phosphorylase DeoD-type

Chain F:  28% 85% 12%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.12Å 110.22Å 88.21Å 90.00° 111.08° 90.00°	Depositor
Resolution (Å)	16.28 – 0.99 16.28 – 0.99	Depositor EDS
% Data completeness (in resolution range)	98.3 (16.28-0.99) 98.3 (16.28-0.99)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 0.99Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.176 , 0.189 0.185 , 0.196	Depositor DCC
R_{free} test set	35993 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	8.6	Xtriage
Anisotropy	0.459	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.47 , 54.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12253	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/1988	0.73	0/2674
1	B	0.50	0/1888	0.72	0/2541
1	C	0.54	0/1890	0.72	0/2546
1	D	0.56	0/1895	0.74	0/2554
1	E	0.58	1/1859 (0.1%)	0.73	0/2506
1	F	0.55	0/1882	0.75	0/2533
All	All	0.55	1/11402 (0.0%)	0.73	0/15354

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	201	THR	C-O	-5.69	1.17	1.24

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	216	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1960	0	1967	28	0
1	B	1859	0	1869	24	0
1	C	1861	0	1863	26	0
1	D	1866	0	1874	41	0
1	E	1829	0	1827	17	0
1	F	1853	0	1859	68	0
2	A	18	0	24	4	0
2	C	12	0	16	7	0
2	D	30	0	40	49	0
2	F	6	0	8	11	0
3	A	178	0	0	6	0
3	B	125	0	0	5	0
3	C	183	0	0	5	0
3	D	195	0	0	4	0
3	E	174	0	0	4	0
3	F	104	0	0	5	0
All	All	12253	0	11347	222	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 222 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:204:ASP:HB2	1:F:211:GLN:CG	1.39	1.48
1:F:212:THR:CG2	1:F:216:GLU:HB2	1.62	1.29
1:F:212:THR:HG23	1:F:216:GLU:CB	1.62	1.27
1:F:204:ASP:CB	1:F:211:GLN:CG	2.26	1.14
1:F:204:ASP:HA	1:F:211:GLN:HA	1.24	1.13

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/237 (107%)	248 (98%)	6 (2%)	0	100	100
1	B	243/237 (102%)	238 (98%)	5 (2%)	0	100	100
1	C	243/237 (102%)	237 (98%)	6 (2%)	0	100	100
1	D	244/237 (103%)	237 (97%)	7 (3%)	0	100	100
1	E	239/237 (101%)	229 (96%)	10 (4%)	0	100	100
1	F	242/237 (102%)	228 (94%)	9 (4%)	5 (2%)	5	0
All	All	1465/1422 (103%)	1417 (97%)	43 (3%)	5 (0%)	30	12

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	209	HIS
1	F	215	ALA
1	F	216	GLU
1	F	211	GLN
1	F	217	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/188 (110%)	201 (98%)	5 (2%)	43	9
1	B	195/188 (104%)	190 (97%)	5 (3%)	40	7
1	C	195/188 (104%)	188 (96%)	7 (4%)	31	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	196/188 (104%)	194 (99%)	2 (1%)	68	33
1	E	191/188 (102%)	185 (97%)	6 (3%)	35	5
1	F	194/188 (103%)	187 (96%)	7 (4%)	31	4
All	All	1177/1128 (104%)	1145 (97%)	32 (3%)	40	7

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	179	GLU
1	F	205	HIS
1	C	41	ASN
1	C	24[B]	ARG
1	F	208	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	6	ASN
1	F	218	GLN
1	D	97	HIS
1	E	222	ASN
1	D	41	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	301[B]	-	5,5,5	0.37	0	5,5,5	1.08	0
2	GOL	A	301[C]	-	5,5,5	0.39	0	5,5,5	0.92	0
2	GOL	D	302	-	5,5,5	0.30	0	5,5,5	0.52	0
2	GOL	A	301[A]	-	5,5,5	0.20	0	5,5,5	1.38	1 (20%)
2	GOL	D	301[B]	-	5,5,5	0.20	0	5,5,5	1.02	0
2	GOL	C	301[B]	-	5,5,5	0.53	0	5,5,5	0.90	0
2	GOL	D	301[C]	-	5,5,5	0.51	0	5,5,5	1.25	1 (20%)
2	GOL	D	301[A]	-	5,5,5	0.18	0	5,5,5	0.64	0
2	GOL	C	301[A]	-	5,5,5	0.28	0	5,5,5	0.94	0
2	GOL	D	303	-	5,5,5	0.85	0	5,5,5	3.45	3 (60%)
2	GOL	F	301	-	5,5,5	0.73	0	5,5,5	1.64	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	301[B]	-	-	4/4/4/4	-
2	GOL	A	301[C]	-	-	2/4/4/4	-
2	GOL	D	302	-	-	0/4/4/4	-
2	GOL	A	301[A]	-	-	3/4/4/4	-
2	GOL	D	301[B]	-	-	0/4/4/4	-
2	GOL	C	301[B]	-	-	0/4/4/4	-
2	GOL	D	301[C]	-	-	2/4/4/4	-
2	GOL	D	301[A]	-	-	4/4/4/4	-
2	GOL	C	301[A]	-	-	1/4/4/4	-
2	GOL	D	303	-	-	4/4/4/4	-
2	GOL	F	301	-	-	2/4/4/4	-

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	D	303	GOL	C3-C2-C1	6.43	135.38	111.80
2	D	303	GOL	O2-C2-C3	-3.15	96.14	109.18
2	D	303	GOL	O2-C2-C1	-2.81	97.55	109.18
2	F	301	GOL	O2-C2-C3	-2.32	99.59	109.18
2	D	301[C]	GOL	C3-C2-C1	2.16	119.72	111.80

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301[A]	GOL	C1-C2-C3-O3
2	A	301[B]	GOL	O2-C2-C3-O3
2	A	301[C]	GOL	C1-C2-C3-O3
2	D	301[A]	GOL	C1-C2-C3-O3
2	D	301[C]	GOL	C1-C2-C3-O3

There are no ring outliers.

9 monomers are involved in 71 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301[C]	GOL	2	0
2	A	301[A]	GOL	2	0
2	D	301[B]	GOL	12	0
2	C	301[B]	GOL	3	0
2	D	301[C]	GOL	18	0
2	D	301[A]	GOL	13	0
2	C	301[A]	GOL	4	0
2	D	303	GOL	34	0
2	F	301	GOL	11	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/237 (100%)	0.48	15 (6%) 26 28	3, 6, 21, 39	20 (8%)
1	B	237/237 (100%)	1.33	62 (26%) 1 1	3, 11, 24, 73	8 (3%)
1	C	237/237 (100%)	0.39	21 (8%) 15 15	2, 7, 18, 24	8 (3%)
1	D	237/237 (100%)	0.30	8 (3%) 48 54	2, 7, 14, 23	9 (3%)
1	E	237/237 (100%)	0.48	20 (8%) 17 17	3, 7, 21, 77	4 (1%)
1	F	237/237 (100%)	1.60	67 (28%) 1 1	3, 10, 36, 93	7 (2%)
All	All	1422/1422 (100%)	0.76	193 (13%) 7 6	2, 8, 22, 93	56 (3%)

The worst 5 of 193 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	214	ALA	12.1
1	E	208	THR	7.9
1	F	213	THR	7.4
1	F	206	ILE	7.2
1	A	215	ALA	7.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	F	301	6/6	0.66	0.19	10,15,27,30	0
2	GOL	A	301[B]	6/6	0.80	0.37	5,10,13,14	6
2	GOL	A	301[C]	6/6	0.80	0.37	8,11,12,16	6
2	GOL	A	301[A]	6/6	0.80	0.37	7,8,11,11	6
2	GOL	C	301[B]	6/6	0.84	0.21	13,14,15,17	6
2	GOL	C	301[A]	6/6	0.84	0.21	7,9,11,19	6
2	GOL	D	303	6/6	0.85	0.17	14,17,19,21	0
2	GOL	D	301[C]	6/6	0.88	0.24	9,10,13,18	6
2	GOL	D	301[A]	6/6	0.88	0.24	3,9,10,10	6
2	GOL	D	301[B]	6/6	0.88	0.24	7,10,12,12	6
2	GOL	D	302	6/6	0.94	0.09	12,12,13,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.