



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 11:07 PM UTC

PDB ID : 1RKY / pdb_00001rky
Title : PPLO + Xe
Authors : Guss, J.M.; Duff, A.P.
Deposited on : 2003-11-24
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1054 (1.68-1.68)
Clashscore	190562	1078 (1.68-1.68)
Ramachandran outliers	187476	1068 (1.68-1.68)
Sidechain outliers	187428	1067 (1.68-1.68)
RSRZ outliers	180081	1055 (1.68-1.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	747	3% 93% 5%
2	B	3	33% 33% 33%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 6853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called lysyl oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	735	Total	C	N	O	S	2	49	0
			6087	3861	974	1236	16			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	314	VAL	ILE	conflict	GB 13936870
A	338	ASP	GLU	conflict	GB 13936870
A	417	SER	ALA	conflict	GB 13936870
A	478	TPQ	TYR	modified residue	GB 13936870
A	549	GLN	LYS	conflict	GB 13936870
A	577	PHE	LEU	conflict	GB 13936870
A	579	ASN	LYS	conflict	GB 13936870
A	650	THR	GLU	conflict	GB 13936870
A	758	VAL	LEU	conflict	GB 13936870
A	761	ALA	PRO	conflict	GB 13936870

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	1
			28	16	2	10		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cu	0	0
			1	1		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Ca	0	0
			2	2		


- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	Mg	0	0
			3	3		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	6	Total Cl 6 6	0	0

- Molecule 8 is IMIDAZOLE (CCD ID: IMD) (formula: C₃H₅N₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C N 5 3 2	0	0

- Molecule 9 is XENON (CCD ID: XE) (formula: Xe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	8	Total Xe 8 8	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	632	Total O 632 632	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: lysyl oxidase



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	140.14Å 66.70Å 108.62Å 90.00° 119.24° 90.00°	Depositor
Resolution (Å)	31.47 – 1.68 31.47 – 1.68	Depositor EDS
% Data completeness (in resolution range)	87.7 (31.47-1.68) 87.7 (31.47-1.68)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 1.68Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.151 , 0.182 0.166 , 0.195	Depositor DCC
R_{free} test set	3316 reflections (1.82%)	wwPDB-VP
Wilson B-factor (Å ²)	23.9	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6853	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XE, NAG, CU, IMD, MG, TPQ, CL, BMA, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/6412	0.84	4/8732 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	119	PRO	O-C-N	6.16	124.14	121.31
1	A	666	PRO	O-C-N	5.66	123.81	121.15
1	A	660	THR	N-CA-C	5.51	117.29	111.28
1	A	725	ASN	N-CA-C	5.33	118.79	112.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6087	0	5600	22	0
2	B	39	0	34	1	0
3	A	70	0	65	0	0
4	A	1	0	0	0	0
5	A	2	0	0	0	0
6	A	3	0	0	0	0
7	A	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	5	0	5	0	0
9	A	8	0	0	2	0
10	A	632	0	0	3	0
All	All	6853	0	5704	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:463:GLN:NE2	10:A:2438:HOH:O	2.31	0.63
1:A:691:GLY:HA2	9:A:907:XE:XE	2.76	0.63
1:A:45:CYS:SG	1:A:758:VAL:HG13	2.39	0.62
1:A:553:LYS:HG2	10:A:2625:HOH:O	2.01	0.61
1:A:380[B]:LEU:HD21	1:A:480:TYR:CZ	2.38	0.58
1:A:451:LEU:C	1:A:451:LEU:HD12	2.31	0.56
1:A:380[B]:LEU:HD12	1:A:532:LEU:CD2	2.38	0.53
1:A:377:LEU:HD21	1:A:380[A]:LEU:HD11	1.92	0.52
1:A:380[B]:LEU:HD22	1:A:398:ASP:OD2	2.11	0.49
1:A:532:LEU:HD11	9:A:908:XE:XE	2.93	0.47
1:A:650:THR:HG23	10:A:2620:HOH:O	2.17	0.44
1:A:631:GLY:N	1:A:632:PRO:HD3	2.34	0.43
1:A:642:LEU:HD22	1:A:688:VAL:HG22	2.01	0.43
1:A:314:VAL:HG13	1:A:318:ASP:HB3	2.01	0.43
1:A:727:ARG:HA	1:A:727:ARG:CZ	2.49	0.42
1:A:231:GLY:HA3	1:A:251:PHE:CE1	2.54	0.42
1:A:504:ALA:CB	1:A:523:LEU:HD11	2.49	0.42
2:B:1:NAG:HO3	2:B:2:NAG:C1	2.33	0.42
1:A:449[B]:SER:HB3	1:A:462:THR:O	2.20	0.41
1:A:735:TYR:CD1	1:A:735:TYR:C	2.98	0.41
1:A:533:ASN:HB2	1:A:637:TRP:CE3	2.56	0.40
1:A:145:LEU:HA	1:A:146:PRO:C	2.47	0.40
1:A:250:PHE:HB2	1:A:269:ILE:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	776/747 (104%)	754 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	686/658 (104%)	678 (99%)	8 (1%)	63	43

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	310	ASP
1	A	388	ASP
1	A	642	LEU
1	A	703	ASN
1	A	725	ASN
1	A	739	ASP
1	A	740	GLU
1	A	769	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	173	HIS
1	A	187	ASN
1	A	464	ASN
1	A	579	ASN
1	A	590	ASN
1	A	592	GLN
1	A	661	ASN
1	A	731	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPQ	A	478	4,1	13,14,15	1.69	4 (30%)	13,19,21	1.05	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPQ	A	478	4,1	-	4/5/22/24	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	478	TPQ	O4-C4	-3.30	1.25	1.33
1	A	478	TPQ	C3-C4	2.63	1.40	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	478	TPQ	C6-C1	2.37	1.40	1.34
1	A	478	TPQ	C3-C2	2.02	1.50	1.44

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	478	TPQ	CB-C1-C2	2.36	122.69	118.56

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	478	TPQ	C-CA-CB-C1
1	A	478	TPQ	N-CA-CB-C1
1	A	478	TPQ	C2-C1-CB-CA
1	A	478	TPQ	C6-C1-CB-CA

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.71	0	17,19,21	1.03	1 (5%)
2	NAG	B	2	2	14,14,15	0.50	0	17,19,21	0.92	0
2	BMA	B	3	2	11,11,12	0.56	0	15,15,17	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	O5-C1-C2	-2.55	107.35	111.29

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	3	BMA	O5-C5-C6-O6
2	B	3	BMA	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	NAG	1	0
2	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 20 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1309	1	14,14,15	0.67	0	17,19,21	0.77	0
3	NAG	A	1434	1	14,14,15	0.60	0	17,19,21	1.01	1 (5%)
3	NAG	A	1191	1	14,14,15	0.38	0	17,19,21	1.06	0
8	IMD	A	821	-	5,5,5	0.69	0	5,5,5	0.58	0
3	NAG	A	1081[B]	1	14,14,15	0.53	0	17,19,21	0.75	0
3	NAG	A	1081[A]	1	14,14,15	0.47	0	17,19,21	0.84	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1309	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1434	1	-	2/6/23/26	0/1/1/1
3	NAG	A	1191	1	-	1/6/23/26	0/1/1/1
8	IMD	A	821	-	-	-	0/1/1/1
3	NAG	A	1081[B]	1	-	2/6/23/26	0/1/1/1
3	NAG	A	1081[A]	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1434	NAG	O5-C1-C2	-3.37	106.08	111.29
3	A	1081[A]	NAG	C1-O5-C5	2.05	114.93	112.19

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1081[B]	NAG	O5-C5-C6-O6
3	A	1191	NAG	O5-C5-C6-O6
3	A	1434	NAG	C8-C7-N2-C2
3	A	1434	NAG	O7-C7-N2-C2
3	A	1081[B]	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	734/747 (98%)	0.14	20 (2%) 56 63	14, 29, 50, 105	48 (6%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	776[A]	ILE	7.7
1	A	43	ALA	4.9
1	A	741	THR	4.6
1	A	769	THR	3.6
1	A	310	ASP	3.2
1	A	223	ALA	2.9
1	A	47	SER	2.7
1	A	774[A]	THR	2.6
1	A	46	VAL	2.6
1	A	770[A]	TYR	2.5
1	A	577	PHE	2.4
1	A	44	GLU	2.2
1	A	311	SER	2.2
1	A	773[A]	GLY	2.1
1	A	277	SER	2.1
1	A	45	CYS	2.1
1	A	212	CYS	2.1
1	A	565	THR	2.1
1	A	777[A]	ASN	2.1
1	A	282	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	TPQ	A	478	14/15	0.87	0.20	25,40,73,108	9

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BMA	B	3	11/12	0.62	0.16	70,85,97,105	0
2	NAG	B	2	14/15	0.81	0.13	39,52,78,88	0
2	NAG	B	1	14/15	0.90	0.11	31,48,62,63	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	1309	14/15	0.69	0.17	66,86,102,103	0
8	IMD	A	821	5/5	0.73	0.20	38,44,48,60	5
6	MG	A	805	1/1	0.75	0.21	37,37,37,37	1
3	NAG	A	1081[B]	14/15	0.77	0.19	30,35,50,57	14
3	NAG	A	1081[A]	14/15	0.77	0.19	35,56,70,80	14
3	NAG	A	1434	14/15	0.78	0.15	44,60,78,89	0
6	MG	A	806	1/1	0.79	0.17	59,59,59,59	1
7	CL	A	814	1/1	0.86	0.16	48,48,48,48	1
3	NAG	A	1191	14/15	0.89	0.11	39,53,71,75	0
9	XE	A	907	1/1	0.91	0.13	39,39,39,39	1
9	XE	A	906	1/1	0.92	0.09	43,43,43,43	1
7	CL	A	815	1/1	0.95	0.13	46,46,46,46	1
9	XE	A	908	1/1	0.95	0.09	39,39,39,39	1
7	CL	A	811	1/1	0.96	0.11	40,40,40,40	0
9	XE	A	905	1/1	0.97	0.06	38,38,38,38	1
7	CL	A	813	1/1	0.97	0.29	43,43,43,43	0
7	CL	A	816	1/1	0.98	0.34	49,49,49,49	1
7	CL	A	812	1/1	0.98	0.15	41,41,41,41	0
6	MG	A	804	1/1	0.98	0.04	31,31,31,31	0
9	XE	A	903	1/1	0.99	0.03	34,34,34,34	1
9	XE	A	904	1/1	0.99	0.05	38,38,38,38	1
4	CU	A	801	1/1	0.99	0.02	32,32,32,32	0
5	CA	A	803	1/1	0.99	0.04	25,25,25,25	0
9	XE	A	901	1/1	0.99	0.04	27,27,27,27	1
9	XE	A	902	1/1	0.99	0.03	41,41,41,41	1
5	CA	A	802	1/1	1.00	0.03	23,23,23,23	0

6.5 Other polymers ⓘ

There are no such residues in this entry.