



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:19 PM UTC

PDB ID : 2RKM / pdb_00002rkm
Title : STRUCTURE OF OPPA COMPLEXED WITH LYS-LYS
Authors : Sleight, S.H.; Tame, J.R.H.; Wilkinson, A.J.
Deposited on : 1997-03-25
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	517	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IUM	A	520	-	-	X	-
3	IUM	A	521	-	-	X	-
3	IUM	A	522	-	-	X	-
3	IUM	A	523	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called OLIGO-PEPTIDE BINDING PROTEIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	517	4189	2682	701	801	5	0	9	0

- Molecule 2 is LYSINE (CCD ID: LYS) (formula: $C_6H_{15}N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	9	6	2	1	0	0
2	A	1	10	6	2	2	0	0

- Molecule 3 is URANYL (VI) ION (CCD ID: IUM) (formula: O_2U).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	U	0	0
			3	2	1		
3	A	1	Total	O	U	0	0
			3	2	1		
3	A	1	Total	O	U	0	0
			3	2	1		
3	A	1	Total	O	U	0	0
			3	2	1		
3	A	1	Total	O	U	0	0
			3	2	1		
3	A	1	Total	O	U	0	0
			3	2	1		
3	A	1	Total	O	U	0	0
			2	1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	519	Total	O	0	0
			519	519		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: OLIGO-PEPTIDE BINDING PROTEIN

Chain A:  86% 13%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	110.47Å 77.15Å 71.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-1.80)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.167 , 0.219	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4750	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IUM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/4344	1.25	29/5922 (0.5%)

There are no bond length outliers.

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	329	GLU	CA-CB-CG	9.59	133.28	114.10
1	A	409	PHE	N-CA-C	7.31	118.25	108.38
1	A	485	TYR	CA-CB-CG	6.57	125.73	113.90
1	A	398	LYS	CA-CB-CG	6.42	126.93	114.10
1	A	224	LEU	CA-C-N	6.37	127.80	119.84
1	A	224	LEU	C-N-CA	6.37	127.80	119.84
1	A	303	VAL	N-CA-C	6.07	116.24	110.53
1	A	330	PRO	N-CA-CB	5.78	108.38	103.35
1	A	260	ILE	CA-C-N	5.75	125.88	119.32
1	A	260	ILE	C-N-CA	5.75	125.88	119.32
1	A	481	PRO	N-CA-CB	5.73	108.41	103.31
1	A	400	PHE	CA-CB-CG	5.56	119.36	113.80
1	A	319	PRO	CA-C-N	5.54	125.21	119.56
1	A	319	PRO	C-N-CA	5.54	125.21	119.56
1	A	306	VAL	N-CA-C	5.51	115.75	110.74
1	A	440	HIS	CA-CB-CG	-5.43	108.37	113.80
1	A	232	ASP	CA-CB-CG	5.42	118.02	112.60
1	A	161	HIS	CA-C-N	5.38	125.19	119.28
1	A	161	HIS	C-N-CA	5.38	125.19	119.28
1	A	159	LEU	N-CA-C	5.34	119.55	113.19
1	A	152	VAL	CA-C-N	5.34	125.15	119.28
1	A	152	VAL	C-N-CA	5.34	125.15	119.28
1	A	485	TYR	N-CA-C	5.24	117.88	110.50
1	A	87	PRO	N-CA-CB	5.19	107.87	103.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	207	ASN	CA-C-N	5.19	124.85	119.56
1	A	207	ASN	C-N-CA	5.19	124.85	119.56
1	A	144	PHE	CA-CB-CG	5.03	118.83	113.80
1	A	259	GLU	N-CA-C	5.01	117.86	111.69
1	A	151	PRO	N-CA-CB	5.01	107.77	103.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4189	0	4103	34	0
2	A	19	0	25	0	0
3	A	23	0	0	9	0
4	A	519	0	0	5	0
All	All	4750	0	4128	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:522:IUM:O1	3:A:522:IUM:U	1.32	1.08
3:A:522:IUM:U	3:A:522:IUM:O2	1.34	1.08
3:A:523:IUM:O1	3:A:523:IUM:U	1.49	0.93
1:A:403:THR:HG23	1:A:408:THR:HB	1.54	0.89
3:A:521:IUM:U	3:A:521:IUM:O1	1.53	0.88
3:A:521:IUM:U	3:A:521:IUM:O2	1.57	0.84
3:A:523:IUM:U	3:A:523:IUM:O2	1.57	0.84
3:A:520:IUM:O2	3:A:520:IUM:U	1.72	0.71
1:A:408:THR:HA	4:A:2494:HOH:O	1.92	0.68
3:A:520:IUM:U	3:A:520:IUM:O1	1.75	0.68
3:A:525:IUM:O1	3:A:525:IUM:U	1.76	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:ILE:HA	1:A:160[B]:VAL:HG13	1.79	0.65
1:A:459:ASP:OD1	1:A:462:ARG:NH1	2.30	0.64
1:A:301:ILE:HA	1:A:305:LYS:HG2	1.83	0.61
1:A:11:ASP:H	1:A:516:LYS:NZ	1.99	0.60
1:A:359:LEU:HB3	1:A:388:VAL:HG12	1.84	0.59
1:A:398:LYS:HD3	4:A:2038:HOH:O	2.04	0.57
1:A:10:ALA:HA	1:A:516:LYS:HD3	1.85	0.57
1:A:11:ASP:H	1:A:516:LYS:HZ3	1.52	0.57
1:A:57[B]:SER:HB3	1:A:58:PRO:HD2	1.92	0.51
1:A:174:LYS:HE3	4:A:2479:HOH:O	2.11	0.51
1:A:254:GLN:HG2	4:A:2468:HOH:O	2.11	0.50
1:A:291:THR:O	1:A:295[B]:LEU:HD23	2.11	0.50
1:A:346:LYS:HE2	1:A:350:GLU:OE2	2.11	0.50
1:A:34:VAL:HB	1:A:35:PRO:HD3	1.95	0.49
1:A:382:TRP:HB3	1:A:388:VAL:HG23	1.96	0.47
1:A:402:ASP:O	1:A:406:GLN:HG3	2.14	0.47
1:A:57[A]:SER:HB2	1:A:58:PRO:HD2	1.95	0.47
1:A:257:LYS:HE3	1:A:261:PRO:HB3	1.96	0.47
1:A:323:ASP:O	1:A:423:PRO:HD3	2.16	0.46
1:A:57[B]:SER:HB3	1:A:58:PRO:CD	2.46	0.45
1:A:248:MET:HA	1:A:248:MET:HE2	1.97	0.45
1:A:382:TRP:HB3	1:A:388:VAL:CG2	2.47	0.44
1:A:115:TYR:CE1	1:A:428:ASN:HB3	2.52	0.44
1:A:3:VAL:HA	1:A:4:PRO:HD3	1.90	0.43
1:A:257:LYS:HA	1:A:257:LYS:HD2	1.87	0.43
1:A:129:LYS:HD3	1:A:133:ASP:OD2	2.20	0.42
1:A:57[A]:SER:HB2	1:A:58:PRO:CD	2.49	0.42
1:A:43:LEU:O	1:A:187:THR:HB	2.20	0.41
1:A:356:ASP:O	1:A:357:LYS:HD2	2.21	0.41
1:A:300:ASP:OD1	1:A:304:ASN:ND2	2.50	0.41
1:A:117:HIS:HE1	4:A:2288:HOH:O	2.04	0.40
1:A:297[B]:LEU:HD11	1:A:378:VAL:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	524/517 (101%)	506 (97%)	17 (3%)	1 (0%)	43 31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	464/455 (102%)	458 (99%)	6 (1%)	61 54

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	174	LYS
1	A	258	LYS
1	A	357	LYS
1	A	398	LYS
1	A	455	LYS
1	A	468	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	HIS
1	A	184	ASN
1	A	199	ASN
1	A	262	ASN
1	A	279	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	308	ASN
1	A	338	GLN
1	A	406	GLN
1	A	436	ASN
1	A	440	HIS
1	A	472	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	IUM	A	520	-	0,2,2	-	-	-		
3	IUM	A	523	-	0,2,2	-	-	-		
2	LYS	A	2001	2	7,8,9	0.56	0	3,8,10	0.88	0
3	IUM	A	518	-	0,2,2	-	-	-		
3	IUM	A	519	-	0,2,2	-	-	-		
2	LYS	A	2002	2	8,9,9	0.64	0	7,10,10	0.44	0
3	IUM	A	521	-	0,2,2	-	-	-		
3	IUM	A	522	-	0,2,2	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	IUM	A	524	-	0,2,2	-	-	-		
3	IUM	A	525	-	0,1,2	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LYS	A	2001	2	-	1/6/7/9	-
2	LYS	A	2002	2	-	2/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2002	LYS	CA-CB-CG-CD
2	A	2002	LYS	CG-CD-CE-NZ
2	A	2001	LYS	N-CA-CB-CG

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	520	IUM	2	0
3	A	523	IUM	2	0
3	A	521	IUM	2	0
3	A	522	IUM	2	0
3	A	525	IUM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.