



Full wwPDB NMR Structure Validation Report ⓘ

Mar 5, 2026 – 02:05 PM UTC

PDB ID : 2RLQ / pdb_00002rlq
Title : NMR structure of CCP modules 2-3 of complement factor H
Authors : Hocking, H.G.; Herbert, A.P.; Pangburn, M.K.; Kavanagh, D.; Barlow, P.N.; Uhrin, D.
Deposited on : 2007-07-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	129	

2 Ensemble composition and analysis

This entry contains 29 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:84-A:159, A:168-A:206 (115)	0.77	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 6, 7, 8, 9, 10, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28
2	5, 29
Single-model clusters	11

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1831 atoms, of which 876 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Complement factor H.

Mol	Chain	Residues	Atoms						Trace
1	A	123	Total	C	H	N	O	S	0
			1831	599	876	155	191	10	

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	78	GLU	-	expression tag	UNP P08603
A	79	ALA	-	expression tag	UNP P08603
A	80	GLU	-	expression tag	UNP P08603
A	81	ALA	-	expression tag	UNP P08603
A	82	ALA	-	expression tag	UNP P08603
A	83	GLY	-	expression tag	UNP P08603

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Complement factor H




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Complement factor H




4.2.2 Score per residue for model 2

- Molecule 1: Complement factor H




4.2.3 Score per residue for model 3

- Molecule 1: Complement factor H




4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: Complement factor H




4.2.5 Score per residue for model 5

- Molecule 1: Complement factor H




4.2.6 Score per residue for model 6

- Molecule 1: Complement factor H



4.2.7 Score per residue for model 7

- Molecule 1: Complement factor H



V206

4.2.8 Score per residue for model 8

- Molecule 1: Complement factor H

Chain A:  78% 9% 6% 5%



4.2.9 Score per residue for model 9

- Molecule 1: Complement factor H

Chain A:  81% 9% 6% 5%



4.2.10 Score per residue for model 10

- Molecule 1: Complement factor H

Chain A:  81% 6% 6% 5%



4.2.11 Score per residue for model 11

- Molecule 1: Complement factor H

Chain A:  79% 10% 6% 5%



4.2.12 Score per residue for model 12

- Molecule 1: Complement factor H

Chain A:  77% 12% 6% 5%



4.2.13 Score per residue for model 13

- Molecule 1: Complement factor H

Chain A:  78% 11% 6% 5%



4.2.14 Score per residue for model 14

- Molecule 1: Complement factor H


Chain A:  71% 18% 6% 5%



4.2.15 Score per residue for model 15

- Molecule 1: Complement factor H


Chain A:  72% 17% 6% 5%



4.2.16 Score per residue for model 16


- Molecule 1: Complement factor H

Chain A:  76% 13% 6% 5%




4.2.17 Score per residue for model 17

- Molecule 1: Complement factor H




4.2.18 Score per residue for model 18

- Molecule 1: Complement factor H




4.2.19 Score per residue for model 19

- Molecule 1: Complement factor H




4.2.20 Score per residue for model 20

- Molecule 1: Complement factor H




4.2.21 Score per residue for model 21

- Molecule 1: Complement factor H




4.2.22 Score per residue for model 22

- Molecule 1: Complement factor H




4.2.23 Score per residue for model 23

- Molecule 1: Complement factor H




4.2.24 Score per residue for model 24

- Molecule 1: Complement factor H




4.2.25 Score per residue for model 25

- Molecule 1: Complement factor H



4.2.26 Score per residue for model 26

- Molecule 1: Complement factor H






4.2.27 Score per residue for model 27

- Molecule 1: Complement factor H

Chain A: 66% 21% 6% 5%



4.2.28 Score per residue for model 28

- Molecule 1: Complement factor H

Chain A: 73% 15% 6% 5%



4.2.29 Score per residue for model 29

- Molecule 1: Complement factor H

Chain A: 75% 13% 6% 5%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics, simulated annealing*.

Of the 100 calculated structures, 29 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.1
CNS	refinement	1.1
CYANA	structure solution	2.1

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.28±0.04	1±1/917 (0.1± 0.1%)	0.90±0.02	0±0/1243 (0.0± 0.0%)
All	All	1.28	26/26593 (0.1%)	0.91	2/36047 (0.0%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	101	GLY	N-CA	-6.75	1.38	1.45	11	2
1	A	101	GLY	C-N	-5.69	1.25	1.33	11	1
1	A	177	VAL	CA-C	-5.57	1.46	1.52	18	1
1	A	137	ASP	C-N	-5.56	1.27	1.33	11	4
1	A	184	ILE	N-CA	-5.47	1.39	1.46	20	2
1	A	178	CYS	N-CA	-5.47	1.39	1.46	15	3
1	A	142	GLU	C-N	-5.46	1.27	1.33	16	2
1	A	111	VAL	N-CA	-5.41	1.40	1.46	27	4
1	A	94	GLY	C-N	-5.41	1.26	1.33	15	1
1	A	142	GLU	CA-C	-5.26	1.46	1.52	16	1
1	A	179	ASN	C-N	-5.14	1.26	1.33	24	1
1	A	111	VAL	C-N	-5.14	1.26	1.33	21	1
1	A	177	VAL	C-N	-5.06	1.26	1.33	5	2
1	A	143	VAL	C-N	-5.03	1.28	1.33	27	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	169	HIS	CA-CB-CG	5.72	119.52	113.80	26	1
1	A	170	PHE	CA-CB-CG	5.04	118.83	113.80	7	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	892	821	818	7±3
All	All	25868	23809	23722	206

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:105:GLU:O	1:A:108:VAL:HG12	0.62	1.95	20	29
1:A:118:TYR:CD1	1:A:143:VAL:HA	0.59	2.33	28	6
1:A:106:TYR:HA	1:A:129:CYS:SG	0.56	2.40	7	3
1:A:184:ILE:HD13	1:A:188:GLU:HA	0.56	1.76	25	2
1:A:143:VAL:O	1:A:145:LYS:HG3	0.54	2.03	27	3
1:A:144:VAL:HG23	1:A:170:PHE:HB2	0.54	1.80	6	6
1:A:94:GLY:HA2	1:A:115:ASN:CG	0.54	2.28	29	9
1:A:99:THR:OG1	1:A:109:LYS:HB2	0.54	2.03	5	4
1:A:175:ARG:HA	1:A:188:GLU:O	0.54	2.03	19	2
1:A:149:VAL:HG12	1:A:150:THR:O	0.54	2.03	6	2
1:A:118:TYR:HE1	1:A:143:VAL:HG22	0.53	1.63	23	7
1:A:118:TYR:HB3	1:A:141:CYS:SG	0.53	2.43	28	8
1:A:144:VAL:CG2	1:A:170:PHE:HB2	0.53	2.34	2	9
1:A:118:TYR:CE1	1:A:143:VAL:HG22	0.53	2.38	23	6
1:A:158:VAL:HG12	1:A:177:VAL:HG22	0.51	1.83	25	6
1:A:182:TYR:HB3	1:A:205:CYS:SG	0.51	2.45	17	1
1:A:94:GLY:HA2	1:A:115:ASN:OD1	0.51	2.05	28	1
1:A:85:CYS:HB3	1:A:134:TRP:NE1	0.50	2.22	7	1
1:A:156:LYS:HG3	1:A:177:VAL:HG23	0.49	1.85	7	1
1:A:158:VAL:HG22	1:A:175:ARG:O	0.49	2.08	21	1
1:A:153:GLU:HG2	1:A:154:ASN:ND2	0.49	2.23	5	4
1:A:187:ASP:CG	1:A:200:LYS:HD3	0.48	2.33	18	1
1:A:98:LEU:HA	1:A:109:LYS:O	0.48	2.08	5	1
1:A:124:ILE:O	1:A:139:PRO:HB3	0.48	2.09	20	5
1:A:174:VAL:HG12	1:A:190:MET:CG	0.48	2.39	23	1
1:A:202:LYS:HE2	1:A:203:PRO:HD2	0.48	1.84	8	1
1:A:121:LEU:O	1:A:140:ILE:HB	0.48	2.09	27	5

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:189:GLU:O	1:A:200:LYS:HG3	0.47	2.10	8	2
1:A:200:LYS:HE3	1:A:201:GLU:O	0.47	2.10	19	1
1:A:85:CYS:SG	1:A:133:GLY:HA2	0.47	2.50	27	1
1:A:103:VAL:O	1:A:108:VAL:HG11	0.45	2.11	27	16
1:A:158:VAL:CG2	1:A:175:ARG:HB3	0.45	2.41	27	5
1:A:174:VAL:HG23	1:A:198:TRP:CH2	0.45	2.46	25	2
1:A:148:PRO:HA	1:A:168:TYR:CD2	0.45	2.47	29	1
1:A:157:ILE:HG12	1:A:176:PHE:CE2	0.45	2.46	24	1
1:A:146:CYS:HA	1:A:196:GLY:O	0.45	2.11	9	3
1:A:88:PRO:HD3	1:A:104:PHE:CD1	0.45	2.47	29	4
1:A:174:VAL:O	1:A:189:GLU:HA	0.45	2.11	4	1
1:A:183:LYS:O	1:A:205:CYS:HA	0.45	2.12	4	4
1:A:153:GLU:CD	1:A:153:GLU:H	0.45	2.20	21	4
1:A:147:LEU:HD12	1:A:148:PRO:HD2	0.44	1.89	23	1
1:A:110:ALA:O	1:A:126:TYR:HA	0.44	2.13	26	7
1:A:130:ASP:OD2	1:A:131:THR:HG22	0.44	2.13	6	1
1:A:153:GLU:OE2	1:A:204:LYS:HG3	0.43	2.13	21	1
1:A:112:TYR:CE2	1:A:141:CYS:SG	0.43	3.11	20	1
1:A:93:PHE:CB	1:A:141:CYS:HB2	0.43	2.44	25	1
1:A:117:GLY:C	1:A:144:VAL:HG22	0.43	2.37	6	2
1:A:117:GLY:HA2	1:A:170:PHE:CD2	0.43	2.48	12	1
1:A:183:LYS:HE2	1:A:184:ILE:O	0.43	2.14	10	1
1:A:201:GLU:H	1:A:201:GLU:CD	0.42	2.22	25	2
1:A:118:TYR:CE1	1:A:143:VAL:HG13	0.42	2.50	15	1
1:A:153:GLU:OE1	1:A:204:LYS:HG3	0.42	2.14	27	1
1:A:174:VAL:HG12	1:A:190:MET:HG3	0.42	1.90	23	1
1:A:119:GLN:OE1	1:A:142:GLU:HB3	0.42	2.15	28	1
1:A:183:LYS:HG2	1:A:206:VAL:OXT	0.42	2.15	18	1
1:A:157:ILE:HG12	1:A:176:PHE:CD2	0.41	2.50	4	1
1:A:130:ASP:OD1	1:A:131:THR:HG22	0.41	2.15	14	1
1:A:152:PRO:HD3	1:A:176:PHE:CD1	0.41	2.50	16	1
1:A:112:TYR:CB	1:A:139:PRO:HG2	0.41	2.45	20	2
1:A:200:LYS:HE2	1:A:200:LYS:HA	0.41	1.93	2	1
1:A:103:VAL:O	1:A:108:VAL:HG21	0.41	2.15	19	1
1:A:127:ARG:HA	1:A:136:ASN:HB2	0.40	1.92	14	1
1:A:193:SER:HB2	1:A:195:ASP:OD1	0.40	2.16	15	1
1:A:144:VAL:O	1:A:169:HIS:HA	0.40	2.17	15	1
1:A:145:LYS:HA	1:A:168:TYR:O	0.40	2.16	16	1
1:A:173:ALA:HA	1:A:190:MET:O	0.40	2.17	25	1
1:A:175:ARG:HD2	1:A:188:GLU:OE2	0.40	2.16	24	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	113/129 (88%)	99±2 (87±2%)	13±2 (12±2%)	1±1 (1±1%)	20	70
All	All	3277/3741 (88%)	2866 (87%)	389 (12%)	22 (1%)	20	70

All 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	150	THR	5
1	A	101	GLY	4
1	A	87	HIS	2
1	A	180	SER	2
1	A	195	ASP	2
1	A	103	VAL	1
1	A	152	PRO	1
1	A	151	ALA	1
1	A	185	GLU	1
1	A	179	ASN	1
1	A	186	GLY	1
1	A	194	ASP	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/107 (92%)	94±1 (95±2%)	4±1 (5±2%)	26	77
All	All	2842/3103 (92%)	2713 (95%)	129 (5%)	26	77

All 20 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	121	LEU	29
1	A	119	GLN	21
1	A	177	VAL	15
1	A	120	LEU	14
1	A	153	GLU	7
1	A	142	GLU	6
1	A	202	LYS	5
1	A	183	LYS	5
1	A	189	GLU	4
1	A	201	GLU	3
1	A	147	LEU	3
1	A	158	VAL	3
1	A	200	LYS	3
1	A	190	MET	3
1	A	172	GLN	3
1	A	204	LYS	1
1	A	85	CYS	1
1	A	195	ASP	1
1	A	174	VAL	1
1	A	184	ILE	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

There are no ligands in this entry.

6.7 Other polymers ⓘ

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided