



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:18 PM UTC

PDB ID : 5RL9 / pdb_00005rl9
Title : PanDDA analysis group deposition – Crystal Structure of SARS-CoV-2 heli-
case in complex with Z1703168683
Authors : Newman, J.A.; Yosaatmadja, Y.; Douangamath, A.; Aimon, A.; Powell, A.J.;
Dias, A.; Fearon, D.; Dunnett, L.; Brandao-Neto, J.; Krojer, T.; Skyner, R.;
Gorrie-Stone, T.; Thompson, W.; von Delft, F.; Arrowsmith, C.H.; Edwards,
A.; Bountra, C.; Gileadi, O.
Deposited on : 2020-09-16
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	601	22% 84% 10% 5%
1	B	601	12% 87% 9% ..

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Helicase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	572	Total	C	N	O	S	0	0	0
			4417	2816	737	832	32			
1	B	585	Total	C	N	O	S	0	1	0
			4508	2875	750	848	35			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		
2	B	3	Total	Zn	0	0
			3	3		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is 1-(3-fluoro-4-methylphenyl)methanesulfonamide (CCD ID: UR7) (formula: $C_8H_{10}FNO_2S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C F N O S 13 8 1 1 2 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	204	Total O 204 204	0	0
5	B	252	Total O 252 252	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Helicase



• Molecule 1: Helicase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	59.12Å 70.18Å 85.41Å 102.61° 96.08° 112.48°	Depositor
Resolution (Å)	81.49 – 1.79 81.49 – 1.79	Depositor EDS
% Data completeness (in resolution range)	91.1 (81.49-1.79) 91.1 (81.49-1.79)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 1.78Å)	Xtriage
Refinement program	BUSTER 2.10.3 (20-MAY-2020)	Depositor
R, R_{free}	0.177 , 0.231 (Not available) , 0.223	Depositor DCC
R_{free} test set	5321 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.439	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9420	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PO4, UR7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/4517 (0.0%)	1.09	9/6156 (0.1%)
1	B	0.88	1/4610 (0.0%)	1.10	14/6283 (0.2%)
All	All	0.87	2/9127 (0.0%)	1.09	23/12439 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	429	MET	SD-CE	-6.29	1.63	1.79
1	A	553	ALA	CA-C	5.10	1.59	1.52

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	86	ASN	CA-CB-CG	7.49	120.09	112.60
1	B	103	VAL	CA-C-N	6.83	129.99	120.29
1	B	103	VAL	C-N-CA	6.83	129.99	120.29
1	A	591	GLU	CA-C-N	6.34	133.11	121.70
1	A	591	GLU	C-N-CA	6.34	133.11	121.70
1	B	95	ASN	CA-CB-CG	6.34	118.94	112.60
1	A	561	PHE	CA-CB-CG	-6.27	107.53	113.80
1	B	561	PHE	CA-CB-CG	-6.26	107.54	113.80
1	B	361	ASN	CA-CB-CG	6.10	118.70	112.60
1	B	580	ASP	CA-CB-CG	5.93	118.53	112.60
1	A	530	THR	N-CA-CB	-5.73	100.63	111.00
1	B	532	THR	N-CA-C	-5.72	103.15	110.53
1	A	145	PHE	CA-CB-CG	-5.64	108.16	113.80
1	B	12	THR	N-CA-C	5.58	117.58	108.76
1	A	532	THR	N-CA-C	-5.43	103.22	110.55
1	B	45	VAL	N-CA-C	-5.36	105.16	110.62
1	B	350	SER	CA-C-N	5.27	128.07	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	350	SER	C-N-CA	5.27	128.07	120.38
1	A	405	LEU	N-CA-C	5.21	116.44	109.83
1	B	561	PHE	CA-C-N	5.03	126.98	120.44
1	B	561	PHE	C-N-CA	5.03	126.98	120.44
1	A	561	PHE	CA-C-N	5.02	126.96	120.44
1	A	561	PHE	C-N-CA	5.02	126.96	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4417	0	4321	31	0
1	B	4508	0	4424	29	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
4	B	13	0	0	0	0
5	A	204	0	0	2	0
5	B	252	0	0	1	0
All	All	9420	0	8745	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:12:THR:HG21	1:B:26:CYS:HA	1.47	0.95
1:A:467:LYS:NZ	5:A:801:HOH:O	2.00	0.93
1:A:158:LEU:HD21	1:A:164:HIS:ND1	1.86	0.90
1:B:12:THR:CG2	1:B:26:CYS:HA	2.07	0.84
1:A:442:ARG:HH11	1:A:464:HIS:CE1	2.04	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:LEU:HD11	1:B:234:PRO:HD3	1.69	0.73
1:A:158:LEU:HD21	1:A:164:HIS:CE1	2.26	0.70
1:A:255:THR:HG23	1:A:300:PRO:HG3	1.76	0.68
1:A:331:SER:HB2	1:A:353:GLU:HG3	1.79	0.64
1:A:176:LEU:HD22	1:A:200:PHE:HB2	1.81	0.62
1:B:474[B]:MET:HG2	1:B:590:LEU:HB2	1.86	0.58
1:B:228:THR:HG22	1:B:230:HIS:NE2	2.19	0.58
1:A:158:LEU:HD12	1:A:162:GLU:HB3	1.85	0.58
1:B:279:THR:HB	1:B:429:MET:HE2	1.84	0.58
1:A:215:THR:HG22	1:B:193:VAL:HG21	1.89	0.54
1:A:12:THR:HG21	1:A:26:CYS:HA	1.89	0.54
1:A:7:LEU:HD22	1:A:103:VAL:HG22	1.89	0.54
1:B:511:PHE:HB3	1:B:530:THR:HG22	1.90	0.54
1:B:7:LEU:HD21	1:B:106:PHE:HB2	1.90	0.53
1:B:371:VAL:HG23	1:B:393:ALA:HB2	1.92	0.52
1:A:7:LEU:HD21	1:A:106:PHE:HB2	1.91	0.51
1:B:155:ARG:HE	1:B:164:HIS:HD2	1.59	0.51
1:B:13:SER:O	1:B:44:SER:HA	2.11	0.50
1:A:279:THR:HB	1:A:429:MET:HE2	1.93	0.50
1:B:185:TYR:CE2	1:B:194:GLN:HG2	2.48	0.49
1:A:19:CYS:HB2	1:A:23:PRO:HD2	1.95	0.49
1:A:519:ASN:HB3	1:A:530:THR:HG23	1.94	0.49
1:B:228:THR:CG2	1:B:230:HIS:NE2	2.76	0.49
1:B:460:LYS:NZ	5:B:801:HOH:O	2.26	0.49
1:B:13:SER:HB3	1:B:92:LEU:H	1.78	0.49
1:B:158:LEU:HD13	1:B:164:HIS:HB2	1.97	0.47
1:A:519:ASN:HB3	1:A:530:THR:CG2	2.44	0.47
1:B:510:VAL:HG21	1:B:541:TYR:CD1	2.51	0.46
1:B:19:CYS:HB2	1:B:23:PRO:HD2	1.96	0.46
1:A:163:LEU:HD11	1:A:200:PHE:HE2	1.81	0.45
1:B:8:CYS:SG	1:B:99:GLY:N	2.89	0.45
1:A:445:PRO:HB3	1:A:468:SER:HB3	1.97	0.44
1:A:510:VAL:HG21	1:A:541:TYR:CD1	2.53	0.44
1:B:177:ASN:OD1	1:B:179:ASN:HB2	2.17	0.43
1:A:163:LEU:HG	1:A:211:TYR:CD2	2.54	0.43
1:A:551:GLU:HG3	1:A:577:SER:HB3	2.01	0.43
1:B:16:CYS:O	1:B:22:ARG:HA	2.20	0.42
1:B:445:PRO:HB3	1:B:468:SER:HB3	2.01	0.42
1:A:512:ILE:O	1:A:546:PHE:HA	2.19	0.42
1:B:185:TYR:HE2	1:B:194:GLN:HG2	1.85	0.42
1:A:14:LEU:HB2	1:A:25:LEU:O	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:279:THR:HB	1:B:429:MET:CE	2.48	0.42
1:A:163:LEU:HD11	1:A:200:PHE:CE2	2.55	0.41
1:B:512:ILE:O	1:B:546:PHE:HA	2.21	0.41
1:A:280:LEU:HD11	1:A:438:LEU:HG	2.03	0.41
1:A:320:LYS:HE3	1:A:324:TYR:HE1	1.86	0.41
1:A:252:LEU:HB3	1:A:299:TYR:CD2	2.57	0.40
1:A:505:ALA:O	1:A:508:LYS:HG2	2.20	0.40
1:B:269:TYR:CD1	1:B:295:LEU:HD13	2.56	0.40
1:B:551:GLU:HG3	1:B:577:SER:HB3	2.02	0.40
1:A:289:SER:HB3	5:A:975:HOH:O	2.20	0.40
1:A:19:CYS:CB	1:A:23:PRO:HD2	2.51	0.40
1:B:175:PRO:HD2	1:B:180:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	564/601 (94%)	546 (97%)	16 (3%)	2 (0%)	30	16
1	B	580/601 (96%)	560 (97%)	20 (3%)	0	100	100
All	All	1144/1202 (95%)	1106 (97%)	36 (3%)	2 (0%)	43	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	484	VAL
1	A	195	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	485/523 (93%)	464 (96%)	21 (4%)	26	7
1	B	498/523 (95%)	482 (97%)	16 (3%)	34	13
All	All	983/1046 (94%)	946 (96%)	37 (4%)	29	9

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	THR
1	A	35	ILE
1	A	46	ASN
1	A	69	SER
1	A	79	ILE
1	A	81	PHE
1	A	86	ASN
1	A	153	THR
1	A	162	GLU
1	A	173	ARG
1	A	209	VAL
1	A	215	THR
1	A	217	TYR
1	A	255	THR
1	A	416	THR
1	A	442	ARG
1	A	502	ARG
1	A	517	SER
1	A	530	THR
1	A	591	GLU
1	A	592	ILE
1	B	8	CYS
1	B	9	ASN
1	B	12	THR
1	B	13	SER
1	B	68	MET
1	B	92	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	95	ASN
1	B	103	VAL
1	B	124	ASN
1	B	191	SER
1	B	192	LYS
1	B	195	ILE
1	B	219	LEU
1	B	247	VAL
1	B	353	GLU
1	B	484	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	179	ASN
1	A	230	HIS
1	A	265	ASN
1	A	459	ASN
1	A	464	HIS
1	A	516	ASN
1	A	531	GLN
1	B	164	HIS
1	B	179	ASN
1	B	245	HIS
1	B	265	ASN
1	B	268	ASN
1	B	361	ASN
1	B	404	GLN
1	B	459	ASN
1	B	464	HIS
1	B	482	HIS
1	B	516	ASN
1	B	531	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	705	-	4,4,4	2.58	2 (50%)	6,6,6	0.83	0
4	UR7	B	701	-	12,13,13	0.39	0	18,19,19	0.76	0
3	PO4	B	705	-	4,4,4	2.56	1 (25%)	6,6,6	0.55	0
3	PO4	B	706	-	4,4,4	2.40	1 (25%)	6,6,6	1.20	0
3	PO4	A	704	-	4,4,4	2.53	1 (25%)	6,6,6	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UR7	B	701	-	-	0/5/5/5	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	705	PO4	P-O1	4.34	1.60	1.50
3	A	704	PO4	P-O1	4.20	1.60	1.50
3	B	706	PO4	P-O1	4.06	1.60	1.50
3	A	705	PO4	P-O1	4.06	1.60	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	705	PO4	P-O3	2.32	1.61	1.54

There are no bond angle outliers.


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	572/601 (95%)	1.20	134 (23%) 2 2	27, 58, 119, 149	0
1	B	585/601 (97%)	0.68	72 (12%) 8 9	19, 45, 95, 132	1 (0%)
All	All	1157/1202 (96%)	0.94	206 (17%) 4 4	19, 51, 110, 149	1 (0%)

All (206) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	219	LEU	7.0
1	A	592	ILE	5.9
1	B	340	VAL	5.7
1	A	225	PHE	5.7
1	A	1	ALA	5.3
1	A	167	TRP	5.0
1	A	228	THR	5.0
1	A	182	PHE	5.0
1	A	226	VAL	5.0
1	A	7	LEU	4.9
1	B	2	VAL	4.8
1	A	258	ILE	4.7
1	A	209	VAL	4.6
1	A	45	VAL	4.6
1	A	6	VAL	4.5
1	A	157	VAL	4.5
1	A	181	VAL	4.4
1	A	221	VAL	4.4
1	A	340	VAL	4.4
1	A	202	LYS	4.3
1	A	151	ILE	4.3
1	B	203	GLY	4.3
1	A	94	LYS	4.3
1	A	81	PHE	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	227	LEU	4.3
1	A	247	VAL	4.2
1	B	336	ALA	4.2
1	B	92	LEU	4.2
1	A	169	VAL	4.2
1	A	336	ALA	4.1
1	B	187	VAL	4.1
1	B	592	ILE	4.1
1	A	149	TYR	4.1
1	B	217	TYR	4.1
1	A	222	GLY	4.0
1	A	152	ALA	4.0
1	A	165	LEU	4.0
1	A	195	ILE	3.9
1	A	176	LEU	3.9
1	A	163	LEU	3.9
1	A	145	PHE	3.9
1	A	210	VAL	3.9
1	B	57	VAL	3.9
1	A	170	GLY	3.9
1	B	193	VAL	3.8
1	A	200	PHE	3.8
1	B	593	PRO	3.8
1	A	2	VAL	3.7
1	B	98	VAL	3.7
1	A	224	TYR	3.7
1	B	79	ILE	3.6
1	A	24	PHE	3.6
1	A	217	TYR	3.6
1	A	266	VAL	3.5
1	A	155	ARG	3.5
1	A	174	PRO	3.5
1	A	215	THR	3.5
1	A	185	TYR	3.5
1	A	208	ALA	3.4
1	B	45	VAL	3.4
1	A	43	LEU	3.4
1	A	256	LEU	3.4
1	B	90	PHE	3.3
1	A	27	CYS	3.3
1	A	241	VAL	3.3
1	B	214	THR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	229	SER	3.3
1	A	214	THR	3.2
1	B	81	PHE	3.2
1	A	153	THR	3.2
1	A	158	LEU	3.2
1	B	97	CYS	3.2
1	A	389	ALA	3.2
1	A	230	HIS	3.2
1	A	89	VAL	3.2
1	A	144	THR	3.2
1	A	92	LEU	3.1
1	A	48	TYR	3.1
1	B	158	LEU	3.1
1	B	12	THR	3.1
1	B	46	ASN	3.0
1	A	154	VAL	3.0
1	A	352	LEU	3.0
1	B	7	LEU	3.0
1	B	102	ASN	3.0
1	B	103	VAL	2.9
1	A	194	GLN	2.9
1	A	180	TYR	2.9
1	B	8	CYS	2.9
1	B	181	VAL	2.9
1	A	14	LEU	2.9
1	B	219	LEU	2.9
1	B	53	PRO	2.9
1	A	28	LYS	2.9
1	A	103	VAL	2.9
1	A	334	ILE	2.9
1	B	195	ILE	2.9
1	A	218	LYS	2.9
1	A	49	VAL	2.8
1	A	508	LYS	2.8
1	B	96	THR	2.8
1	B	215	THR	2.8
1	B	228	THR	2.8
1	A	223	ASP	2.8
1	A	42	VAL	2.8
1	A	179	ASN	2.8
1	A	12	THR	2.8
1	A	175	PRO	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	91	GLY	2.7
1	B	218	LYS	2.7
1	A	31	TYR	2.7
1	A	93	TYR	2.7
1	A	84	CYS	2.7
1	B	54	GLY	2.7
1	B	196	GLY	2.7
1	A	338	ALA	2.6
1	A	220	ASN	2.6
1	A	198	TYR	2.6
1	B	6	VAL	2.6
1	B	4	ALA	2.6
1	B	91	GLY	2.6
1	A	90	PHE	2.6
1	A	147	LEU	2.6
1	A	305	VAL	2.6
1	A	20	ILE	2.6
1	A	178	ARG	2.6
1	A	231	THR	2.5
1	A	351	THR	2.5
1	A	343	PHE	2.5
1	B	198	TYR	2.5
1	A	47	PRO	2.5
1	B	208	ALA	2.5
1	A	25	LEU	2.5
1	A	34	VAL	2.5
1	A	46	ASN	2.5
1	A	172	PRO	2.5
1	A	339	ARG	2.4
1	A	85	ALA	2.4
1	A	11	GLN	2.4
1	A	18	ALA	2.4
1	B	229	SER	2.4
1	A	250	THR	2.4
1	A	253	TYR	2.4
1	B	180	TYR	2.4
1	B	221	VAL	2.4
1	B	67	GLY	2.4
1	A	8	CYS	2.4
1	A	23	PRO	2.4
1	B	44	SER	2.4
1	B	161	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	150	GLY	2.3
1	B	231	THR	2.3
1	B	200	PHE	2.3
1	B	93	TYR	2.3
1	A	371	VAL	2.3
1	B	247	VAL	2.3
1	A	207	ASP	2.3
1	B	78	PRO	2.3
1	A	44	SER	2.3
1	A	255	THR	2.3
1	B	189	LYS	2.3
1	A	328	ASP	2.3
1	A	57	VAL	2.3
1	A	348	VAL	2.3
1	B	157	VAL	2.3
1	A	127	THR	2.2
1	B	341	GLU	2.2
1	A	330	CYS	2.2
1	B	63	LEU	2.2
1	A	346	PHE	2.2
1	B	24	PHE	2.2
1	B	66	GLY	2.2
1	A	249	ILE	2.2
1	A	9	ASN	2.2
1	A	183	THR	2.2
1	A	254	PRO	2.2
1	A	166	SER	2.2
1	B	100	SER	2.2
1	B	84	CYS	2.2
1	B	227	LEU	2.2
1	B	70	TYR	2.2
1	B	49	VAL	2.1
1	B	52	ALA	2.1
1	A	41	LEU	2.1
1	B	73	LYS	2.1
1	A	104	THR	2.1
1	A	216	THR	2.1
1	B	145	PHE	2.1
1	A	232	VAL	2.1
1	A	245	HIS	2.1
1	A	19	CYS	2.1
1	A	342	CYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	235	LEU	2.1
1	B	211	TYR	2.1
1	B	3	GLY	2.1
1	B	463	ALA	2.1
1	B	210	VAL	2.1
1	A	65	LEU	2.1
1	A	325	LEU	2.1
1	A	507	ARG	2.1
1	A	524	LYS	2.1
1	B	35	ILE	2.1
1	A	156	GLU	2.0
1	B	458	ASP	2.0
1	B	51	ASN	2.0
1	A	22	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	UR7	B	701	13/13	0.94	0.11	28,30,32,32	13
3	PO4	B	705	5/5	0.96	0.08	43,48,50,50	0
3	PO4	B	706	5/5	0.97	0.08	43,45,47,47	0
2	ZN	A	703	1/1	0.98	0.04	99,99,99,99	0
3	PO4	A	704	5/5	0.98	0.06	42,45,47,47	0
3	PO4	A	705	5/5	0.98	0.06	42,43,44,45	0
2	ZN	B	704	1/1	0.99	0.04	93,93,93,93	0
2	ZN	A	702	1/1	0.99	0.03	58,58,58,58	0
2	ZN	B	703	1/1	0.99	0.06	74,74,74,74	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	B	702	1/1	1.00	0.01	46,46,46,46	0
2	ZN	A	701	1/1	1.00	0.03	59,59,59,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.