



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:11 PM UTC

PDB ID : 6RMF / pdb_00006rmf
Title : Crystal structure of NDM-1 with VNRX-5133
Authors : Hinchliffe, P.; Spencer, J.; Brem, J.; Schofield, C.J.
Deposited on : 2019-05-06
Resolution : 1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5890 (1.54-1.50)
Clashscore	190562	6116 (1.54-1.50)
Ramachandran outliers	187476	6002 (1.54-1.50)
Sidechain outliers	187428	5999 (1.54-1.50)
RSRZ outliers	180081	5891 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	 6% 93%
1	B	244	 2% 89% 5% 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7424 atoms, of which 3437 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metallo-beta-lactamase type 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	234	Total	C	H	N	O	S	0	2	0
			3447	1094	1700	311	333	9			
1	B	228	Total	C	H	N	O	S	0	5	0
			3403	1080	1678	307	329	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	GLY	-	expression tag	UNP C7C422
A	28	SER	-	expression tag	UNP C7C422
B	27	GLY	-	expression tag	UNP C7C422
B	28	SER	-	expression tag	UNP C7C422

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is (10aR)-2-(((1r,4R)-4-((2-aminoethyl)amino)cyclohexyl)methyl)-6-carboxy-4-hydroxy-4,10a-dihydro-10H-benzo[5,6][1,2]oxaborinino[2,3-b][1,4,2]oxazaborol-4-uide (CCD ID: K9K) (formula: $C_{19}H_{27}BN_3O_5$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	B	C	H	N	O	
			43	1	17	19	1	5	0
4	B	1	Total	B	C	H	N	O	
			43	1	17	19	1	5	0

- Molecule 5 is (4 {R})-4-[2-[4-(2-azanylethylamino)cyclohexyl]ethanoylamino]-3,3-bis(oxidanyl)-2-oxa-3-boranuidabicyclo[4.4.0]deca-1(10),6,8-triene-10-carboxylic acid (CCD ID: K9B) (formula: C₁₉H₂₉BN₃O₆) (labeled as "Ligand of Interest" by depositor).





3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Metallo-beta-lactamase type 2



- Molecule 1: Metallo-beta-lactamase type 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.82Å 73.84Å 77.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.70 – 1.51 42.70 – 1.51	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.70-1.51) 100.0 (42.70-1.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.51Å)	Xtriage
Refinement program	PHENIX (1.15.2_3472: ???)	Depositor
R, R_{free}	0.146 , 0.172 0.146 , 0.172	Depositor DCC
R_{free} test set	3422 reflections (5.30%)	wwPDB-VP
Wilson B-factor (Å ²)	10.2	Xtriage
Anisotropy	0.627	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7424	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, K9K, K9B, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/1792	0.56	0/2438
1	B	0.34	0/1769	0.55	0/2407
All	All	0.37	0/3561	0.55	0/4845

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1747	1700	1687	4	0
1	B	1725	1678	1662	8	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	15	0	0	0	0
4	A	24	19	0	0	0
4	B	24	19	0	0	0
5	A	25	21	0	0	0
6	A	211	0	0	2	4
6	B	212	0	0	3	2
All	All	3987	3437	3349	12	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:ASP:OD1	6:B:401:HOH:O	1.82	0.94
1:A:223:ASP:OD2	6:A:401:HOH:O	2.06	0.72
1:B:223:ASP:OD2	6:B:402:HOH:O	2.14	0.65
1:A:126:MET:O	1:A:129:MET:HG2	2.09	0.52
1:B:126:MET:O	1:B:129:MET:HG2	2.13	0.48
1:B:218:LEU:CD1	1:B:269:LEU:HD11	2.45	0.47
1:B:104:TRP:NE1	1:B:108:GLU:HG3	2.31	0.45
1:A:242:LYS:HG2	6:A:546:HOH:O	2.17	0.44
1:B:41:THR:HG21	6:B:587:HOH:O	2.17	0.43
1:A:36:GLY:HA3	1:A:74:ALA:HB3	2.01	0.42
1:B:218:LEU:HD12	1:B:269:LEU:HD11	2.03	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:513:HOH:O	6:A:516:HOH:O[2_554]	2.02	0.18
6:A:604:HOH:O	6:B:566:HOH:O[3_645]	2.11	0.09
6:A:477:HOH:O	6:A:489:HOH:O[2_554]	2.13	0.07
6:A:432:HOH:O	6:B:445:HOH:O[1_545]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/244 (95%)	230 (99%)	2 (1%)	0	100	100
1	B	229/244 (94%)	227 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	461/488 (94%)	457 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/183 (97%)	177 (99%)	1 (1%)	78	61
1	B	176/183 (96%)	173 (98%)	3 (2%)	53	24
All	All	354/366 (97%)	350 (99%)	4 (1%)	63	40

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	193	ASN
1	B	193	ASN
1	B	199	ASP
1	B	212	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	107	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	305	-	4,4,4	0.24	0	6,6,6	0.13	0
4	K9K	B	303	2	26,27,31	2.36	5 (19%)	27,40,45	1.82	7 (25%)
3	SO4	A	303	-	4,4,4	0.39	0	6,6,6	0.36	0
3	SO4	A	304	-	4,4,4	0.26	0	6,6,6	0.25	0
4	K9K	A	306[B]	2	26,27,31	2.30	5 (19%)	27,40,45	1.87	7 (25%)
5	K9B	A	307[A]	2	27,27,31	2.15	10 (37%)	28,39,44	1.80	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	K9K	A	306[B]	2	-	4/8/40/46	0/3/4/4
4	K9K	B	303	2	-	5/8/40/46	0/3/4/4
5	K9B	A	307[A]	2	-	2/11/35/41	0/2/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	306[B]	K9K	O15-C10	7.84	1.37	1.28
4	B	303	K9K	O15-C10	7.52	1.37	1.28
4	B	303	K9K	C10-N11	5.96	1.33	1.27
4	A	306[B]	K9K	C10-N11	5.56	1.32	1.27
5	A	307[A]	K9B	B05-C04	5.26	1.69	1.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	306[B]	K9K	B13-O14	-4.50	1.38	1.47
4	B	303	K9K	C26-C12	-3.91	1.49	1.53
5	A	307[A]	K9B	C02-N03	3.71	1.41	1.34
5	A	307[A]	K9B	C15-C14	3.43	1.44	1.38
5	A	307[A]	K9B	C16-C17	3.08	1.44	1.39
5	A	307[A]	K9B	O08-C09	2.91	1.40	1.35
4	B	303	K9K	B13-O14	-2.80	1.42	1.47
5	A	307[A]	K9B	O13-C11	2.65	1.30	1.22
5	A	307[A]	K9B	C15-C16	2.58	1.43	1.38
5	A	307[A]	K9B	C29-C20	2.46	1.59	1.52
4	A	306[B]	K9K	O16-C17	2.44	1.39	1.35
4	B	303	K9K	C26-C25	2.33	1.55	1.51
5	A	307[A]	K9B	C19-C02	-2.32	1.46	1.51
5	A	307[A]	K9B	B05-O06	-2.23	1.43	1.47
4	A	306[B]	K9K	C26-C12	-2.02	1.51	1.53

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	303	K9K	C08-C09-C10	4.43	124.56	114.00
5	A	307[A]	K9B	O12-C11-O13	-3.97	114.82	123.35
4	A	306[B]	K9K	O16-C17-C18	-3.87	116.19	118.82
4	B	303	K9K	O16-C17-C18	-3.59	116.38	118.82
4	A	306[B]	K9K	C08-C09-C10	3.54	122.43	114.00
5	A	307[A]	K9B	C19-C02-N03	3.46	120.98	116.25
4	A	306[B]	K9K	C05-C06-C07	3.42	118.45	111.42
4	A	306[B]	K9K	C12-N11-C10	3.30	108.86	106.65
4	A	306[B]	K9K	C06-C07-C08	3.24	118.89	112.08
4	A	306[B]	K9K	B13-O16-C17	-3.23	112.08	121.15
5	A	307[A]	K9B	O08-C09-C10	3.19	121.00	118.82
4	B	303	K9K	C05-C28-C27	3.04	117.66	111.42
5	A	307[A]	K9B	C21-C20-C19	-3.01	107.56	111.48
5	A	307[A]	K9B	O01-C02-C19	-2.96	117.20	121.54
4	B	303	K9K	C12-N11-C10	2.72	108.47	106.65
4	B	303	K9K	C18-C17-C25	2.55	123.53	119.46
4	B	303	K9K	C07-C08-C09	2.42	114.63	111.48
5	A	307[A]	K9B	C18-C17-C09	-2.39	114.97	119.78
4	A	306[B]	K9K	O14-B13-O15	-2.35	102.82	109.15
5	A	307[A]	K9B	O12-C11-C10	2.31	121.84	115.28
4	B	303	K9K	O14-B13-O15	-2.04	103.66	109.15

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	306[B]	K9K	C07-C08-C09-C10
4	B	303	K9K	C07-C08-C09-C10
4	B	303	K9K	C17-C18-C19-O21
4	B	303	K9K	C17-C18-C19-O20
4	A	306[B]	K9K	C27-C08-C09-C10
4	A	306[B]	K9K	C17-C18-C19-O21
5	A	307[A]	K9B	C09-C10-C11-O12
4	A	306[B]	K9K	C17-C18-C19-O20
5	A	307[A]	K9B	C09-C10-C11-O13
4	B	303	K9K	C22-C18-C19-O21
4	B	303	K9K	C22-C18-C19-O20

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand K9K B 303



Ligand K9K A 306 (B)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	234/244 (95%)	-0.19	15 (6%)	25 30	7, 13, 47, 72	1 (0%)
1	B	228/244 (93%)	-0.26	6 (2%)	57 63	6, 14, 40, 66	4 (1%)
All	All	462/488 (94%)	-0.23	21 (4%)	38 44	6, 13, 43, 72	5 (1%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	34	THR	4.7
1	A	35	ILE	4.4
1	A	33	PRO	3.5
1	B	71	GLY	3.0
1	A	242	LYS	2.8
1	A	90	ASP	2.7
1	A	71	GLY	2.6
1	B	67	MET	2.5
1	A	38	GLN	2.5
1	B	270	ARG	2.4
1	A	39	MET	2.4
1	A	41	THR	2.3
1	A	66	ASP	2.2
1	A	72	ALA	2.2
1	A	42	GLY	2.2
1	B	41	THR	2.1
1	A	270	ARG	2.1
1	B	66	ASP	2.1
1	B	90	ASP	2.1
1	A	37	GLN	2.1
1	A	36	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	305	5/5	0.90	0.11	65,65,65,65	0
5	K9B	A	307[A]	25/29	0.92	0.09	10,23,31,40	46
4	K9K	B	303	24/28	0.97	0.07	10,16,39,42	0
4	K9K	A	306[B]	24/28	0.97	0.06	5,16,32,32	43
3	SO4	A	303	5/5	0.98	0.08	19,20,21,22	0
3	SO4	A	304	5/5	0.98	0.06	22,22,24,25	0
2	ZN	B	301	1/1	1.00	0.01	9,9,9,9	0
2	ZN	B	302	1/1	1.00	0.02	11,11,11,11	0
2	ZN	A	301	1/1	1.00	0.01	11,11,11,11	0
2	ZN	A	302	1/1	1.00	0.01	10,10,10,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around K9B A 307 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K9K B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around K9K A 306 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.