



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:09 PM UTC

PDB ID : 7RMM / pdb_00007rmm
Title : Structure of N74D mutant of disulfide stabilized HIV-1 CA hexamer
Authors : Bester, S.M.; Kvaratskhelia, M.
Deposited on : 2021-07-27
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	10% 88% 5% 7%
1	B	231	9% 84% 7% 8%
1	C	231	10% 77% 8% 14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5243 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CAPSID PROTEIN P24.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	215	Total	C	N	O	S	0	3	0
			1679	1053	295	316	15			
1	B	213	Total	C	N	O	S	0	1	0
			1656	1042	289	311	14			
1	C	198	Total	C	N	O	S	0	0	0
			1523	962	262	286	13			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	14	CYS	ALA	engineered mutation	UNP B6DRA0
A	45	CYS	GLU	engineered mutation	UNP B6DRA0
A	74	ASP	ASN	engineered mutation	UNP B6DRA0
A	184	ALA	TRP	engineered mutation	UNP B6DRA0
A	185	ALA	MET	engineered mutation	UNP B6DRA0
B	14	CYS	ALA	engineered mutation	UNP B6DRA0
B	45	CYS	GLU	engineered mutation	UNP B6DRA0
B	74	ASP	ASN	engineered mutation	UNP B6DRA0
B	184	ALA	TRP	engineered mutation	UNP B6DRA0
B	185	ALA	MET	engineered mutation	UNP B6DRA0
C	14	CYS	ALA	engineered mutation	UNP B6DRA0
C	45	CYS	GLU	engineered mutation	UNP B6DRA0
C	74	ASP	ASN	engineered mutation	UNP B6DRA0
C	184	ALA	TRP	engineered mutation	UNP B6DRA0
C	185	ALA	MET	engineered mutation	UNP B6DRA0

- Molecule 2 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total I 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total I 2 2	0	0
2	C	2	Total I 2 2	0	0

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	6	Total Cl 6 6	0	0
3	B	6	Total Cl 6 6	0	0
3	C	5	Total Cl 5 5	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	127	Total O 127 127	0	0
4	B	132	Total O 132 132	0	0
4	C	103	Total O 103 103	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CAPSID PROTEIN P24



• Molecule 1: CAPSID PROTEIN P24



• Molecule 1: CAPSID PROTEIN P24



4 Data and refinement statistics

Property	Value	Source
Space group	P 6	Depositor
Cell constants a, b, c, α , β , γ	157.36Å 157.36Å 56.58Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.43 – 1.97 45.43 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.43-1.97) 99.9 (45.43-1.97)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.97Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.218 , 0.246 0.218 , 0.246	Depositor DCC
R_{free} test set	2788 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.042 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5243	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.7671e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/1713	0.47	0/2327
1	B	0.29	0/1690	0.47	0/2294
1	C	0.27	0/1555	0.46	0/2112
All	All	0.28	0/4958	0.47	0/6733

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1679	0	1658	6	0
1	B	1656	0	1649	13	0
1	C	1523	0	1499	11	0
2	A	2	0	0	0	0
2	B	2	0	0	1	0
2	C	2	0	0	0	0
3	A	6	0	0	0	0
3	B	6	0	0	0	0
3	C	5	0	0	0	0
4	A	127	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	132	0	0	1	0
4	C	103	0	0	1	0
All	All	5243	0	4806	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:37:ILE:HD12	1:B:138:LEU:HB3	1.84	0.59
1:C:2:ILE:HG22	1:C:10:MET:HE3	1.86	0.56
1:C:37:ILE:HD12	1:C:138:LEU:HB3	1.86	0.56
1:A:192:GLN:HA	1:A:199:LYS:HE3	1.90	0.54
1:B:2:ILE:HG22	1:B:10:MET:HE3	1.90	0.53
1:B:2:ILE:HD11	1:B:115:ILE:HG12	1.92	0.52
1:A:37:ILE:HB	1:A:38:PRO:HD3	1.92	0.51
1:A:172:LEU:HD23	1:A:182:LYS:HG3	1.92	0.51
1:A:37:ILE:HD12	1:A:138:LEU:HB3	1.94	0.50
1:B:26:VAL:HG21	1:B:39:MET:HG2	1.95	0.48
1:B:37:ILE:HB	1:B:38:PRO:HD3	1.96	0.48
1:A:202:LEU:HD13	1:A:214:MET:HE3	1.96	0.47
1:C:47:ALA:HB1	1:C:51:ASP:HB2	1.96	0.47
1:C:202:LEU:HD13	1:C:214:MET:HE3	1.97	0.46
1:B:172:LEU:HG	1:B:177:ALA:HB3	1.98	0.45
1:C:154:ARG:HG2	1:C:193:ASN:HB3	1.97	0.45
1:C:144:MET:HE3	1:C:144:MET:HB2	1.76	0.45
1:C:136:LEU:O	1:C:140:LYS:HG3	2.17	0.45
1:C:156:GLY:HA3	4:C:455:HOH:O	2.17	0.45
1:A:23:TRP:CZ3	1:A:40:PHE:HB2	2.54	0.43
1:B:96:MET:HE2	4:B:451:HOH:O	2.19	0.42
1:B:96:MET:HE3	1:B:96:MET:HB2	1.91	0.42
1:B:23:TRP:CZ3	1:B:40:PHE:HB2	2.55	0.42
1:C:192:GLN:HA	1:C:199:LYS:HE2	2.02	0.42
1:B:202:LEU:HD13	1:B:214:MET:HE3	2.02	0.41
1:B:25:LYS:HD3	1:B:25:LYS:HA	1.90	0.41
1:C:153:ILE:HG21	1:C:168:PHE:HA	2.02	0.41
1:C:203:LYS:HD3	1:C:203:LYS:HA	1.88	0.41
1:B:144:MET:HE3	1:B:144:MET:HB2	1.93	0.40
1:B:18:ARG:NE	2:B:302:IOD:I	3.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/231 (93%)	211 (99%)	3 (1%)	0	100	100
1	B	210/231 (91%)	208 (99%)	2 (1%)	0	100	100
1	C	190/231 (82%)	189 (100%)	1 (0%)	0	100	100
All	All	614/693 (89%)	608 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/193 (94%)	181 (100%)	0	100	100
1	B	180/193 (93%)	179 (99%)	1 (1%)	78	76
1	C	165/193 (86%)	165 (100%)	0	100	100
All	All	526/579 (91%)	525 (100%)	1 (0%)	87	88

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	26	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	HIS
1	A	179	GLN
1	B	62	HIS
1	B	84	HIS
1	B	120	HIS
1	C	50	GLN
1	C	84	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 23 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	215/231 (93%)	0.55	23 (10%)	11 15	16, 34, 75, 97	3 (1%)
1	B	213/231 (92%)	0.52	21 (9%)	13 18	19, 34, 65, 78	1 (0%)
1	C	198/231 (85%)	0.65	22 (11%)	10 14	25, 37, 69, 80	0
All	All	626/693 (90%)	0.57	66 (10%)	11 15	16, 35, 70, 97	4 (0%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	96	MET	5.5
1	A	91	ILE	4.9
1	B	185	ALA	4.5
1	A	86	VAL	4.4
1	B	188	THR	4.3
1	A	205	LEU	4.2
1	A	6	LEU	4.1
1	B	177	ALA	4.0
1	A	207	PRO	4.0
1	B	184	ALA	3.9
1	B	179	GLN	3.8
1	B	178	SER	3.7
1	C	220	GLY	3.6
1	C	176	GLN	3.6
1	A	209	ALA	3.5
1	C	86	VAL	3.5
1	B	181	VAL	3.4
1	A	206	GLY	3.4
1	A	203	LYS	3.3
1	C	207	PRO	3.2
1	B	93	PRO	3.1
1	B	8	GLY	3.1
1	A	204	ALA	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	157	PRO	3.0
1	C	189	LEU	2.9
1	B	187	GLU	2.9
1	B	186	THR	2.9
1	B	86	VAL	2.8
1	A	182	LYS	2.8
1	B	83	LEU	2.8
1	A	85	PRO	2.8
1	A	177	ALA	2.7
1	C	149	SER	2.7
1	A	7	GLN	2.7
1	C	11	VAL	2.7
1	B	6	LEU	2.7
1	C	2	ILE	2.7
1	C	93	PRO	2.6
1	C	150	ILE	2.5
1	A	178	SER	2.5
1	C	95	GLN	2.5
1	B	183	ASN	2.5
1	C	5	ASN	2.4
1	C	147	PRO	2.4
1	A	200	THR	2.4
1	A	175	GLU	2.4
1	B	7	GLN	2.4
1	A	191	VAL	2.4
1	A	145	TYR	2.3
1	C	148	THR	2.3
1	C	3	VAL	2.3
1	C	121	ASN	2.3
1	C	10	MET	2.3
1	A	208	GLY	2.2
1	C	94	GLY	2.2
1	B	94	GLY	2.2
1	B	96	MET	2.1
1	C	175	GLU	2.1
1	B	176	GLN	2.1
1	A	11	VAL	2.1
1	B	182	LYS	2.1
1	A	219	GLN	2.1
1	A	201	ILE	2.1
1	B	4	GLN	2.1
1	A	157	PRO	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	151	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	A	306	1/1	0.86	0.14	61,61,61,61	0
3	CL	B	303	1/1	0.86	0.13	61,61,61,61	0
3	CL	A	303	1/1	0.91	0.10	59,59,59,59	0
3	CL	B	305	1/1	0.91	0.11	64,64,64,64	0
2	IOD	A	302	1/1	0.92	0.17	75,75,75,75	0
2	IOD	C	302	1/1	0.92	0.26	82,82,82,82	0
3	CL	C	306	1/1	0.94	0.08	53,53,53,53	0
3	CL	B	306	1/1	0.95	0.07	49,49,49,49	0
3	CL	C	304	1/1	0.95	0.15	44,44,44,44	0
2	IOD	B	302	1/1	0.95	0.20	67,67,67,67	0
3	CL	A	307	1/1	0.97	0.05	42,42,42,42	0
3	CL	A	308	1/1	0.98	0.15	25,25,25,25	0
3	CL	B	307	1/1	0.98	0.10	34,34,34,34	0
3	CL	A	304	1/1	0.98	0.08	32,32,32,32	0
3	CL	C	305	1/1	0.98	0.11	31,31,31,31	0
3	CL	A	305	1/1	0.98	0.09	31,31,31,31	0
3	CL	C	307	1/1	0.98	0.15	20,20,20,20	0
3	CL	B	308	1/1	0.99	0.12	15,15,15,15	0
3	CL	C	303	1/1	0.99	0.04	31,31,31,31	0
2	IOD	B	301	1/1	1.00	0.02	33,33,33,33	0
2	IOD	A	301	1/1	1.00	0.11	46,46,46,46	0
3	CL	B	304	1/1	1.00	0.14	21,21,21,21	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IOD	C	301	1/1	1.00	0.04	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.