



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:42 PM UTC

PDB ID : 8RMW / pdb_00008rmw
Title : Alpha-Methylacyl-CoA racemase from Mycobacterium tuberculosis.
Authors : Mojanaga, O.O.; Acharya, K.R.; Lloyd, M.D.
Deposited on : 2024-01-09
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	6% 91% 6% .
1	B	365	4% 91% 6% ...

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11138 atoms, of which 5311 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-methylacyl-CoA racemase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	354	Total	C	H	N	O	S	56	3	0
			5333	1690	2641	481	505	16			
1	B	354	Total	C	H	N	O	S	56	5	0
			5337	1691	2642	482	506	16			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP A0A045IZ15
A	361	GLY	-	expression tag	UNP A0A045IZ15
A	362	SER	-	expression tag	UNP A0A045IZ15
A	363	GLY	-	expression tag	UNP A0A045IZ15
A	364	CYS	-	expression tag	UNP A0A045IZ15
B	0	MET	-	initiating methionine	UNP A0A045IZ15
B	361	GLY	-	expression tag	UNP A0A045IZ15
B	362	SER	-	expression tag	UNP A0A045IZ15
B	363	GLY	-	expression tag	UNP A0A045IZ15
B	364	CYS	-	expression tag	UNP A0A045IZ15

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	1	0
			10	2	6	2		
2	B	1	Total	C	H	O	1	0
			10	2	6	2		
2	B	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	1	0
			17	4	10	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	206	Total 206	O 206	0	0
4	B	215	Total 215	O 215	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-methylacyl-CoA racemase



- Molecule 1: Alpha-methylacyl-CoA racemase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	181.05Å 79.09Å 59.19Å 90.00° 92.01° 90.00°	Depositor
Resolution (Å)	90.47 – 1.65 90.47 – 1.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (90.47-1.65) 99.9 (90.47-1.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.14 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.8.0419, REFMAC 5.8.0419	Depositor
R, R_{free}	0.194 , 0.225 0.195 , 0.226	Depositor DCC
R_{free} test set	5093 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.110 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11138	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/2767	1.11	5/3763 (0.1%)
1	B	0.72	0/2780	1.11	11/3781 (0.3%)
All	All	0.71	0/5547	1.11	16/7544 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	28	ASP	CA-CB-CG	6.33	118.94	112.60
1	A	54	ARG	CB-CA-C	-6.31	97.81	110.30
1	B	54	ARG	CD-NE-CZ	6.29	133.21	124.40
1	A	28	ASP	CA-CB-CG	6.24	118.84	112.60
1	B	123	GLN	N-CA-CB	-6.24	100.47	110.63
1	B	127	ASP	CA-CB-CG	6.17	118.77	112.60
1	A	306	GLU	CB-CA-C	-5.98	99.24	109.65
1	B	54	ARG	CB-CA-C	-5.96	96.97	110.07
1	B	85	ARG	CD-NE-CZ	5.95	132.73	124.40
1	A	276	THR	CA-CB-OG1	-5.69	101.06	109.60
1	B	348	THR	CA-CB-OG1	-5.54	101.30	109.60
1	B	255	ASP	CA-CB-CG	5.51	118.11	112.60
1	A	356	THR	CA-CB-OG1	-5.39	101.51	109.60
1	B	198	MET	CG-SD-CE	5.21	112.36	100.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	190	ASP	CA-CB-CG	5.07	117.67	112.60
1	B	85	ARG	N-CA-CB	5.00	117.84	109.98

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	212	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2692	2641	2635	13	0
1	B	2695	2642	2630	7	0
2	A	4	6	6	0	0
2	B	8	12	12	0	0
3	A	7	10	10	0	0
4	A	206	0	0	2	0
4	B	215	0	0	0	0
All	All	5827	5311	5293	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:MET:HE3	1:A:202:MET:CE	2.24	0.68
1:B:138:HIS:O	1:B:212:ARG:HD3	2.01	0.61
1:A:199:MET:HE3	1:A:202:MET:HE1	1.82	0.60
1:A:270:GLU:OE1	4:A:501:HOH:O	2.17	0.59
1:B:16:ILE:O	1:B:18:PRO:HD2	2.06	0.55
1:A:259:LEU:HD22	1:A:274:LEU:HD13	1.89	0.54
1:B:259:LEU:HD22	1:B:274:LEU:HD13	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:196:ILE:HG12	1:B:199:MET:HB2	2.00	0.44
1:B:212:ARG:HB2	1:B:212:ARG:NH1	2.33	0.43
1:B:113:GLY:HA3	1:B:130:TYR:CZ	2.54	0.43
1:A:199:MET:HE3	1:A:202:MET:HE2	1.99	0.42
1:A:260:PRO:HG2	1:A:271:LEU:HB2	2.01	0.42
1:A:230:ALA:HB3	1:A:283[B]:ASP:OD1	2.19	0.42
1:A:151:LEU:HA	1:B:217:LEU:O	2.20	0.41
1:A:196:ILE:HG12	1:A:199:MET:HB2	2.02	0.41
1:A:16:ILE:O	1:A:18:PRO:HD2	2.21	0.41
1:A:113:GLY:HA3	1:A:130:TYR:CZ	2.55	0.41
1:A:176:GLN:NE2	4:A:516:HOH:O	2.53	0.40
1:A:105:ARG:HG2	1:A:105:ARG:HH11	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	353/365 (97%)	339 (96%)	14 (4%)	0	100	100
1	B	355/365 (97%)	343 (97%)	11 (3%)	1 (0%)	36	21
All	All	708/730 (97%)	682 (96%)	25 (4%)	1 (0%)	48	30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	151	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/278 (99%)	272 (99%)	2 (1%)	76	64
1	B	275/278 (99%)	272 (99%)	3 (1%)	65	48
All	All	549/556 (99%)	544 (99%)	5 (1%)	70	56

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	ARG
1	A	199	MET
1	B	133	LEU
1	B	199	MET
1	B	340	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	176	GLN
1	A	263	ASN
1	A	282	HIS
1	A	327	GLN
1	B	309	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	401	-	3,3,3	0.24	0	2,2,2	0.13	0
3	PEG	A	402	-	6,6,6	0.46	0	5,5,5	0.56	0
2	EDO	B	402	-	3,3,3	0.76	0	2,2,2	0.29	0
2	EDO	B	401	-	3,3,3	0.40	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	401	-	-	0/1/1/1	-
3	PEG	A	402	-	-	2/4/4/4	-
2	EDO	B	402	-	-	0/1/1/1	-
2	EDO	B	401	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	PEG	C1-C2-O2-C3
3	A	402	PEG	O2-C3-C4-O4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/365 (96%)	0.25	23 (6%) 25 28	17, 30, 59, 92	3 (0%)
1	B	354/365 (96%)	0.20	14 (3%) 42 46	17, 29, 57, 87	5 (1%)
All	All	708/730 (96%)	0.23	37 (5%) 33 36	17, 30, 58, 92	8 (1%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	323	ASN	5.3
1	A	324	GLY	4.7
1	B	324	GLY	4.6
1	B	39	PRO	4.4
1	A	45	ILE	4.3
1	B	16	ILE	4.2
1	A	49	ALA	4.0
1	A	16	ILE	4.0
1	A	39	PRO	3.9
1	B	325	GLY	3.8
1	A	47	ARG	3.7
1	A	349	ILE	3.5
1	B	17	GLY	3.2
1	B	323	ASN	3.1
1	B	49	ALA	3.0
1	A	322	ALA	2.9
1	B	349	ILE	2.9
1	A	350	ASP	2.9
1	B	359	ASP	2.9
1	B	351	ILE	2.8
1	A	38	ARG	2.7
1	A	176	GLN	2.7
1	B	69	LEU	2.7
1	A	325	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	47	ARG	2.6
1	A	346	ALA	2.5
1	A	347	ALA	2.4
1	A	50	MET	2.3
1	A	17	GLY	2.3
1	B	355	LEU	2.2
1	A	345	PRO	2.2
1	A	62	LYS	2.2
1	A	48	ASP	2.2
1	A	353	ALA	2.1
1	A	351	ILE	2.1
1	B	358	TRP	2.1
1	A	359	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PEG	A	402	7/7	0.88	0.14	38,41,47,49	1
2	EDO	B	402	4/4	0.89	0.13	32,37,39,40	1
2	EDO	B	401	4/4	0.96	0.06	23,25,31,31	1
2	EDO	A	401	4/4	0.97	0.06	24,28,35,35	1

6.5 Other polymers [i](#)

There are no such residues in this entry.