



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:26 AM UTC

PDB ID : 3RO2 / pdb_00003ro2
Title : Structures of the LGN/NuMA complex
Authors : Shang, Y.; Wei, Z.
Deposited on : 2011-04-25
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	338	
2	B	28	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2792 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called G-protein-signaling modulator 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	328	Total	C	N	O	S	0	2	0
			2485	1557	444	477	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	GLY	-	expression tag	UNP Q8VDU0
A	14	SER	-	expression tag	UNP Q8VDU0

- Molecule 2 is a protein called peptide of Nuclear mitotic apparatus protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	22	Total	C	N	O	S	0	0	0
			186	116	29	39	2			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	96	Total O 96 96	0	0
4	B	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: G-protein-signaling modulator 2



• Molecule 2: peptide of Nuclear mitotic apparatus protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	91.31Å 91.31Å 178.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.14 – 2.30 36.14 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.1 (36.14-2.30) 97.2 (36.14-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.24 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.215 , 0.271 0.245 , 0.301	Depositor DCC
R_{free} test set	1012 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2792	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/2537	0.83	0/3432
2	B	0.67	0/190	0.80	0/257
All	All	0.51	0/2727	0.83	0/3689

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2485	0	2318	23	0
2	B	186	0	161	4	0
3	A	18	0	24	4	0
4	A	96	0	0	0	0
4	B	7	0	0	0	0
All	All	2792	0	2503	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:235:ARG:HD2	1:A:273:VAL:HG11	1.52	0.89
1:A:63:ASN:HB3	2:B:2004:LEU:HD21	1.71	0.72
1:A:83:THR:O	1:A:87:THR:HG23	2.05	0.57
1:A:131:LYS:HD2	1:A:184:LEU:HD11	1.86	0.56
1:A:37:VAL:HG23	1:A:64:ALA:HB1	1.88	0.56
1:A:291:TYR:O	1:A:295:ILE:HD12	2.06	0.55
1:A:238:TYR:HB2	1:A:261:THR:HG22	1.88	0.54
1:A:143:ASN:HB3	3:A:3:GOL:H32	1.92	0.51
1:A:257:TYR:O	1:A:261:THR:HG23	2.12	0.50
1:A:163:PHE:HB2	1:A:164:PRO:CD	2.44	0.48
1:A:40:PHE:HB3	1:A:61:LEU:HG	1.96	0.48
1:A:177:LEU:HD21	3:A:1:GOL:H32	1.96	0.47
1:A:21:ALA:HB1	2:B:2010:ILE:HD13	1.95	0.47
1:A:163:PHE:HB2	1:A:164:PRO:HD3	1.95	0.47
1:A:258:TYR:HB3	1:A:281:LEU:HG	1.98	0.44
1:A:303:ALA:HA	1:A:306:GLN:OE1	2.19	0.43
1:A:182:LEU:HD13	1:A:197:ALA:HB3	2.00	0.42
1:A:147:ALA:HB2	3:A:3:GOL:H12	2.00	0.42
1:A:61:LEU:HD13	1:A:77:TYR:CE1	2.55	0.41
1:A:238:TYR:CB	1:A:261:THR:HG22	2.50	0.41
1:A:63:ASN:ND2	3:A:2:GOL:O3	2.53	0.41
1:A:63:ASN:HB3	2:B:2004:LEU:CD2	2.45	0.40
1:A:66:PHE:CE1	2:B:2002:GLU:HG2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/338 (96%)	322 (99%)	3 (1%)	1 (0%)	36	46
2	B	20/28 (71%)	18 (90%)	1 (5%)	1 (5%)	1	1
All	All	346/366 (94%)	340 (98%)	4 (1%)	2 (1%)	21	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	1997	CYS
1	A	330	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/265 (88%)	226 (97%)	8 (3%)	32	49
2	B	20/26 (77%)	17 (85%)	3 (15%)	3	3
All	All	254/291 (87%)	243 (96%)	11 (4%)	25	39

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	VAL
1	A	55	SER
1	A	122	LEU
1	A	201	LEU
1	A	223	LEU
1	A	276	GLN
1	A	287	LEU
1	A	295	ILE
2	B	1996	THR
2	B	2006	ASP
2	B	2013	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	ASN
1	A	80	HIS
1	A	220	GLN
1	A	240	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	3	-	5,5,5	0.38	0	5,5,5	0.58	0
3	GOL	A	1	-	5,5,5	0.37	0	5,5,5	0.60	0
3	GOL	A	2	-	5,5,5	0.36	0	5,5,5	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	3	-	-	4/4/4/4	-
3	GOL	A	1	-	-	4/4/4/4	-
3	GOL	A	2	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1	GOL	O1-C1-C2-C3
3	A	1	GOL	C1-C2-C3-O3
3	A	3	GOL	O1-C1-C2-C3
3	A	3	GOL	C1-C2-C3-O3
3	A	1	GOL	O1-C1-C2-O2
3	A	3	GOL	O2-C2-C3-O3
3	A	2	GOL	C1-C2-C3-O3
3	A	1	GOL	O2-C2-C3-O3
3	A	2	GOL	O2-C2-C3-O3
3	A	3	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	3	GOL	2	0
3	A	1	GOL	1	0
3	A	2	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/338 (97%)	0.98	69 (21%) 2 3	17, 42, 97, 114	2 (0%)
2	B	22/28 (78%)	3.30	16 (72%) 0 0	60, 94, 109, 129	0
All	All	350/366 (95%)	1.12	85 (24%) 2 2	17, 45, 100, 129	2 (0%)

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	344	ILE	6.7
1	A	337	PHE	6.3
1	A	299	LEU	5.7
2	B	1996	THR	5.6
1	A	321	LEU	5.4
2	B	2007	TRP	5.3
2	B	2013	LEU	5.2
1	A	156	GLY	5.0
1	A	326	THR	5.0
2	B	1992	PHE	4.9
1	A	334	ALA	4.9
1	A	342	LEU	4.7
2	B	2010	ILE	4.5
1	A	320	SER	4.5
1	A	324	ALA	4.4
1	A	329	GLY	4.3
2	B	2011	ALA	4.3
2	B	1994	MET	4.3
1	A	319	TRP	4.2
1	A	338	ALA	4.2
1	A	13	GLY	4.1
1	A	331	HIS	4.1
2	B	2004	LEU	4.1
1	A	297	TYR	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	325	TYR	4.0
1	A	158	GLN	4.0
1	A	330	ASN	3.9
1	A	152	PHE	3.8
1	A	163	PHE	3.8
1	A	318	CYS	3.8
2	B	2005	ASP	3.7
1	A	335	MET	3.7
2	B	1997	CYS	3.7
1	A	322	GLY	3.7
1	A	301	HIS	3.6
1	A	327	ALA	3.5
2	B	2006	ASP	3.5
1	A	154	CYS	3.5
1	A	294	ALA	3.5
1	A	275	ALA	3.4
1	A	293	LYS	3.4
1	A	271	ARG	3.4
1	A	308	LEU	3.3
1	A	314	GLU	3.3
1	A	292	GLU	3.2
1	A	317	ALA	3.2
1	A	312	ILE	3.2
2	B	1993	TYR	3.1
1	A	295	ILE	3.1
2	B	2009	ARG	3.1
1	A	290	ASP	3.0
1	A	305	ALA	3.0
1	A	323	ASN	3.0
1	A	157	PRO	3.0
1	A	285	TYR	2.9
2	B	1995	GLY	2.9
1	A	332	ASP	2.9
1	A	300	LYS	2.8
2	B	1998	GLN	2.8
1	A	333	GLN	2.8
1	A	339	GLU	2.8
1	A	272	ALA	2.8
1	A	268	LEU	2.7
1	A	288	LEU	2.7
1	A	291	TYR	2.7
1	A	315	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	341	HIS	2.6
1	A	313	GLY	2.5
1	A	296	ASP	2.5
1	A	340	LYS	2.5
1	A	310	ASP	2.5
1	A	88	ILE	2.4
1	A	298	HIS	2.4
1	A	273	VAL	2.3
1	A	276	GLN	2.3
1	A	278	CYS	2.3
1	A	328	LEU	2.3
1	A	304	ILE	2.3
1	A	336	HIS	2.2
1	A	67	TYR	2.2
1	A	155	PRO	2.2
1	A	307	GLU	2.2
1	A	287	LEU	2.1
2	B	2012	GLU	2.1
1	A	235	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	3	6/6	0.91	0.15	49,52,57,58	0
3	GOL	A	1	6/6	0.93	0.09	41,44,48,53	0
3	GOL	A	2	6/6	0.94	0.12	36,37,41,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.