



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:30 PM UTC

PDB ID : 3ROO / pdb_00003roo
Title : Murine class I major histocompatibility complex H-2Kb in complex with immunodominant LCMV-derived gp34-41 peptide
Authors : Madhurantakam, C.; Duru, A.D.; Leong, C.; Sandalova, T.; Webb, J.R.; Achour, A.
Deposited on : 2011-04-26
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	17% 95% . .
1	C	275	19% 93% 6% .
2	B	99	7% 94% 6%
2	D	99	27% 90% 9% .
3	E	8	12% 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	8	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MPD	B	5276	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, K-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	4	0
			2258	1425	397	427	9			
1	C	271	Total	C	N	O	S	0	5	0
			2222	1404	388	420	10			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	3	0
			830	529	138	155	8			
2	D	99	Total	C	N	O	S	0	3	0
			824	526	135	155	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	85	ASP	ALA	variant	UNP P01887
D	85	ASP	ALA	variant	UNP P01887

- Molecule 3 is a protein called Pre-glycoprotein polypeptide GP complex.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	8	Total	C	N	O	S	0	0	0
			64	42	9	12	1			
3	F	8	Total	C	N	O	S	0	0	0
			64	42	9	12	1			

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			8	6	2		
5	C	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	P	0	0
			5	4	1		
6	C	1	Total	O	P	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	146	Total	O	0	0
			146	146		
7	B	74	Total	O	0	0
			74	74		
7	C	126	Total	O	0	0
			126	126		
7	D	34	Total	O	0	0
			34	34		
7	E	7	Total	O	0	0
			7	7		
7	F	7	Total	O	0	0
			7	7		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain



- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin



- Molecule 3: Pre-glycoprotein polyprotein GP complex

Chain E:  12%
100%



- Molecule 3: Pre-glycoprotein polypeptide GP complex

Chain F:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.45Å 92.59Å 128.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.26 – 2.00 57.26 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (57.26-2.00) 99.9 (57.26-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.225 , 0.267 0.230 , 0.271	Depositor DCC
R_{free} test set	3633 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6694	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MPD, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/2336	0.71	0/3171
1	C	0.54	0/2302	0.67	0/3121
2	B	0.55	0/870	0.72	0/1178
2	D	0.55	0/864	0.74	0/1171
3	E	0.55	0/65	0.68	0/86
3	F	0.75	0/65	0.77	0/86
All	All	0.55	0/6502	0.70	0/8813

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2146	9	0
1	C	2222	0	2113	9	0
2	B	830	0	800	7	0
2	D	824	0	789	9	0
3	E	64	0	61	0	0
3	F	64	0	61	0	0
4	A	6	0	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	6	0	8	0	0
5	B	8	0	14	6	0
5	C	8	0	14	5	0
6	B	5	0	0	0	0
6	C	5	0	0	0	0
7	A	146	0	0	1	0
7	B	74	0	0	0	0
7	C	126	0	0	0	0
7	D	34	0	0	0	0
7	E	7	0	0	0	0
7	F	7	0	0	0	0
All	All	6694	0	6014	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:5276:MPD:H11	2:D:33:PRO:O	1.83	0.79
2:D:12:ARG:CB	2:D:22:ILE:HG23	2.12	0.78
2:B:54[B]:MET:CE	5:B:5276:MPD:HM2	2.24	0.66
1:C:23[B]:MET:SD	2:D:54[B]:MET:SD	2.99	0.60
1:C:117:ALA:HB2	2:D:60:TRP:CE2	2.38	0.59
2:B:54[B]:MET:HE2	5:B:5276:MPD:H11	1.86	0.58
1:A:23:MET:HE3	5:B:5276:MPD:H53	1.85	0.58
1:C:94[B]:THR:HG21	2:D:33:PRO:CD	2.34	0.58
1:C:181:ARG:NH2	1:C:239:GLY:O	2.35	0.56
5:C:5276:MPD:O4	5:C:5276:MPD:H12	2.06	0.56
1:A:215:LEU:HD22	1:A:261:VAL:HG22	1.88	0.54
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.43	0.54
2:B:54[B]:MET:HE2	5:B:5276:MPD:HM2	1.89	0.54
2:B:27:VAL:HG21	2:B:37:ILE:HD13	1.89	0.53
5:C:5276:MPD:H4	2:D:33:PRO:HB2	1.93	0.49
2:D:1:ILE:O	2:D:1:ILE:HG23	2.13	0.48
2:B:54[B]:MET:HE1	5:B:5276:MPD:HM2	1.97	0.47
2:D:22:ILE:HD11	2:D:67:HIS:HB2	1.98	0.46
1:A:182:THR:HG21	1:A:265:GLY:HA2	1.97	0.46
1:C:216:THR:OG1	1:C:260:HIS:HB2	2.18	0.44
2:B:54[B]:MET:HE1	5:B:5276:MPD:H32	2.00	0.44
1:C:251:LEU:HD12	1:C:252:GLY:H	1.82	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:21:ASN:OD1	2:D:22:ILE:N	2.45	0.44
1:A:182:THR:HG23	1:A:265:GLY:HA3	2.00	0.43
1:C:23[A]:MET:SD	5:C:5276:MPD:H53	2.59	0.43
1:C:251:LEU:HD12	1:C:252:GLY:N	2.35	0.42
1:A:228:MET:HG2	1:A:247:VAL:HG12	2.02	0.42
1:A:215:LEU:CD2	1:A:261:VAL:HG22	2.50	0.41
1:A:54:GLN:NE2	7:A:289:HOH:O	2.53	0.41
1:C:203:CYS:HB2	1:C:217:TRP:CZ2	2.56	0.41
5:C:5276:MPD:O4	5:C:5276:MPD:C1	2.69	0.40
1:A:141:LEU:HD12	1:A:141:LEU:HA	2.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/275 (101%)	268 (97%)	9 (3%)	0	100	100
1	C	270/275 (98%)	263 (97%)	7 (3%)	0	100	100
2	B	100/99 (101%)	98 (98%)	2 (2%)	0	100	100
2	D	100/99 (101%)	100 (100%)	0	0	100	100
3	E	6/8 (75%)	6 (100%)	0	0	100	100
3	F	6/8 (75%)	6 (100%)	0	0	100	100
All	All	759/764 (99%)	741 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/233 (102%)	230 (97%)	7 (3%)	36	38
1	C	234/233 (100%)	229 (98%)	5 (2%)	47	52
2	B	97/94 (103%)	95 (98%)	2 (2%)	47	52
2	D	96/94 (102%)	93 (97%)	3 (3%)	35	37
3	E	6/6 (100%)	6 (100%)	0	100	100
3	F	6/6 (100%)	6 (100%)	0	100	100
All	All	676/666 (102%)	659 (98%)	17 (2%)	40	45

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	TYR
1	A	114	GLN
1	A	141	LEU
1	A	182	THR
1	A	196	GLU
1	A	199	VAL
1	A	228	MET
2	B	56	PHE
2	B	58	LYS
1	C	111	ARG
1	C	114	GLN
1	C	141	LEU
1	C	230	LEU
1	C	255	GLN
2	D	22	ILE
2	D	56	PHE
2	D	58	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	115	GLN
1	A	174	ASN
1	A	218	GLN
2	B	38	GLN
1	C	86	ASN
1	C	114	GLN
1	C	115	GLN
1	C	174	ASN
2	D	29	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PO4	C	276	-	4,4,4	1.02	0	6,6,6	0.45	0
4	GOL	D	3968	-	5,5,5	0.44	0	5,5,5	0.40	0
5	MPD	C	5276	-	7,7,7	0.46	0	9,10,10	0.62	0
5	MPD	B	5276	-	7,7,7	0.24	0	9,10,10	0.69	0
4	GOL	A	3968	-	5,5,5	0.33	0	5,5,5	0.57	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PO4	B	100	-	4,4,4	0.94	0	6,6,6	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	3968	-	-	2/4/4/4	-
4	GOL	A	3968	-	-	4/4/4/4	-
5	MPD	B	5276	-	-	4/5/5/5	-
5	MPD	C	5276	-	-	4/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	3968	GOL	O1-C1-C2-C3
4	D	3968	GOL	C1-C2-C3-O3
5	C	5276	MPD	C1-C2-C3-C4
5	C	5276	MPD	O2-C2-C3-C4
4	A	3968	GOL	C1-C2-C3-O3
4	D	3968	GOL	O2-C2-C3-O3
4	A	3968	GOL	O2-C2-C3-O3
4	A	3968	GOL	O1-C1-C2-O2
5	B	5276	MPD	C1-C2-C3-C4
5	B	5276	MPD	CM-C2-C3-C4
5	B	5276	MPD	O2-C2-C3-C4
5	B	5276	MPD	C2-C3-C4-O4
5	C	5276	MPD	C2-C3-C4-O4
5	C	5276	MPD	CM-C2-C3-C4

There are no ring outliers.

3 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	5276	MPD	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	5276	MPD	6	0
4	A	3968	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	0.95	47 (17%) 4 4	20, 37, 64, 69	5 (1%)
1	C	271/275 (98%)	1.03	51 (18%) 3 3	18, 40, 67, 75	6 (2%)
2	B	99/99 (100%)	0.48	7 (7%) 22 20	22, 34, 48, 54	3 (3%)
2	D	99/99 (100%)	1.31	27 (27%) 1 1	22, 43, 59, 62	3 (3%)
3	E	8/8 (100%)	0.15	1 (12%) 8 7	25, 28, 33, 33	0
3	F	8/8 (100%)	-0.01	0 100 100	27, 29, 32, 33	0
All	All	760/764 (99%)	0.95	133 (17%) 4 3	18, 38, 64, 75	17 (2%)

All (133) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	219	LEU	5.9
1	C	256	TYR	5.6
1	A	199	VAL	5.5
1	A	256	TYR	5.4
2	D	22	ILE	5.1
1	C	224	LEU	4.8
1	A	272	LEU	4.7
1	C	270	LEU	4.5
2	B	1	ILE	4.5
1	C	266	LEU	4.5
1	A	195	PRO	4.4
1	A	219	LEU	4.4
1	C	222	GLU	4.3
1	C	257	TYR	4.1
1	C	272	LEU	4.0
1	C	258	THR	3.9
1	C	199	VAL	3.9
1	A	251	LEU	3.9
1	A	249	VAL	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	88	ALA	3.8
1	A	198	LYS	3.8
1	A	226	GLN	3.8
2	D	14	PRO	3.8
1	A	274	TRP	3.7
1	C	274	TRP	3.7
1	A	259	CYS	3.7
1	C	195	PRO	3.7
1	C	228	MET	3.6
1	A	269	PRO	3.6
2	D	15	PRO	3.6
1	C	221	GLY	3.6
1	C	255	GLN	3.5
1	A	225	ILE	3.5
2	D	20	PRO	3.5
1	A	220	ASN	3.4
1	C	267	PRO	3.4
1	A	260	HIS	3.3
2	B	48	LYS	3.3
1	C	190	THR	3.2
1	A	257	TYR	3.2
1	A	262	TYR	3.2
1	A	250	PRO	3.2
2	D	58	LYS	3.1
1	C	273	ARG	3.1
1	C	251	LEU	3.1
2	D	1	ILE	3.1
2	D	16	GLU	3.0
2	D	19	LYS	3.0
2	D	12	ARG	3.0
2	D	71	THR	3.0
1	C	250	PRO	3.0
1	A	222	GLU	3.0
1	C	186	LYS	3.0
1	C	269	PRO	3.0
1	A	191	HIS	3.0
1	C	252	GLY	2.9
1	A	227	ASP	2.9
1	A	196	GLU	2.9
1	C	237	GLY	2.9
2	D	67	HIS	2.9
1	A	258	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	266	LEU	2.8
2	D	49	VAL	2.8
1	A	58	GLU	2.8
1	A	270	LEU	2.8
1	A	224	LEU	2.7
1	A	217	TRP	2.7
1	C	58	GLU	2.7
2	D	18	GLY	2.7
1	A	273	ARG	2.7
1	C	249	VAL	2.7
1	A	221	GLY	2.7
1	A	268	GLU	2.6
1	C	189	VAL	2.6
1	C	259	CYS	2.6
1	C	265	GLY	2.6
1	A	247	VAL	2.6
1	C	248	VAL	2.6
1	C	268	GLU	2.6
2	D	13	HIS	2.6
1	C	217	TRP	2.6
1	A	41	GLU	2.6
1	C	201	LEU	2.6
1	A	228	MET	2.5
2	B	58	LYS	2.5
1	A	193	SER	2.5
1	C	196	GLU	2.5
1	C	107	GLY	2.5
1	C	218	GLN	2.4
1	C	18	GLY	2.4
1	A	197	ASP	2.4
1	A	271	THR	2.4
2	D	17	ASN	2.4
2	D	97	ARG	2.4
1	C	1	GLY	2.4
1	C	42	ASN	2.4
2	D	21	ASN	2.4
2	B	85	ASP	2.4
1	C	260	HIS	2.4
1	C	113	TYR	2.3
1	C	188	HIS	2.3
1	A	254	GLU	2.3
1	A	200	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	E	4	ASN	2.3
1	C	231	VAL	2.3
1	C	271	THR	2.3
1	A	170	ARG	2.3
1	C	105	SER	2.2
1	C	253	LYS	2.2
1	C	203	CYS	2.2
1	A	255	GLN	2.2
1	A	253	LYS	2.2
2	D	94	TYR	2.2
2	D	11	SER	2.2
1	A	108	ARG	2.2
2	D	45	LYS	2.2
1	C	128	GLU	2.1
2	D	7	ILE	2.1
1	C	262	TYR	2.1
1	A	188	HIS	2.1
1	C	108	ARG	2.1
2	D	3	LYS	2.1
2	B	4	THR	2.1
1	A	261	VAL	2.1
2	D	70	PHE	2.1
1	A	275	GLU	2.0
1	C	138	MET	2.0
2	D	69	GLU	2.0
2	D	98	ASP	2.0
2	D	8	GLN	2.0
2	D	48	LYS	2.0
1	A	267	PRO	2.0
2	B	90	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	PO4	C	276	5/5	0.84	0.14	65,66,66,66	0
4	GOL	D	3968	6/6	0.89	0.17	58,60,60,61	0
6	PO4	B	100	5/5	0.89	0.12	71,72,73,73	0
4	GOL	A	3968	6/6	0.89	0.15	29,38,40,40	0
5	MPD	C	5276	8/8	0.90	0.13	34,37,39,41	0
5	MPD	B	5276	8/8	0.94	0.11	33,35,38,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.