



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:42 AM UTC

PDB ID : 7RP0 / pdb_00007rp0
Title : Structural Snapshots of Intermediates in the Gating of a K⁺ Channel
Authors : Reddi, R.; Matulef, K.; Riederer, E.A.; Valiyaveetil, F.I.
Deposited on : 2021-08-02
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	27% 90% 10%
2	B	212	23% 92% 8%
3	C	103	6% 95% 5%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4115 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called KcsA Fab chain A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1610	1023	269	312	6			

- Molecule 2 is a protein called KcsA Fab chain B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	212	Total	C	N	O	S	0	0	0
			1646	1022	282	337	5			


- Molecule 3 is a protein called pH-gated potassium channel KcsA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	103	Total	C	N	O	S	0	0	0
			760	498	130	131	1			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	82	ALA	TYR	engineered mutation	UNP P0A334

- Molecule 4 is NONAN-1-OL (CCD ID: F09) (formula: C₉H₂₀O).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	9	1		

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	4	Total	K	0	0
			4	4		

- Molecule 6 is TETRABUTYLAMMONIUM ION (CCD ID: TBA) (formula: C₁₆H₃₆N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	N	0	0
			17	16	1		

- Molecule 7 is DIACYL GLYCEROL (CCD ID: DGA) (formula: $C_{39}H_{76}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			31	26	5		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	8	Total	O	0	0
			8	8		
8	B	22	Total	O	0	0
			22	22		
8	C	7	Total	O	0	0
			7	7		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: KcsA Fab chain A



• Molecule 2: KcsA Fab chain B



• Molecule 3: pH-gated potassium channel KcsA



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	155.33Å 155.33Å 76.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.92 – 2.48 54.92 – 2.48	Depositor EDS
% Data completeness (in resolution range)	90.0 (54.92-2.48) 89.7 (54.92-2.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 2.48Å)	Xtriage
Refinement program	PHENIX 1.14 _3260	Depositor
R, R_{free}	0.241 , 0.271 0.242 , 0.271	Depositor DCC
R_{free} test set	3262 reflections (5.56%)	wwPDB-VP
Wilson B-factor (Å ²)	37.4	Xtriage
Anisotropy	0.723	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.038 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4115	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: F09, TBA, K, DGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.09	0/1654	0.27	0/2267
2	B	0.09	0/1683	0.27	0/2283
3	C	0.09	0/778	0.26	0/1069
All	All	0.09	0/4115	0.27	0/5619

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1610	0	1545	11	0
2	B	1646	0	1572	10	0
3	C	760	0	772	5	0
4	A	10	0	20	1	0
5	C	4	0	0	0	0
6	C	17	0	36	0	0
7	C	31	0	44	2	0
8	A	8	0	0	0	0
8	B	22	0	0	0	0
8	C	7	0	0	0	0
All	All	4115	0	3989	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:37:GLN:HB2	2:B:47:LEU:HD11	1.71	0.72
1:A:83:LEU:HB3	1:A:86:LEU:HD21	1.81	0.62
4:A:301:F09:H62	3:C:46:LEU:HD22	1.86	0.58
1:A:40:ARG:HG2	1:A:92:ALA:HB2	1.88	0.56
2:B:13:VAL:HG21	2:B:19:VAL:HB	1.88	0.54
3:C:86:LEU:HB2	7:C:506:DGA:HG11	1.90	0.53
1:A:124:PRO:HB3	1:A:150:TYR:HB3	1.92	0.52
3:C:51:GLU:HG3	3:C:59:LEU:HB3	1.92	0.51
1:A:38:LYS:HB2	1:A:48:ILE:HD11	1.94	0.49
3:C:86:LEU:HD22	7:C:506:DGA:HB21	1.94	0.49
1:A:172:PRO:HD3	2:B:164:THR:HG22	1.97	0.46
2:B:150:ILE:HD11	2:B:179:LEU:HD21	1.97	0.46
1:A:12:VAL:HG11	1:A:86:LEU:HD12	1.99	0.44
1:A:193:TRP:CD1	1:A:194:PRO:HA	2.53	0.44
2:B:39:ARG:HG2	2:B:84:ALA:HB2	2.00	0.43
1:A:24:ALA:HB1	1:A:98:ARG:HH21	1.83	0.43
1:A:31:SER:HB2	3:C:62:TYR:CE1	2.54	0.43
2:B:147:LYS:HB3	2:B:195:GLU:HB2	2.01	0.43
2:B:149:LYS:HB2	2:B:193:THR:HB	2.01	0.41
2:B:195:GLU:HG2	2:B:206:VAL:HG22	2.03	0.41
2:B:35:TRP:CE2	2:B:73:LEU:HB2	2.56	0.41
1:A:148:LYS:HE2	1:A:148:LYS:HB2	1.92	0.40
2:B:13:VAL:O	2:B:106:ILE:HA	2.22	0.40
1:A:199:THR:HG22	1:A:214:LYS:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/219 (99%)	207 (95%)	10 (5%)	0	100	100
2	B	210/212 (99%)	201 (96%)	9 (4%)	0	100	100
3	C	101/103 (98%)	100 (99%)	1 (1%)	0	100	100
All	All	528/534 (99%)	508 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/185 (93%)	168 (98%)	4 (2%)	44	69
2	B	189/190 (100%)	188 (100%)	1 (0%)	81	91
3	C	71/74 (96%)	71 (100%)	0	100	100
All	All	432/449 (96%)	427 (99%)	5 (1%)	63	81

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	156	THR
1	A	157	VAL
1	A	158	THR
1	A	174	VAL
2	B	19	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
2	B	38	GLN
2	B	124	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	DGA	C	506	-	30,30,43	1.43	3 (10%)	32,32,45	1.26	2 (6%)
6	TBA	C	505	-	16,16,16	0.96	0	18,18,18	0.60	0
4	F09	A	301	-	9,9,9	0.32	0	8,8,8	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	DGA	C	506	-	-	18/32/32/45	-
6	TBA	C	505	-	-	6/20/20/20	-
4	F09	A	301	-	-	5/7/7/7	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	506	DGA	OG1-CA1	3.62	1.43	1.33
7	C	506	DGA	OG2-CB1	2.98	1.42	1.34
7	C	506	DGA	OG2-CG2	-2.89	1.39	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	506	DGA	OG2-CB1-CB2	4.05	120.25	111.48
7	C	506	DGA	OG1-CA1-CA2	2.67	119.98	111.83

There are no chirality outliers.

All (29) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	C	506	DGA	OB1-CB1-OG2-CG2
7	C	506	DGA	OG1-CG1-CG2-OG2
7	C	506	DGA	OG2-CG2-CG3-OXT
7	C	506	DGA	CB2-CB1-OG2-CG2
7	C	506	DGA	CB6-CB7-CB8-CB9
6	C	505	TBA	N1-C11-C12-C13
4	A	301	F09	C3-C4-C5-C6
7	C	506	DGA	CBB-CAB-CB9-CB8
7	C	506	DGA	CA4-CA5-CA6-CA7
7	C	506	DGA	CB9-CAB-CBB-CCB
7	C	506	DGA	CA2-CA1-OG1-CG1
7	C	506	DGA	CB7-CB8-CB9-CAB
4	A	301	F09	C2-C3-C4-C5
6	C	505	TBA	C12-C11-N1-C41
6	C	505	TBA	C12-C11-N1-C21
7	C	506	DGA	OA1-CA1-OG1-CG1
4	A	301	F09	C6-C7-C8-C9
7	C	506	DGA	CB3-CB4-CB5-CB6
7	C	506	DGA	CA6-CA7-CA8-CA9
7	C	506	DGA	CG1-CG2-CG3-OXT
7	C	506	DGA	CB4-CB5-CB6-CB7
6	C	505	TBA	C12-C11-N1-C31
4	A	301	F09	C1-C2-C3-C4
7	C	506	DGA	OG1-CG1-CG2-CG3
6	C	505	TBA	C21-C22-C23-C24
6	C	505	TBA	C41-C42-C43-C44
4	A	301	F09	C5-C6-C7-C8
7	C	506	DGA	CA3-CA4-CA5-CA6
7	C	506	DGA	CB5-CB6-CB7-CB8

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	506	DGA	2	0
4	A	301	F09	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/219 (100%)	1.52	60 (27%)  	37, 63, 79, 85	0
2	B	212/212 (100%)	1.23	48 (22%)  	31, 58, 83, 93	1 (0%)
3	C	103/103 (100%)	0.46	6 (5%)  	28, 35, 52, 58	0
All	All	534/534 (100%)	1.20	114 (21%)  	28, 57, 79, 93	1 (0%)

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	GLN	5.0
2	B	210	ASN	4.5
1	A	2	VAL	4.4
1	A	219	ASP	4.4
1	A	134	ALA	4.4
1	A	136	GLN	4.0
1	A	131	PRO	4.0
1	A	24	ALA	3.9
1	A	178	ASP	3.9
2	B	208	SER	3.9
2	B	179	LEU	3.8
1	A	120	LYS	3.8
1	A	132	GLY	3.8
3	C	22	SER	3.8
2	B	209	PHE	3.7
1	A	199	THR	3.7
1	A	137	THR	3.6
1	A	84	SER	3.6
1	A	164	LEU	3.5
1	A	166	SER	3.5
1	A	138	ASN	3.4
1	A	196	GLU	3.4
2	B	124	GLN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	212	ASN	3.3
1	A	82	GLN	3.3
1	A	119	ALA	3.2
1	A	159	TRP	3.2
1	A	5	GLN	3.2
1	A	198	VAL	3.2
2	B	133	VAL	3.1
2	B	119	PRO	3.1
2	B	163	TRP	3.1
2	B	156	GLN	3.1
2	B	118	PHE	3.0
1	A	85	SER	3.0
2	B	132	VAL	2.9
2	B	125	LEU	2.9
2	B	117	ILE	2.9
1	A	55	TYR	2.9
1	A	177	SER	2.9
1	A	211	VAL	2.9
2	B	148	TRP	2.8
1	A	169	HIS	2.8
2	B	121	SER	2.8
1	A	142	THR	2.8
1	A	13	LYS	2.8
1	A	3	GLN	2.8
2	B	191	SER	2.7
2	B	116	SER	2.7
1	A	129	LEU	2.7
2	B	130	ALA	2.7
1	A	65	GLN	2.7
2	B	64	GLY	2.6
2	B	205	ILE	2.6
2	B	182	THR	2.6
2	B	190	ASN	2.6
1	A	135	ALA	2.6
2	B	192	TYR	2.6
3	C	123	GLY	2.6
1	A	145	CYS	2.6
2	B	126	THR	2.5
3	C	23	ALA	2.5
1	A	22	CYS	2.5
2	B	128	GLY	2.5
1	A	193	TRP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	12	VAL	2.5
2	B	180	THR	2.5
2	B	159	VAL	2.4
1	A	210	LYS	2.4
1	A	214	LYS	2.4
2	B	183	LYS	2.4
1	A	212	ASP	2.4
2	B	73	LEU	2.4
1	A	157	VAL	2.4
2	B	122	SER	2.4
1	A	11	LEU	2.4
1	A	152	PRO	2.4
1	A	195	SER	2.4
2	B	152	GLY	2.4
2	B	115	VAL	2.4
1	A	217	PRO	2.3
2	B	12	SER	2.3
2	B	181	LEU	2.3
3	C	35	LEU	2.3
2	B	120	PRO	2.3
2	B	188	ARG	2.3
3	C	122	ARG	2.3
1	A	206	ALA	2.3
1	A	187	THR	2.3
3	C	124	HIS	2.3
1	A	175	LEU	2.3
2	B	76	ASN	2.2
1	A	42	GLY	2.2
1	A	80	PHE	2.2
2	B	135	PHE	2.2
1	A	27	TYR	2.2
1	A	122	THR	2.2
1	A	179	LEU	2.2
2	B	105	GLU	2.2
2	B	41	ASN	2.2
1	A	176	GLN	2.2
1	A	141	VAL	2.1
2	B	189	HIS	2.1
1	A	67	LYS	2.1
2	B	17	GLU	2.1
1	A	127	TYR	2.1
1	A	197	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	127	SER	2.1
1	A	143	LEU	2.0
2	B	129	GLY	2.0
2	B	202	THR	2.0
2	B	206	VAL	2.0
1	A	83	LEU	2.0
2	B	153	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	DGA	C	506	31/44	0.82	0.20	36,54,65,69	0
4	F09	A	301	10/10	0.87	0.19	42,52,55,60	0
5	K	C	502	1/1	0.93	0.06	36,36,36,36	1
6	TBA	C	505	17/17	0.95	0.11	29,30,31,31	17
5	K	C	503	1/1	0.95	0.08	35,35,35,35	1
5	K	C	504	1/1	0.96	0.08	29,29,29,29	1
5	K	C	501	1/1	0.98	0.03	32,32,32,32	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around DGA C 506:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around K C 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K C 503:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around K C 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.