



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:27 AM UTC

PDB ID : 1RQI / pdb_00001rqi
Title : Active Conformation of Farnesyl Pyrophosphate Synthase Bound to Isopentyl Pyrophosphate and Dimethylallyl S-Thiolodiphosphate
Authors : Hosfield, D.J.; Zhang, Y.; Dougan, D.R.; Brooun, A.; Tari, L.W.; Swanson, R.V.; Finn, J.
Deposited on : 2003-12-05
Resolution : 2.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6562 (2.44-2.40)
Ramachandran outliers	187476	6481 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	300	 81% 17% •
1	B	300	 84% 14% ••

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4852 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Geranyltranstransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	300	Total	C	N	O	S	0	7	0
			2279	1418	408	440	13			
1	B	298	Total	C	N	O	S	0	11	0
			2265	1411	405	437	12			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	ALA	-	cloning artifact	UNP P22939
B	21	ALA	-	cloning artifact	UNP P22939

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Mg	0	0
			3	3		
2	B	3	Total	Mg	0	0
			3	3		

- Molecule 3 is DIMETHYLALLYL S-THIOLODIPHOSPHATE (CCD ID: DST) (formula: C₅H₁₂O₆P₂S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	O	P	S	0	0
			14	5	6	2	1		
3	B	1	Total	C	O	P	S	0	0
			14	5	6	2	1		

- Molecule 4 is ISOPENTYL PYROPHOSPHATE (CCD ID: IPR) (formula: C₅H₁₄O₇P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	O	P		0	0
			14	5	7	2			
4	B	1	Total	C	O	P		0	0
			14	5	7	2			

- Molecule 5 is DIPHOSPHATE (CCD ID: DPO) (formula: O_7P_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			9	7	2		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	107	Total	O	0	0
			107	107		
6	B	130	Total	O	0	0
			130	130		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Geranyltranstransferase



- Molecule 1: Geranyltranstransferase



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	88.84Å 88.84Å 174.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.03 – 2.42	Depositor
% Data completeness (in resolution range)	100.0 (43.03-2.42)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.203 , 0.259	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4852	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DST, MG, IPR, DPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	6/2332 (0.3%)	1.59	14/3151 (0.4%)
1	B	0.69	0/2318	0.92	0/3133
All	All	0.94	6/4650 (0.1%)	1.30	14/6284 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	41	ARG	C-N	30.39	1.80	1.33
1	A	150	ASP	C-N	-20.03	1.09	1.33
1	A	21	ALA	C-N	18.35	1.63	1.33
1	A	40	SER	C-N	-12.78	1.14	1.33
1	A	149	ALA	C-N	8.87	1.45	1.33
1	A	25	PRO	CB-CG	8.73	1.93	1.49

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	21	ALA	CA-C-N	-36.09	65.57	122.23
1	A	21	ALA	C-N-CA	-36.09	65.57	122.23
1	A	21	ALA	O-C-N	-30.80	73.72	123.00
1	A	150	ASP	O-C-N	-15.63	104.56	123.01
1	A	25	PRO	N-CA-CB	15.53	120.96	103.26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	40	SER	CA-C-N	14.41	148.04	121.52
1	A	40	SER	C-N-CA	14.41	148.04	121.52
1	A	40	SER	O-C-N	-13.78	106.10	122.22
1	A	41	ARG	O-C-N	11.82	139.14	122.36
1	A	150	ASP	CA-C-N	10.83	132.44	122.37
1	A	150	ASP	C-N-CA	10.83	132.44	122.37
1	A	25	PRO	CA-CB-CG	-8.97	87.45	104.50
1	A	149	ALA	CA-C-N	-6.14	112.64	121.42
1	A	149	ALA	C-N-CA	-6.14	112.64	121.42

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	21	ALA	Mainchain,Peptide
1	A	41	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2279	0	2281	38	0
1	B	2265	0	2261	28	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	14	0	8	1	0
3	B	14	0	8	2	0
4	A	14	0	10	1	0
4	B	14	0	11	2	0
5	A	9	0	0	0	0
6	A	107	0	0	1	0
6	B	130	0	0	5	1
All	All	4852	0	4579	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:PRO:CB	1:A:25:PRO:CG	1.93	1.41
1:A:41:ARG:C	1:A:42:PHE:N	1.80	1.39
1:A:41:ARG:HH21	1:A:41:ARG:HB3	1.37	0.86
1:A:37:GLN:O	1:A:41:ARG:HD3	1.80	0.82
1:A:41:ARG:HB3	1:A:41:ARG:NH2	1.97	0.79
1:A:41:ARG:HH21	1:A:41:ARG:CB	1.96	0.78
1:B:100:TYR:CE2	1:B:139:GLN:HG3	2.23	0.73
1:B:175[B]:MET:CE	1:B:206:LEU:HD22	2.18	0.73
1:B:29:GLU:HG2	6:B:908:HOH:O	1.90	0.71
1:B:223:ARG:NE	6:B:940:HOH:O	2.27	0.68
1:A:41:ARG:HH21	1:A:41:ARG:CG	2.08	0.67
1:A:41:ARG:CA	1:A:42:PHE:N	2.57	0.65
1:B:89:ALA:HB3	1:B:90:PRO:CD	2.27	0.65
1:A:41:ARG:NH2	1:A:41:ARG:CB	2.59	0.64
1:B:97:ILE:HG22	1:B:175[B]:MET:HE1	1.80	0.63
1:A:97:ILE:HG22	1:A:175[B]:MET:HE1	1.82	0.60
1:A:169:ALA:HB2	1:A:208[B]:ARG:NH2	2.16	0.60
1:B:175[B]:MET:HE2	1:B:206:LEU:HD22	1.82	0.60
1:A:150:ASP:OD2	1:A:152:PRO:HG3	2.01	0.60
1:B:89:ALA:HB3	1:B:90:PRO:HD3	1.81	0.59
1:A:46:LEU:HD13	1:B:172:ILE:HG21	1.84	0.59
1:B:33:LYS:NZ	6:B:910:HOH:O	2.34	0.58
1:A:317:GLN:NE2	6:A:861:HOH:O	2.36	0.58
1:A:35:ALA:O	1:A:39:LEU:HD12	2.04	0.57
1:A:69:ARG:NH2	1:A:98:HIS:CE1	2.73	0.56
1:B:100:TYR:CD2	1:B:139:GLN:HG3	2.40	0.56
1:A:41:ARG:N	1:A:42:PHE:N	2.54	0.55
1:B:175[B]:MET:HE1	1:B:206:LEU:HD22	1.87	0.53
1:B:79:MET:HE1	1:B:229:LEU:HD21	1.90	0.53
1:A:169:ALA:HB2	1:A:208[B]:ARG:HH21	1.74	0.51
1:A:97:ILE:CG2	1:A:175[B]:MET:HE1	2.41	0.50
1:A:69:ARG:CZ	1:A:98:HIS:CE1	2.94	0.50
1:B:199:HIS:CD2	1:B:241:GLN:HG3	2.47	0.49
1:A:89:ALA:HB3	1:A:90:PRO:CD	2.42	0.49
1:A:41:ARG:C	1:A:42:PHE:CA	2.81	0.49
3:B:400:DST:C10	4:B:500:IPR:HC13	2.43	0.48
1:A:101:SER:HB2	3:A:401:DST:H131	1.95	0.47
1:B:157[A]:ARG:HG3	6:B:936:HOH:O	2.13	0.47
1:A:66:LYS:NZ	4:A:501:IPR:O13	2.41	0.46
1:A:53:VAL:HG21	1:B:180:ALA:CB	2.46	0.46
1:A:104:HIS:O	1:A:110:MET:HG3	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:TYR:OH	1:B:136:ASP:HB3	2.17	0.45
1:B:292[A]:ARG:CZ	6:B:869:HOH:O	2.64	0.45
1:B:97:ILE:HG13	1:B:166:LEU:HD11	1.99	0.45
3:B:400:DST:H101	4:B:500:IPR:HC13	1.99	0.45
1:B:208:ARG:HA	1:B:233:ALA:HB1	1.99	0.44
1:B:77:GLY:O	1:B:82[B]:VAL:HG13	2.18	0.44
1:A:53:VAL:HG21	1:B:180:ALA:HB2	2.00	0.43
1:A:68:LEU:HD22	1:A:311:LEU:HD11	2.00	0.43
1:A:181:LEU:HD11	1:B:48:PHE:CZ	2.54	0.43
1:B:82[C]:VAL:HG21	1:B:217[C]:SER:OG	2.19	0.43
1:B:77:GLY:O	1:B:82[C]:VAL:HG22	2.19	0.42
1:A:245:ASP:HB3	1:A:272:PRO:HD2	2.02	0.42
1:B:82[C]:VAL:HG23	1:B:87:LEU:HD11	2.01	0.42
1:B:100:TYR:CD1	1:B:100:TYR:C	2.98	0.42
1:A:44:ALA:N	1:A:45:PRO:CD	2.84	0.41
1:A:232:TYR:CE1	1:A:295:LEU:HD11	2.55	0.41
1:A:172:ILE:HG21	1:B:46:LEU:HD13	2.02	0.41
1:A:72:LEU:CD1	1:A:207:ILE:HG23	2.50	0.41
1:A:144:SER:O	1:A:148:ASP:HB2	2.21	0.41
1:A:41:ARG:HH21	1:A:41:ARG:HG2	1.82	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:939:HOH:O	6:B:941:HOH:O[6_565]	1.64	0.56

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	303/300 (101%)	300 (99%)	3 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	301/300 (100%)	295 (98%)	5 (2%)	1 (0%)	36	49
All	All	604/600 (101%)	595 (98%)	8 (1%)	1 (0%)	43	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	300	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/234 (102%)	221 (92%)	18 (8%)	12	20
1	B	238/234 (102%)	215 (90%)	23 (10%)	8	11
All	All	477/468 (102%)	436 (91%)	41 (9%)	10	15

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	MET
1	A	26	GLN
1	A	39	LEU
1	A	41	ARG
1	A	63	LEU
1	A	100	TYR
1	A	115	LEU
1	A	157	ARG
1	A	183	LEU
1	A	192	LEU
1	A	197	ARG
1	A	213	LEU
1	A	231	LYS
1	A	242	VAL
1	A	264	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	282	LYS
1	A	289	ASP
1	A	292	ARG
1	B	27	GLN
1	B	28	LEU
1	B	37	GLN
1	B	63	LEU
1	B	82[B]	VAL
1	B	82[C]	VAL
1	B	100	TYR
1	B	115	LEU
1	B	125	LYS
1	B	139	GLN
1	B	183	LEU
1	B	192	LEU
1	B	207	ILE
1	B	213	LEU
1	B	231[A]	LYS
1	B	242	VAL
1	B	248	ASP
1	B	252	ASP
1	B	259	ARG
1	B	278	GLU
1	B	292[A]	ARG
1	B	296	LYS
1	B	302	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	130	ASN
1	A	139	GLN
1	A	241	GLN
1	A	260	GLN
1	A	279	GLN
1	A	293	GLN
1	A	317	GLN
1	B	34	GLN
1	B	37	GLN
1	B	58	GLN
1	B	139	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	279	GLN
1	B	317	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	IPR	A	501	-	12,13,13	0.75	0	16,19,19	1.07	1 (6%)
3	DST	A	401	2	10,13,13	3.54	1 (10%)	12,19,19	1.82	3 (25%)
5	DPO	A	606	-	6,8,8	0.77	0	12,13,13	0.98	0
3	DST	B	400	2	10,13,13	4.06	2 (20%)	12,19,19	2.15	4 (33%)
4	IPR	B	500	-	12,13,13	0.57	0	16,19,19	1.01	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IPR	A	501	-	-	3/13/13/13	-
3	DST	A	401	2	-	0/7/13/13	-
5	DPO	A	606	-	-	0/6/6/6	-
3	DST	B	400	2	-	0/7/13/13	-
4	IPR	B	500	-	-	4/13/13/13	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	400	DST	C10-S9	-12.26	1.69	1.84
3	A	401	DST	C10-S9	-10.79	1.70	1.84
3	B	400	DST	P3-O8	-2.37	1.50	1.56

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	400	DST	C11-C10-S9	4.44	133.55	111.87
3	B	400	DST	O8-P3-O7	4.03	119.47	109.86
3	A	401	DST	O8-P3-O7	3.75	118.80	109.86
3	A	401	DST	C11-C10-S9	3.43	128.59	111.87
4	A	501	IPR	C3-C2-C1	2.93	123.61	110.53
3	B	400	DST	C14-C12-C13	2.72	120.85	114.59
4	B	500	IPR	C3-C2-C1	2.45	121.44	110.53
3	A	401	DST	C14-C12-C13	2.23	119.73	114.59
3	B	400	DST	O6-P1-O4	2.06	115.52	107.80

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	501	IPR	C5-O6-P7-O9
4	B	500	IPR	C5-O6-P7-O9
4	B	500	IPR	C2-C4-C5-O6
4	A	501	IPR	C1-C2-C4-C5
4	B	500	IPR	C1-C2-C4-C5
4	A	501	IPR	C2-C4-C5-O6
4	B	500	IPR	P11-O10-P7-O9


There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	501	IPR	1	0
3	A	401	DST	1	0
3	B	400	DST	2	0
4	B	500	IPR	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	4

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	41:ARG	C	42:PHE	N	1.80
1	A	21:ALA	C	22:MET	N	1.63
1	A	40:SER	C	41:ARG	N	1.14
1	A	150:ASP	C	151:MET	N	1.09

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.