



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:12 PM UTC

PDB ID : 3RQL / pdb_00003rql
Title : Structure of the neuronal nitric oxide synthase heme domain in complex with 6-(((3R,4R)-4-(2-((1S,2R/1R,2S)-2-(3-Clorophenyl)cyclopropylamino)ethoxy)pyrrolidin-3-yl)methyl)-4-methylpyridin-2-amine
Authors : Li, H.; Delker, S.L.; Poulos, T.L.
Deposited on : 2011-04-28
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	4% 83% 13% ..
1	B	422	2% 84% 12% ..

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7193 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	409	Total	C	N	O	S	0	2	0
			3337	2136	571	608	22			
1	B	411	Total	C	N	O	S	0	4	0
			3359	2149	575	613	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (CCD ID: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 6-{{[(3R,4R)-4-(2-{{[(1R,2S)-2-(3-chlorophenyl)cyclopropyl]amino}ethoxy)pyrrolidin-3-yl]methyl}-4-methylpyridin-2-amine (CCD ID: X2D) (formula: C₂₂H₂₉ClN₄O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	0	0
			28	22	1	4	1		
5	B	1	Total	C	Cl	N	O	0	0
			28	22	1	4	1		

- Molecule 6 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	131	Total	O	0	0
			131	131		
7	B	181	Total	O	0	0
			181	181		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.70Å 110.78Å 163.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.64 – 1.93 38.64 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.1 (38.64-1.93) 99.5 (38.64-1.93)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 1.92Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.190 , 0.235 0.204 , 0.242	Depositor DCC
R_{free} test set	3545 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7193	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, H4B, X2D, ZN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	1/3436 (0.0%)	1.01	5/4661 (0.1%)
1	B	1.00	4/3464 (0.1%)	1.04	10/4696 (0.2%)
All	All	0.96	5/6900 (0.1%)	1.03	15/9357 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	570	MET	CG-SD	-5.96	1.65	1.80
1	B	677	VAL	C-O	5.25	1.30	1.24
1	B	566	ALA	C-O	5.23	1.30	1.23
1	A	615	ASP	CG-OD2	5.15	1.35	1.25
1	B	567	VAL	C-O	-5.06	1.19	1.24

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	707	GLN	CA-C-N	-9.18	110.94	120.03
1	A	707	GLN	C-N-CA	-9.18	110.94	120.03
1	B	707	GLN	CA-C-N	-7.18	112.38	119.78
1	B	707	GLN	C-N-CA	-7.18	112.38	119.78
1	B	570	MET	CA-CB-CG	-6.94	100.21	114.10
1	A	701	THR	CB-CA-C	6.06	117.99	108.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	570	MET	CA-CB-CG	-5.89	102.33	114.10
1	B	512	ALA	CA-C-N	5.67	125.52	119.28
1	B	512	ALA	C-N-CA	5.67	125.52	119.28
1	B	520	LEU	CA-C-N	5.55	126.03	120.14
1	B	520	LEU	C-N-CA	5.55	126.03	120.14
1	B	680	VAL	CA-C-N	5.36	123.57	119.66
1	B	680	VAL	C-N-CA	5.36	123.57	119.66
1	B	486	LYS	N-CA-C	-5.27	100.64	109.24
1	A	589	MET	N-CA-C	-5.15	100.85	109.24

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	513	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3337	0	3253	44	0
1	B	3359	0	3280	31	1
2	A	43	0	30	1	0
2	B	43	0	30	3	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
5	A	28	0	29	2	0
5	B	28	0	29	2	0
6	A	1	0	0	0	0
7	A	131	0	0	3	0
7	B	181	0	0	9	1
All	All	7193	0	6687	75	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:ARG:HB2	1:A:318:LEU:HD21	1.53	0.90
1:A:350:THR:H	1:A:353:GLN:NE2	1.67	0.90
1:A:592:GLU:OE2	5:A:800:X2D:H14	1.81	0.80
2:B:750:HEM:HHC	2:B:750:HEM:HBB2	1.65	0.79
5:A:800:X2D:H26	7:A:1004:HOH:O	1.88	0.72
1:B:592:GLU:OE1	5:B:800:X2D:H14	1.90	0.71
1:B:355:PHE:HB3	7:B:1046:HOH:O	1.90	0.71
1:B:322:LEU:HD22	1:B:699:ARG:HH21	1.58	0.68
1:A:371:ARG:CG	1:A:371:ARG:HH21	2.07	0.67
1:A:382:GLU:HG3	1:A:386:LYS:HE2	1.75	0.67
1:A:513:PRO:O	1:A:515:GLY:N	2.28	0.65
1:A:569:ASN:O	1:A:707:GLN:HG2	1.98	0.63
1:A:332:MET:HE1	1:B:301:LEU:HD22	1.80	0.63
1:B:355:PHE:CB	7:B:1046:HOH:O	2.48	0.59
1:A:380:ARG:HD3	1:A:400:GLU:OE1	2.02	0.59
1:A:371:ARG:HH21	1:A:371:ARG:HG3	1.66	0.59
1:A:317:HIS:O	1:A:320:SER:HB3	2.03	0.58
1:A:601:ASN:HB2	7:A:1020:HOH:O	2.02	0.58
1:A:350:THR:N	1:A:353:GLN:NE2	2.45	0.57
5:B:800:X2D:H26	7:B:1004:HOH:O	2.04	0.57
1:A:371:ARG:CG	1:A:371:ARG:NH2	2.66	0.56
1:A:350:THR:H	1:A:353:GLN:HE21	1.53	0.56
1:A:469:LYS:O	1:A:470:HIS:HD2	1.89	0.55
1:B:478:GLN:HA	1:B:566:ALA:O	2.08	0.53
1:B:507:GLN:NE2	7:B:1114:HOH:O	2.36	0.53
1:A:609:GLU:HG3	7:A:1008:HOH:O	2.07	0.53
1:A:694:GLU:HB3	1:B:335:ILE:HD13	1.90	0.53
2:A:750:HEM:HBB2	2:A:750:HEM:HHC	1.90	0.53
1:B:299:ARG:HB3	1:B:299:ARG:CZ	2.38	0.52
1:B:608:GLU:HB2	7:B:1164:HOH:O	2.11	0.51
1:B:510:TRP:CE2	1:B:521:PRO:HD3	2.46	0.51
1:A:332:MET:CE	1:B:301:LEU:HD22	2.41	0.51
1:B:519:VAL:HG21	1:B:541:VAL:HG11	1.93	0.50
1:A:460:THR:O	1:A:583:PRO:HD2	2.13	0.48
1:A:632:GLU:OE2	1:B:628:GLN:NE2	2.41	0.48
1:B:445:HIS:C	1:B:445:HIS:CD2	2.92	0.48
1:B:615:ASP:HA	7:B:1047:HOH:O	2.14	0.47
1:A:349:ARG:HA	1:A:353:GLN:HE22	1.80	0.47
2:B:750:HEM:HBB2	2:B:750:HEM:CHC	2.39	0.47
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:348:VAL:HG12	7:B:1013:HOH:O	2.15	0.46
1:A:675:ASP:OD2	1:A:675:ASP:C	2.56	0.46
1:B:595:VAL:O	1:B:599:CYS:HB2	2.15	0.46
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.03	0.46
1:A:371:ARG:NH2	1:A:371:ARG:HG2	2.31	0.45
1:B:448:TYR:CD2	1:B:448:TYR:C	2.94	0.45
1:A:469:LYS:C	1:A:470:HIS:CD2	2.95	0.45
1:A:603:ARG:HH11	1:A:603:ARG:HG3	1.81	0.45
1:A:299:ARG:O	1:A:318:LEU:HD11	2.16	0.45
1:A:350:THR:N	1:A:353:GLN:HE21	2.11	0.45
1:B:337:LEU:HD21	1:B:706:TYR:CD2	2.53	0.44
1:A:307:GLU:HG3	7:B:1050:HOH:O	2.17	0.44
1:B:516:ARG:NH2	1:B:557:LEU:O	2.51	0.43
1:B:625:TRP:CZ3	1:B:626:LYS:HG2	2.54	0.43
1:A:349:ARG:HA	1:A:353:GLN:NE2	2.33	0.43
1:A:469:LYS:C	1:A:470:HIS:HD2	2.27	0.42
1:A:628:GLN:HG3	1:B:631:VAL:HG11	2.00	0.42
1:A:696:LEU:HB3	1:B:330:ILE:HD11	2.02	0.42
1:A:320:SER:HA	1:A:700:LEU:HD23	2.00	0.42
1:A:450:THR:HA	1:A:455:LEU:HD22	2.01	0.42
1:B:494:GLY:O	1:B:496:PRO:HD3	2.20	0.42
1:B:535:GLN:HG3	7:B:1090:HOH:O	2.19	0.42
2:B:750:HEM:HHC	2:B:750:HEM:CBB	2.41	0.42
1:A:455:LEU:HD12	1:A:587:TRP:HB3	2.02	0.42
1:B:524:LEU:HD12	1:B:534:PHE:CD1	2.55	0.41
1:A:523:LEU:HD22	1:A:531:PRO:HB2	2.01	0.41
1:A:299:ARG:HB2	1:A:318:LEU:CD2	2.35	0.41
1:A:304:LYS:HE2	1:A:306:TRP:CE3	2.54	0.41
1:A:478:GLN:HA	1:A:566:ALA:O	2.20	0.41
1:B:336:MET:HE2	1:B:336:MET:HB3	1.85	0.41
1:A:403:TYR:CE1	1:A:407:HIS:CE1	3.09	0.41
1:A:510:TRP:CE2	1:A:521:PRO:HD3	2.56	0.41
1:A:686:SER:HA	1:A:691:PHE:CG	2.55	0.40
1:B:423:LYS:O	1:B:457[B]:SER:OG	2.38	0.40
1:B:701:THR:HA	1:B:702:PRO:C	2.46	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:668[B]:CYS:SG	7:B:1012:HOH:O[1_455]	2.07	0.13

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/422 (96%)	393 (97%)	12 (3%)	2 (0%)	24	15
1	B	411/422 (97%)	405 (98%)	6 (2%)	0	100	100
All	All	818/844 (97%)	798 (98%)	18 (2%)	2 (0%)	43	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	514	ARG
1	A	714	HIS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/377 (97%)	358 (98%)	9 (2%)	42	29
1	B	370/377 (98%)	361 (98%)	9 (2%)	43	31
All	All	737/754 (98%)	719 (98%)	18 (2%)	42	31

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	319	LYS
1	A	320	SER
1	A	348	VAL
1	A	371	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	380	ARG
1	A	507	GLN
1	A	555	LYS
1	A	569	ASN
1	A	620	LYS
1	B	299	ARG
1	B	348	VAL
1	B	351	LYS
1	B	353	GLN
1	B	392	SER
1	B	455	LEU
1	B	486	LYS
1	B	608	GLU
1	B	620	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	454	ASN
1	A	470	HIS
1	A	508	GLN
1	A	527	ASN
1	A	569	ASN
1	A	601	ASN
1	A	605	ASN
1	A	628	GLN
1	A	697	ASN
1	A	707	GLN
1	A	712	ASN
1	B	385	ASN
1	B	407	HIS
1	B	454	ASN
1	B	507	GLN
1	B	605	ASN
1	B	642	GLN
1	B	697	ASN
1	B	707	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	H4B	B	760	-	17,18,18	1.26	4 (23%)	14,26,26	1.46	4 (28%)
4	ACT	A	860	-	3,3,3	0.84	0	3,3,3	0.87	0
4	ACT	B	860	-	3,3,3	0.68	0	3,3,3	1.29	0
5	X2D	A	800	-	29,31,31	1.19	3 (10%)	29,43,43	2.30	8 (27%)
3	H4B	A	760	-	17,18,18	1.15	1 (5%)	14,26,26	1.90	3 (21%)
2	HEM	A	750	1	50,50,50	2.10	12 (24%)	67,82,82	1.66	11 (16%)
2	HEM	B	750	1	50,50,50	1.95	9 (18%)	67,82,82	1.81	16 (23%)
5	X2D	B	800	-	29,31,31	1.04	2 (6%)	29,43,43	2.00	8 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	X2D	A	800	-	-	1/15/30/30	0/4/4/4
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
2	HEM	A	750	1	-	1/14/54/54	-
2	HEM	B	750	1	-	1/14/54/54	-
5	X2D	B	800	-	-	2/15/30/30	0/4/4/4

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C3D-C2D	7.48	1.52	1.36
2	A	750	HEM	FE-NB	6.17	2.13	1.94
2	B	750	HEM	C3D-C2D	5.99	1.49	1.36
2	B	750	HEM	FE-NB	5.62	2.12	1.94
2	A	750	HEM	FE-ND	5.53	2.11	1.94
2	B	750	HEM	FE-NC	5.50	2.13	1.95
2	B	750	HEM	FE-ND	4.68	2.09	1.94
2	A	750	HEM	FE-NC	3.85	2.07	1.95
5	A	800	X2D	C15-C13	3.25	1.53	1.49
5	B	800	X2D	C15-C13	3.22	1.53	1.49
3	A	760	H4B	C4-N3	-3.20	1.32	1.38
2	B	750	HEM	CMA-C3A	2.83	1.56	1.50
2	B	750	HEM	CMB-C2B	2.69	1.56	1.50
3	B	760	H4B	C4-N3	-2.65	1.33	1.38
2	A	750	HEM	CAB-C3B	2.64	1.54	1.47
2	A	750	HEM	CMA-C3A	2.52	1.55	1.50
2	A	750	HEM	CAC-C3C	2.47	1.54	1.47
2	A	750	HEM	CMC-C2C	2.45	1.55	1.50
5	A	800	X2D	C23-CL23	2.41	1.80	1.74
2	A	750	HEM	C4C-NC	-2.37	1.35	1.39
2	A	750	HEM	C2A-C3A	-2.36	1.32	1.38
2	A	750	HEM	CMB-C2B	2.34	1.55	1.50
2	A	750	HEM	C4D-ND	-2.32	1.36	1.40
2	B	750	HEM	CAC-C3C	2.32	1.53	1.47
5	B	800	X2D	C15-C14	2.32	1.54	1.50
3	B	760	H4B	O4-C4	2.30	1.27	1.23
5	A	800	X2D	C15-C14	2.21	1.54	1.50
2	B	750	HEM	C2A-C3A	-2.18	1.33	1.38
2	B	750	HEM	CMC-C2C	2.12	1.55	1.50
3	B	760	H4B	C7-N8	2.05	1.48	1.46
3	B	760	H4B	C2-N1	2.04	1.38	1.33

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	800	X2D	C15-C14-C21	-7.23	108.73	122.27
5	B	800	X2D	C15-C14-C21	-6.59	109.94	122.27
2	A	750	HEM	C4D-ND-C1D	6.57	112.99	105.21
2	B	750	HEM	C3B-C4B-NB	-6.21	105.01	109.47
5	A	800	X2D	C02-N01-C06	5.29	122.03	118.07
2	B	750	HEM	C4D-ND-C1D	4.31	110.31	105.21
5	B	800	X2D	C02-N01-C06	4.20	121.21	118.07
3	A	760	H4B	C2-N1-C8A	4.15	120.68	113.36
5	A	800	X2D	C08-C06-N01	4.08	121.57	116.57
2	B	750	HEM	CBD-CAD-C3D	-3.92	101.70	112.53
2	B	750	HEM	CAD-C3D-C4D	3.78	131.28	124.70
2	A	750	HEM	CHC-C1C-NC	3.57	128.34	124.45
2	B	750	HEM	CBA-CAA-C2A	-3.43	103.05	112.53
2	B	750	HEM	C1B-NB-C4B	3.20	108.99	105.21
2	A	750	HEM	C3B-C4B-NB	-3.19	107.17	109.47
2	B	750	HEM	C4B-C3B-C2B	3.10	110.13	107.28
2	A	750	HEM	CBA-CAA-C2A	-3.00	104.25	112.53
5	A	800	X2D	N02-C02-N01	2.99	121.39	116.59
3	A	760	H4B	C4A-C4-N3	2.91	120.11	112.13
2	A	750	HEM	C4C-NC-C1C	2.89	110.53	105.82
3	A	760	H4B	C2-N3-C4	-2.85	119.95	125.11
2	A	750	HEM	C4C-C3C-C2C	2.69	109.15	106.81
2	B	750	HEM	CHC-C4B-NB	2.61	127.23	124.42
5	A	800	X2D	C22-C23-CL23	-2.58	115.93	119.17
3	B	760	H4B	O4-C4-C4A	-2.57	121.05	127.26
2	A	750	HEM	CBD-CAD-C3D	-2.57	105.42	112.53
2	A	750	HEM	CHC-C4B-C3B	2.55	130.17	125.07
2	A	750	HEM	CHD-C4C-C3C	2.53	129.48	125.21
2	B	750	HEM	CAD-C3D-C2D	-2.45	123.28	127.87
5	B	800	X2D	C11-N12-C13	2.45	117.61	114.05
2	B	750	HEM	C4A-CHB-C1B	-2.43	120.54	126.25
5	B	800	X2D	C5'-N1'-C2'	2.42	111.46	105.45
5	B	800	X2D	C08-C06-N01	2.40	119.51	116.57
2	B	750	HEM	C4C-C3C-C2C	2.38	108.88	106.81
3	B	760	H4B	C2-N3-C4	-2.38	120.80	125.11
2	B	750	HEM	C1A-C2A-C3A	2.38	110.55	106.87
2	B	750	HEM	C2A-C1A-NA	-2.35	107.55	110.15
3	B	760	H4B	C2-N1-C8A	2.33	117.47	113.36
5	A	800	X2D	C05-C06-N01	-2.30	120.11	122.73
2	B	750	HEM	CMD-C2D-C1D	2.29	128.62	125.03
5	B	800	X2D	C10-C11-N12	2.22	116.97	111.50
5	A	800	X2D	C26-C25-C24	-2.19	117.42	120.24
3	B	760	H4B	C4A-C4-N3	2.18	118.13	112.13

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	C4C-NC-C1C	2.17	109.37	105.82
5	A	800	X2D	O09-C4'-C5'	-2.13	105.19	111.23
2	A	750	HEM	CAB-C3B-C2B	-2.13	121.52	128.43
5	B	800	X2D	C05-C06-N01	-2.09	120.35	122.73
5	B	800	X2D	N02-C02-N01	2.06	119.90	116.59
2	B	750	HEM	CHA-C1A-NA	2.03	127.54	123.86
2	A	750	HEM	CAA-CBA-CGA	-2.02	108.30	113.67

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	800	X2D	C15-C14-C21-C26
2	A	750	HEM	C4B-C3B-CAB-CBB
2	B	750	HEM	C4B-C3B-CAB-CBB
5	B	800	X2D	C15-C14-C21-C22
5	A	800	X2D	O09-C10-C11-N12


There are no ring outliers.


4 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	800	X2D	2	0
2	A	750	HEM	1	0
2	B	750	HEM	3	0
5	B	800	X2D	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	409/422 (96%)	0.39	16 (3%) 43 49	27, 51, 90, 117	2 (0%)
1	B	411/422 (97%)	0.08	9 (2%) 62 69	20, 42, 66, 83	4 (0%)
All	All	820/844 (97%)	0.23	25 (3%) 52 58	20, 46, 83, 117	6 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PHE	4.5
1	B	350	THR	4.3
1	A	300	PHE	4.1
1	A	321	THR	4.0
1	A	322	LEU	3.6
1	B	322	LEU	3.6
1	A	355	PHE	3.4
1	A	716	TRP	3.0
1	B	348	VAL	2.9
1	A	488	PRO	2.9
1	B	321	THR	2.7
1	A	715	VAL	2.7
1	A	552	ASP	2.7
1	A	348	VAL	2.6
1	A	713	THR	2.6
1	B	352	ASP	2.4
1	A	601	ASN	2.3
1	A	318	LEU	2.3
1	A	551	PHE	2.2
1	B	601	ASN	2.1
1	B	355	PHE	2.1
1	A	330	ILE	2.0
1	B	713	THR	2.0
1	A	490	GLY	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	351	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	X2D	A	800	28/28	0.89	0.10	27,33,36,46	0
4	ACT	A	860	4/4	0.90	0.15	59,60,62,64	0
4	ACT	B	860	4/4	0.91	0.12	48,50,51,52	0
5	X2D	B	800	28/28	0.92	0.09	35,41,45,52	0
3	H4B	A	760	17/17	0.94	0.07	30,33,40,40	0
3	H4B	B	760	17/17	0.96	0.06	28,33,38,40	0
2	HEM	A	750	43/43	0.97	0.07	29,34,38,39	0
2	HEM	B	750	43/43	0.98	0.06	25,32,39,44	0
6	ZN	A	900	1/1	1.00	0.03	38,38,38,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around X2D A 800:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around X2D B 800:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 750:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.