



Full wwPDB NMR Structure Validation Report ⓘ

Mar 9, 2026 – 04:46 AM UTC

PDB ID : 2RRN / pdb_00002rrn
BMRB ID : 11426
Title : Solution structure of SecDF periplasmic domain P4
Authors : Tanaka, T.; Tsukazaki, T.; Echizen, Y.; Nureki, O.; Kohno, T.
Deposited on : 2011-01-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 87%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	92	 67% 15% • 16%

2 Ensemble composition and analysis

This entry contains 20 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:480-A:556 (77)	0.30	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 3 single-model clusters were found.

Cluster number	Models
1	4, 6, 8, 10, 11, 12, 17, 18, 20
2	1, 5, 15
3	7, 9, 14
4	2, 3
Single-model clusters	13; 16; 19

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1455 atoms, of which 730 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Probable SecDF protein-export membrane protein.

Mol	Chain	Residues	Atoms						Trace
1	A	92	Total	C	H	N	O	S	0
			1455	464	730	121	139	1	

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	468	MET	-	initiating methionine	UNP Q5SKE6
A	469	VAL	-	expression tag	UNP Q5SKE6

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Probable SecDF protein-export membrane protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.2 Score per residue for model 2

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.3 Score per residue for model 3

- Molecule 1: Probable SecDF protein-export membrane protein



4.2.4 Score per residue for model 4

- Molecule 1: Probable SecDF protein-export membrane protein



4.2.5 Score per residue for model 5

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.6 Score per residue for model 6 (medoid)

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.7 Score per residue for model 7

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.8 Score per residue for model 8

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.9 Score per residue for model 9

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.10 Score per residue for model 10

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.11 Score per residue for model 11

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.12 Score per residue for model 12

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.13 Score per residue for model 13

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.14 Score per residue for model 14

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.15 Score per residue for model 15

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.16 Score per residue for model 16

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.17 Score per residue for model 17

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.18 Score per residue for model 18

- Molecule 1: Probable SecDF protein-export membrane protein




4.2.19 Score per residue for model 19

- Molecule 1: Probable SecDF protein-export membrane protein



4.2.20 Score per residue for model 20

- Molecule 1: Probable SecDF protein-export membrane protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *Energy Minimization*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
TALOS	structure solution	
ProcheckNMR	structure solution	
CYANA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1086
Number of shifts mapped to atoms	1086
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	87%

6 Model quality

6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	613	627	627	12±2
All	All	12260	12540	12540	246

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:505:PHE:CD2	1:A:540:LEU:HD22	0.68	2.24	3	6
1:A:527:VAL:HG12	1:A:529:LEU:HD21	0.67	1.65	16	17
1:A:487:ALA:HB3	1:A:525:PHE:CE2	0.67	2.25	13	8
1:A:500:LEU:HD12	1:A:512:ILE:HD11	0.65	1.68	15	4
1:A:482:ALA:HB3	1:A:556:GLU:CG	0.62	2.24	19	7
1:A:496:LEU:HB3	1:A:512:ILE:HD13	0.61	1.73	15	15
1:A:482:ALA:HB1	1:A:526:LEU:HD21	0.61	1.70	16	1
1:A:482:ALA:HB1	1:A:526:LEU:HD11	0.59	1.74	9	5
1:A:482:ALA:HB3	1:A:556:GLU:HG2	0.59	1.73	7	5
1:A:503:LYS:NZ	1:A:543:LEU:HD13	0.58	2.13	11	1
1:A:482:ALA:HB3	1:A:556:GLU:HG3	0.57	1.76	2	1
1:A:524:GLU:OE1	1:A:553:LEU:HD12	0.57	1.99	9	1
1:A:484:THR:HG22	1:A:524:GLU:CD	0.57	2.25	12	1
1:A:503:LYS:NZ	1:A:543:LEU:HD22	0.56	2.16	14	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:496:LEU:CB	1:A:512:ILE:HD13	0.56	2.31	19	17
1:A:486:ARG:HB2	1:A:553:LEU:HD11	0.55	1.77	7	2
1:A:487:ALA:HB3	1:A:525:PHE:CD2	0.54	2.37	3	8
1:A:527:VAL:CG1	1:A:529:LEU:HD21	0.54	2.33	16	20
1:A:484:THR:HG22	1:A:524:GLU:OE2	0.53	2.03	9	1
1:A:510:ALA:HB1	1:A:528:LYS:O	0.53	2.04	10	14
1:A:519:THR:HG23	1:A:522:TYR:O	0.53	2.04	2	13
1:A:496:LEU:HD13	1:A:512:ILE:CG2	0.52	2.35	10	9
1:A:493:VAL:HG13	1:A:512:ILE:HB	0.51	1.83	12	1
1:A:510:ALA:HB1	1:A:529:LEU:HD23	0.50	1.83	15	20
1:A:527:VAL:HG12	1:A:529:LEU:CD2	0.49	2.37	16	8
1:A:484:THR:HG22	1:A:524:GLU:CG	0.49	2.38	9	1
1:A:524:GLU:CD	1:A:553:LEU:HD12	0.48	2.34	9	1
1:A:488:GLU:HG3	1:A:491:VAL:HG23	0.48	1.85	4	1
1:A:486:ARG:HD3	1:A:553:LEU:HD11	0.48	1.86	10	1
1:A:503:LYS:HZ1	1:A:543:LEU:HD13	0.48	1.69	11	1
1:A:486:ARG:CD	1:A:553:LEU:HD11	0.47	2.39	10	1
1:A:488:GLU:HG2	1:A:491:VAL:HG23	0.47	1.85	3	2
1:A:503:LYS:HE2	1:A:543:LEU:HD13	0.47	1.87	19	2
1:A:484:THR:OG1	1:A:526:LEU:HD23	0.47	2.09	16	1
1:A:484:THR:HG23	1:A:526:LEU:HD23	0.46	1.88	12	2
1:A:526:LEU:HD13	1:A:526:LEU:C	0.45	2.36	16	1
1:A:491:VAL:HG11	1:A:548:LEU:HD21	0.45	1.87	2	2
1:A:517:ALA:HB1	1:A:518:PRO:HD2	0.44	1.88	9	20
1:A:488:GLU:CG	1:A:491:VAL:HG23	0.44	2.42	12	1
1:A:528:LYS:O	1:A:529:LEU:HD23	0.43	2.13	1	4
1:A:540:LEU:HD11	1:A:544:PHE:CE2	0.43	2.48	3	2
1:A:515:VAL:HG11	1:A:524:GLU:OE2	0.43	2.13	13	1
1:A:503:LYS:CE	1:A:543:LEU:HD13	0.43	2.42	19	1
1:A:506:PRO:O	1:A:510:ALA:HB2	0.43	2.13	16	2
1:A:484:THR:HG23	1:A:526:LEU:CD2	0.43	2.44	10	5
1:A:517:ALA:HB2	1:A:524:GLU:CD	0.43	2.39	11	1
1:A:543:LEU:O	1:A:543:LEU:HD23	0.42	2.14	3	1
1:A:486:ARG:HB2	1:A:553:LEU:HD21	0.42	1.90	3	1
1:A:487:ALA:HB3	1:A:525:PHE:CE1	0.41	2.51	6	1
1:A:503:LYS:HZ3	1:A:543:LEU:HD22	0.40	1.76	14	1
1:A:496:LEU:HD13	1:A:512:ILE:HG21	0.40	1.92	15	1
1:A:496:LEU:HB2	1:A:512:ILE:HD13	0.40	1.93	6	1
1:A:484:THR:HB	1:A:554:ALA:HB3	0.40	1.94	16	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	77/92 (84%)	70±2 (91±2%)	7±2 (9±2%)	0±0 (0±0%)	100	100
All	All	1540/1840 (84%)	1394 (91%)	146 (9%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	65/77 (84%)	59±1 (90±2%)	6±1 (10±2%)	10	55
All	All	1300/1540 (84%)	1174 (90%)	126 (10%)	10	55

All 23 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	548	LEU	20
1	A	526	LEU	19
1	A	513	THR	15
1	A	484	THR	14
1	A	546	SER	9
1	A	516	GLN	8
1	A	549	LYS	6
1	A	503	LYS	4
1	A	486	ARG	3
1	A	509	GLU	3
1	A	533	SER	3
1	A	508	LYS	3
1	A	543	LEU	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	534	ASP	3
1	A	528	LYS	2
1	A	523	ARG	2
1	A	532	LEU	2
1	A	538	LEU	2
1	A	556	GLU	1
1	A	488	GLU	1
1	A	541	GLU	1
1	A	542	ARG	1
1	A	494	GLU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 87% for the well-defined parts and 85% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1086
Number of shifts mapped to atoms	1086
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	9

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	92	-0.25 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	86	0.09 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}'$	89	-0.03 ± 0.09	None needed (< 0.5 ppm)
^{15}N	86	-0.41 ± 0.29	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 87%, i.e. 942 atoms were assigned a chemical shift out of a possible 1089. 0 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	375/378 (99%)	152/152 (100%)	151/154 (98%)	72/72 (100%)
Sidechain	555/653 (85%)	367/424 (87%)	178/201 (89%)	10/28 (36%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	12/58 (21%)	12/28 (43%)	0/30 (0%)	0/0 (—%)
Overall	942/1089 (87%)	531/604 (88%)	329/385 (85%)	82/100 (82%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 85%, i.e. 1086 atoms were assigned a chemical shift out of a possible 1274. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	449/456 (98%)	182/185 (98%)	181/184 (98%)	86/87 (99%)
Sidechain	619/731 (85%)	410/476 (86%)	199/226 (88%)	10/29 (34%)
Aromatic	18/87 (21%)	18/42 (43%)	0/45 (0%)	0/0 (—%)
Overall	1086/1274 (85%)	610/703 (87%)	380/455 (84%)	96/116 (83%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	523	ARG	NE	117.21	76.53 – 92.65	20.2
1	A	486	ARG	NE	116.58	76.53 – 92.65	19.8
1	A	498	ARG	NE	116.58	76.53 – 92.65	19.8
1	A	536	ARG	NE	115.60	76.53 – 92.65	19.2
1	A	542	ARG	NE	115.35	76.53 – 92.65	19.1
1	A	497	ARG	NE	114.21	76.53 – 92.65	18.4
1	A	537	ARG	NE	113.11	76.53 – 92.65	17.7
1	A	537	ARG	CZ	179.49	141.81 – 177.93	5.4
1	A	486	ARG	HD2	1.97	1.97 – 4.26	-5.0

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

