



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:36 AM UTC

PDB ID : 3RRC / pdb_00003rrc
Title : Crystal Structure of Region II from Plasmodium vivax Duffy Binding Protein
Authors : Tolia, N.H.
Deposited on : 2011-04-29
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	11% 74% 12% • 13%
1	B	317	4% 74% 12% • 12%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9711 atoms, of which 4660 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Duffy receptor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	276	Total	C	H	N	O	S	0	1	0
			4584	1453	2284	404	425	18			
1	B	279	Total	C	H	N	O	S	0	0	0
			4633	1471	2310	409	425	18			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	209	MET	-	expression tag	UNP P22290
A	210	GLY	-	expression tag	UNP P22290
B	209	MET	-	expression tag	UNP P22290
B	210	GLY	-	expression tag	UNP P22290

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	171	Total	O	0	0
			171	171		
4	B	193	Total	O	0	0
			193	193		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Duffy receptor



• Molecule 1: Duffy receptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.50Å 66.97Å 91.83Å 90.00° 108.06° 90.00°	Depositor
Resolution (Å)	19.70 – 1.95 19.70 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.5 (19.70-1.95) 99.4 (19.70-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 1.94Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.195 , 0.241 0.187 , 0.233	Depositor DCC
R_{free} test set	2559 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.389	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 58.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9711	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.74 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2428e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/2345	0.75	2/3152 (0.1%)
1	B	0.59	0/2367	0.78	2/3181 (0.1%)
All	All	0.58	0/4712	0.77	4/6333 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	386	GLU	CA-C-N	-6.99	112.76	119.89
1	B	386	GLU	C-N-CA	-6.99	112.76	119.89
1	A	386	GLU	CA-C-N	-5.23	114.39	119.78
1	A	386	GLU	C-N-CA	-5.23	114.39	119.78

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2300	2284	2284	33	0
1	B	2323	2310	2310	34	0
2	A	24	36	36	0	0
2	B	20	30	30	0	0
3	A	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	10	0	0	0	0
4	A	171	0	0	1	0
4	B	193	0	0	1	0
All	All	5051	4660	4660	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:295:TYR:HH	1:A:377:CYS:N	1.90	0.69
1:B:468:THR:O	1:B:468:THR:CG2	2.42	0.68
1:A:358:TRP:CZ2	1:A:362:MET:HG3	2.33	0.64
1:A:277:ILE:HD12	1:B:274:ARG:HD2	1.82	0.60
1:A:358:TRP:CH2	1:A:362:MET:HG3	2.36	0.60
1:B:271:TYR:CE2	1:B:275:LYS:HD2	2.38	0.59
1:B:271:TYR:CZ	1:B:275:LYS:HD2	2.39	0.57
1:A:314:ILE:O	1:A:346:ARG:HD2	2.05	0.57
1:A:266:THR:N	1:B:367:LYS:CE	2.68	0.56
1:B:321:GLY:O	1:B:326:LYS:HD3	2.06	0.56
1:A:271:TYR:CZ	1:A:275:LYS:HE2	2.42	0.55
1:A:386:GLU:HG2	1:A:390:TYR:HB2	1.89	0.55
1:A:266:THR:N	1:B:367:LYS:HD2	2.21	0.55
1:B:398:ARG:HD2	1:B:481:GLU:OE2	2.06	0.54
1:B:468:THR:O	1:B:468:THR:HG22	2.08	0.54
1:A:277:ILE:CD1	1:B:274:ARG:HD2	2.38	0.54
1:B:249:GLU:HB3	1:B:275:LYS:HB3	1.88	0.53
1:A:314:ILE:O	1:A:346:ARG:CD	2.57	0.53
1:A:266:THR:C	1:A:268:ARG:H	2.17	0.53
1:B:213:VAL:HG22	1:B:324:TYR:CE1	2.44	0.53
1:A:429:VAL:HG12	1:A:431:PRO:HD2	1.91	0.51
1:B:298:ASP:OD1	1:B:398:ARG:NH2	2.43	0.51
1:B:428:LYS:O	1:B:430:PRO:HD3	2.10	0.51
1:A:358:TRP:CE3	1:A:379:LEU:HD12	2.45	0.51
1:B:461:LYS:O	1:B:464:GLU:HG3	2.11	0.50
1:A:266:THR:N	1:B:367:LYS:HE3	2.26	0.50
1:A:379:LEU:HD23	1:A:379:LEU:C	2.36	0.50
1:B:473:THR:HB	1:B:474:PRO:HD2	1.94	0.49
1:B:368:ARG:N	1:B:368:ARG:HD3	2.27	0.49
1:B:253:LEU:HD21	1:B:335:ILE:CD1	2.42	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:THR:N	1:B:367:LYS:CD	2.76	0.48
1:B:429:VAL:HG12	1:B:431:PRO:HD2	1.94	0.48
1:A:221:ARG:HD2	1:A:240:ASP:OD2	2.14	0.48
1:B:253:LEU:HD21	1:B:335:ILE:HD11	1.94	0.48
1:B:274:ARG:NH1	4:B:115:HOH:O	2.46	0.48
1:A:464:GLU:HG3	1:A:466:VAL:HG23	1.96	0.46
1:A:274:ARG:HH11	1:A:274:ARG:HG3	1.79	0.46
1:A:224:ARG:CG	1:A:403:GLU:OE2	2.63	0.46
1:A:464:GLU:O	1:A:465:LYS:C	2.60	0.45
1:A:401:VAL:CG1	1:A:482:LEU:HD13	2.47	0.45
1:A:378:LYS:HB2	1:A:381:VAL:HB	1.99	0.44
1:B:228:TRP:CZ3	1:B:239:PRO:HD3	2.52	0.44
1:A:401:VAL:HG11	1:A:482:LEU:HD13	2.00	0.43
1:B:411:LEU:HD11	1:B:505:CYS:SG	2.58	0.43
1:B:394:ARG:O	1:B:398:ARG:HG3	2.18	0.42
1:A:364:SER:HB3	4:A:560:HOH:O	2.19	0.42
1:B:468:THR:O	1:B:468:THR:HG23	2.19	0.42
1:A:295:TYR:CE1	1:A:365:VAL:HG11	2.54	0.42
1:A:249:GLU:HB3	1:A:275:LYS:HD2	2.01	0.42
1:B:407:GLU:HG3	1:B:442:TRP:CD2	2.55	0.42
1:B:253:LEU:HD23	1:B:331:ASN:HB3	2.02	0.41
1:A:429:VAL:CG1	1:A:431:PRO:HD2	2.50	0.41
1:B:430:PRO:HB2	1:B:431:PRO:HD3	2.02	0.41
1:B:216:ASN:HB2	1:B:244:GLN:O	2.20	0.41
1:B:301:LYS:CD	1:B:398:ARG:NE	2.84	0.41
1:B:386:GLU:O	1:B:387:PRO:C	2.60	0.41
1:A:295:TYR:CD1	1:A:365:VAL:HG11	2.57	0.40
1:A:340:GLU:HG3	1:A:341:LYS:HG3	2.02	0.40
1:A:345:ARG:HA	1:A:345:ARG:HD3	1.97	0.40
1:A:266:THR:C	1:A:268:ARG:N	2.79	0.40
1:B:249:GLU:O	1:B:275:LYS:HE3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/317 (86%)	262 (97%)	8 (3%)	1 (0%)	30	21
1	B	273/317 (86%)	264 (97%)	8 (3%)	1 (0%)	30	21
All	All	544/634 (86%)	526 (97%)	16 (3%)	2 (0%)	30	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	339	ASP
1	A	267	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/292 (87%)	247 (97%)	7 (3%)	38	30
1	B	255/292 (87%)	245 (96%)	10 (4%)	28	18
All	All	509/584 (87%)	492 (97%)	17 (3%)	33	24

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	214	MET
1	A	215	LYS
1	A	223	ARG
1	A	274	ARG
1	A	320	GLU
1	A	383	VAL
1	A	484	GLU
1	B	211	ASN
1	B	213	VAL
1	B	253	LEU
1	B	320	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	330	ASN
1	B	374	ILE
1	B	384	ASN
1	B	464	GLU
1	B	468	THR
1	B	471	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	11	-	3,3,3	0.47	0	2,2,2	0.27	0
3	PO4	A	1	-	4,4,4	0.92	0	6,6,6	0.56	0
2	EDO	A	3	-	3,3,3	0.41	0	2,2,2	0.25	0
2	EDO	B	9	-	3,3,3	0.44	0	2,2,2	0.30	0
2	EDO	A	2	-	3,3,3	0.47	0	2,2,2	0.24	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	B	1	-	3,3,3	0.48	0	2,2,2	0.46	0
3	PO4	B	3	-	4,4,4	1.04	0	6,6,6	0.72	0
2	EDO	A	13	-	3,3,3	0.46	0	2,2,2	0.35	0
3	PO4	B	526	-	4,4,4	1.02	0	6,6,6	0.42	0
2	EDO	B	12	-	3,3,3	0.42	0	2,2,2	0.68	0
2	EDO	A	7	-	3,3,3	0.44	0	2,2,2	0.29	0
2	EDO	A	5	-	3,3,3	0.43	0	2,2,2	0.47	0
2	EDO	B	4	-	3,3,3	0.40	0	2,2,2	0.33	0
2	EDO	A	6	-	3,3,3	0.44	0	2,2,2	0.70	0
3	PO4	A	526	-	4,4,4	0.95	0	6,6,6	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	11	-	-	1/1/1/1	-
2	EDO	A	3	-	-	1/1/1/1	-
2	EDO	B	9	-	-	0/1/1/1	-
2	EDO	A	2	-	-	0/1/1/1	-
2	EDO	B	1	-	-	0/1/1/1	-
2	EDO	A	13	-	-	1/1/1/1	-
2	EDO	B	12	-	-	0/1/1/1	-
2	EDO	A	7	-	-	0/1/1/1	-
2	EDO	A	5	-	-	1/1/1/1	-
2	EDO	B	4	-	-	0/1/1/1	-
2	EDO	A	6	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	5	EDO	O1-C1-C2-O2
2	A	13	EDO	O1-C1-C2-O2
2	A	3	EDO	O1-C1-C2-O2
2	A	6	EDO	O1-C1-C2-O2
2	B	11	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/317 (87%)	0.39	36 (13%) 7 8	12, 39, 115, 208	1 (0%)
1	B	279/317 (88%)	-0.02	12 (4%) 40 46	12, 32, 82, 129	0
All	All	555/634 (87%)	0.19	48 (8%) 16 19	12, 35, 97, 208	1 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	219	TYR	6.2
1	A	379	LEU	4.9
1	B	271	TYR	4.9
1	A	278	TYR	4.9
1	B	214	MET	4.8
1	A	421	TYR	4.6
1	A	466	VAL	4.2
1	B	468	THR	4.2
1	B	210	GLY	4.1
1	A	213	VAL	4.1
1	A	377	CYS	4.1
1	A	253	LEU	3.8
1	A	365	VAL	3.8
1	A	381	VAL	3.8
1	B	219	TYR	3.7
1	A	214	MET	3.7
1	A	363	TYR	3.5
1	A	218	ASN	3.4
1	A	464	GLU	3.3
1	A	266	THR	3.3
1	A	380	ASN	3.3
1	B	248	LYS	3.1
1	A	324	TYR	3.0
1	A	378	LYS	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	290	LEU	2.9
1	A	465	LYS	2.8
1	A	210	GLY	2.7
1	A	289	LYS	2.6
1	A	382	ALA	2.6
1	A	383	VAL	2.6
1	A	469	ALA	2.5
1	A	358	TRP	2.4
1	A	226	ARG	2.4
1	A	344	GLN	2.4
1	A	351	ASN	2.4
1	A	345	ARG	2.4
1	B	421	TYR	2.3
1	A	340	GLU	2.3
1	B	464	GLU	2.2
1	A	341	LYS	2.2
1	A	347	LYS	2.2
1	B	253	LEU	2.2
1	A	267	PHE	2.2
1	A	364	SER	2.2
1	B	341	LYS	2.2
1	A	484	GLU	2.0
1	B	367	LYS	2.0
1	B	471	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	A	13	4/4	0.79	0.15	57,68,75,76	0
3	PO4	B	526	5/5	0.80	0.11	76,77,82,84	0
3	PO4	A	1	5/5	0.84	0.11	92,95,98,98	0
2	EDO	A	5	4/4	0.86	0.11	39,48,54,58	0
3	PO4	A	526	5/5	0.87	0.10	81,86,93,96	0
2	EDO	A	7	4/4	0.89	0.12	38,56,65,73	0
2	EDO	B	12	4/4	0.90	0.11	41,49,59,59	0
2	EDO	B	11	4/4	0.91	0.13	39,52,59,64	0
2	EDO	B	9	4/4	0.92	0.08	37,44,53,53	0
2	EDO	A	6	4/4	0.95	0.08	24,32,42,42	0
2	EDO	A	3	4/4	0.96	0.07	28,41,49,49	0
2	EDO	B	1	4/4	0.96	0.08	26,32,63,75	0
3	PO4	B	3	5/5	0.96	0.07	32,38,48,52	0
2	EDO	B	4	4/4	0.96	0.06	27,35,42,42	0
2	EDO	A	2	4/4	0.98	0.04	18,26,32,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.