



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 11:33 PM UTC

PDB ID : 6RRU / pdb_00006rru
Title : GOLGI ALPHA-MANNOSIDASE II in complex with (5R,6R,7S,8S)-5,6,7,8-tetrahydro-5-(hydroxymethyl)-3-(3-phenylpropyl)imidazo[1,2-a]pyridine-6,7,8-triol
Authors : Armstrong, Z.; Lahav, D.; Johnson, R.; Kuo, C.L.; Beenakker, T.J.M.; de Boer, C.; Wong, C.S.; van Rijssel, E.R.; Debets, M.; Geurink, P.P.; Ovaa, H.; van der Stelt, M.; Codee, J.D.C.; Aerts, J.M.F.G.; Wu, L.; Overkleeft, H.S.; Davies, G.J.
Deposited on : 2019-05-20
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1033	8% 87% 9% .

Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8689 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-mannosidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	998	8154	5188	1428	1496	42	0	16	0

- Molecule 2 is (5 {R},6 {R},7 {S},8 {S})-5-(hydroxymethyl)-2-(3-phenylpropyl)-5,6,7,8-tetrahydroimidazo[1,2-a]pyridine-6,7,8-triol (CCD ID: KG8) (formula: C₁₇H₂₂N₂O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	23	17	2	4	0	0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	510	Total	O	0	0
			510	510		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-mannosidase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.96Å 91.91Å 131.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.51 – 1.90 75.39 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (75.51-1.90) 99.9 (75.39-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.192 , 0.242 0.200 , 0.246	Depositor DCC
R_{free} test set	4306 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtriage
Anisotropy	0.615	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.008 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8689	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, KG8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	0/8404	1.29	9/11401 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	46	LEU	CA-C-N	5.81	127.99	120.44
1	A	46	LEU	C-N-CA	5.81	127.99	120.44
1	A	637	SER	CA-C-N	5.73	128.28	120.54
1	A	637	SER	C-N-CA	5.73	128.28	120.54
1	A	689	SER	CA-C-N	5.73	128.53	120.28
1	A	689	SER	C-N-CA	5.73	128.53	120.28
1	A	836	PHE	CA-CB-CG	5.62	119.42	113.80
1	A	409	ASP	CA-CB-CG	5.27	117.87	112.60
1	A	211	THR	CA-CB-OG1	-5.10	101.95	109.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8154	0	7963	49	0
2	A	23	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
4	A	510	0	0	3	0
All	All	8689	0	7963	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:985:MET:HE1	4:A:1565:HOH:O	1.81	0.78
1:A:1008:ARG:NH1	1:A:1010:GLU:OE1	2.19	0.76
1:A:698:GLN:NE2	1:A:700:THR:O	2.25	0.68
1:A:640:LYS:HD3	1:A:646:TYR:HE1	1.62	0.64
1:A:174:HIS:CE1	1:A:176:ARG:HD3	2.35	0.62
1:A:844:LEU:HD11	1:A:894:LEU:HD11	1.85	0.59
1:A:640:LYS:HD3	1:A:646:TYR:CE1	2.39	0.57
1:A:839:ASP:O	1:A:904:ARG:NH1	2.41	0.54
1:A:344:PHE:H	1:A:349:GLU:CD	2.18	0.51
1:A:93:PRO:HD2	1:A:470:HIS:CE1	2.46	0.51
1:A:43:VAL:O	1:A:400:LEU:HA	2.10	0.51
1:A:1008:ARG:HB3	1:A:1042:SER:HB2	1.93	0.50
1:A:60:GLY:O	1:A:63:LYS:HD2	2.12	0.50
1:A:116:ARG:HD3	1:A:347:ASN:HD21	1.75	0.50
1:A:956:ASP:OD1	1:A:981[B]:ARG:NH1	2.45	0.50
1:A:96:ILE:HG23	1:A:479:LYS:HE2	1.93	0.50
1:A:979[A]:LEU:HD21	1:A:999:VAL:HG11	1.93	0.49
1:A:981[B]:ARG:HH21	1:A:1030:GLU:HG2	1.77	0.48
1:A:902:CYS:HB3	1:A:985:MET:HG2	1.96	0.48
1:A:566:GLU:HA	1:A:622:ALA:O	2.15	0.47
1:A:651:LEU:CD1	1:A:659:LEU:CD1	2.92	0.47
1:A:743:PRO:HB2	1:A:758:VAL:HG23	1.96	0.46
1:A:155:ASN:C	1:A:155:ASN:OD1	2.59	0.46
1:A:567:GLN:NE2	4:A:1226:HOH:O	2.47	0.46
1:A:740:LEU:HD22	1:A:760:LEU:HD22	1.99	0.45
1:A:698:GLN:HE21	1:A:700:THR:C	2.23	0.45
1:A:979[B]:LEU:HD23	1:A:1031:VAL:HG13	1.99	0.45
1:A:498:GLN:HG2	1:A:526:LEU:HD13	1.99	0.44
1:A:117:HIS:ND1	1:A:347:ASN:OD1	2.50	0.44
1:A:419:TYR:C	1:A:426:LYS:HE2	2.43	0.44
1:A:592:VAL:HA	1:A:617:ARG:O	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:957:LEU:HD11	1:A:979[B]:LEU:HG	2.00	0.43
1:A:304:ARG:NH1	4:A:1240:HOH:O	2.51	0.43
1:A:498:GLN:HG2	1:A:526:LEU:CD1	2.48	0.43
1:A:579:SER:HB3	1:A:641:PRO:HB3	2.01	0.43
1:A:153:VAL:HG21	1:A:193[B]:MET:HE3	2.00	0.42
1:A:251:TRP:CD1	1:A:251:TRP:C	2.97	0.42
1:A:341:ASP:OD2	2:A:1101:KG8:N2	2.52	0.42
1:A:794:LEU:O	1:A:861:GLU:HA	2.20	0.41
1:A:204:ASP:H	1:A:205:PRO:HD3	1.86	0.41
1:A:87:PRO:HA	1:A:338:LEU:O	2.21	0.41
1:A:678:LEU:HD12	1:A:769[B]:MET:HE1	2.03	0.41
1:A:743:PRO:HB2	1:A:758:VAL:CG2	2.51	0.41
1:A:472:ASP:OD2	2:A:1101:KG8:O1	2.39	0.40
1:A:655:ASN:O	1:A:656:PRO:C	2.65	0.40
1:A:698:GLN:NE2	1:A:700:THR:C	2.79	0.40
1:A:651:LEU:HD13	1:A:659:LEU:HD12	2.03	0.40
1:A:570:ASP:HA	1:A:618:ILE:O	2.20	0.40
1:A:650:LEU:HB3	1:A:747:VAL:HB	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1004/1033 (97%)	966 (96%)	35 (4%)	3 (0%)	36 29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	95	TRP
1	A	708	VAL
1	A	204	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	903/918 (98%)	881 (98%)	22 (2%)	43 38

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	117	HIS
1	A	154	LYS
1	A	275	CYS
1	A	317	LEU
1	A	526	LEU
1	A	585	ASN
1	A	600	HIS
1	A	654	LYS
1	A	682	ASN
1	A	696	SER
1	A	699	LEU
1	A	755	SER
1	A	758	VAL
1	A	848[A]	THR
1	A	848[B]	THR
1	A	859	SER
1	A	901	ASN
1	A	906	SER
1	A	940	ILE
1	A	943	GLN
1	A	1009	CYS
1	A	1031	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	A	119	HIS
1	A	148	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	242	GLN
1	A	249	GLN
1	A	309	GLN
1	A	380	GLN
1	A	394	GLN
1	A	610	GLN
1	A	698	GLN
1	A	709	HIS
1	A	720	HIS
1	A	812	GLN
1	A	919	HIS
1	A	943	GLN
1	A	1019	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	KG8	A	1101	3	24,25,25	0.50	0	24,35,35	1.04	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	KG8	A	1101	3	-	2/8/28/28	0/3/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	KG8	C5-C4-C3	-2.65	108.94	112.93
2	A	1101	KG8	C17-C7-N2	-2.41	108.59	110.65

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1101	KG8	C11-C10-C9-C8
2	A	1101	KG8	C7-C8-C9-C10

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1101	KG8	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	998/1033 (96%)	0.42	79 (7%)	18 20	10, 32, 58, 103	16 (1%)

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	988	GLY	4.4
1	A	659	LEU	4.2
1	A	606	THR	4.0
1	A	741	GLY	4.0
1	A	652	LEU	3.9
1	A	692	GLY	3.8
1	A	671	GLY	3.8
1	A	654	LYS	3.6
1	A	706	VAL	3.5
1	A	656	PRO	3.4
1	A	295	LEU	3.4
1	A	38	VAL	3.4
1	A	549	ILE	3.2
1	A	738	VAL	3.2
1	A	683	GLY	3.2
1	A	687	ALA	3.1
1	A	676	ILE	3.1
1	A	607	ILE	3.0
1	A	703	SER	3.0
1	A	673	PRO	3.0
1	A	1025	GLY	2.9
1	A	784	SER	2.9
1	A	1030	GLU	2.9
1	A	797	HIS	2.9
1	A	906	SER	2.9
1	A	347	ASN	2.9
1	A	670	PHE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	688	PHE	2.8
1	A	858	ALA	2.8
1	A	971	LYS	2.8
1	A	745	VAL	2.8
1	A	699	LEU	2.8
1	A	702	ASP	2.8
1	A	668	VAL	2.7
1	A	744	VAL	2.7
1	A	117	HIS	2.7
1	A	900	ASN	2.7
1	A	996	LYS	2.7
1	A	794	LEU	2.6
1	A	708	VAL	2.6
1	A	651	LEU	2.6
1	A	740	LEU	2.6
1	A	707	PRO	2.6
1	A	710	PHE	2.6
1	A	785	LEU	2.6
1	A	650	LEU	2.5
1	A	660	PRO	2.5
1	A	760	LEU	2.5
1	A	608	HIS	2.5
1	A	720	HIS	2.5
1	A	691	GLN	2.5
1	A	600	HIS	2.4
1	A	841	ASN	2.4
1	A	803	ILE	2.3
1	A	661	LEU	2.3
1	A	684	PRO	2.3
1	A	954[A]	ARG	2.3
1	A	704	PRO	2.3
1	A	896	LEU	2.3
1	A	669	LYS	2.2
1	A	392	ALA	2.2
1	A	709	HIS	2.2
1	A	1029	PRO	2.2
1	A	116	ARG	2.2
1	A	31	CYS	2.1
1	A	678	LEU	2.1
1	A	677[A]	SER	2.1
1	A	796	THR	2.1
1	A	674	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	705	HIS	2.1
1	A	903	VAL	2.1
1	A	743	PRO	2.1
1	A	686	LEU	2.1
1	A	713	LEU	2.1
1	A	697	ILE	2.0
1	A	842	THR	2.0
1	A	712	PHE	2.0
1	A	698	GLN	2.0
1	A	253	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	KG8	A	1101	23/23	0.94	0.12	20,26,87,89	0
3	ZN	A	1103	1/1	0.94	0.06	37,37,37,37	0
3	ZN	A	1102	1/1	1.00	0.02	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.