



Full wwPDB NMR Structure Validation Report ⓘ

Mar 27, 2026 – 11:34 AM UTC

PDB ID : 2RSQ / pdb_00002rsq
BMRB ID : 11502
Title : Copper(I) loaded form of the first domain of the human copper chaperone for SOD1, CCS
Authors : Banci, L.; Bertini, I.; Cantini, F.; Kozyreva, T.; Rubino, J.T.
Deposited on : 2012-05-15

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 93%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	89	71% 7% • 20%

2 Ensemble composition and analysis

This entry contains 20 models. Model 13 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:10-A:78 (69)	0.43	13

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 4, 5, 6, 7, 8, 9, 10, 11, 13, 15, 17, 18, 19, 20
2	2, 16
3	3, 14
Single-model clusters	12

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1053 atoms, of which 529 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Copper chaperone for superoxide dismutase.

Mol	Chain	Residues	Atoms						Trace
1	A	71	Total	C	H	N	O	S	0
			1052	322	529	91	104	6	

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP O14618
A	-2	SER	-	expression tag	UNP O14618
A	-1	PHE	-	expression tag	UNP O14618
A	0	THR	-	expression tag	UNP O14618

- Molecule 2 is COPPER (I) ION (CCD ID: CU1) (formula: Cu).


Mol	Chain	Residues	Atoms	
2	A	1	Total	Cu
			1	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Copper chaperone for superoxide dismutase




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.2 Score per residue for model 2

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.3 Score per residue for model 3

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.4 Score per residue for model 4

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.5 Score per residue for model 5

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.6 Score per residue for model 6

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.7 Score per residue for model 7

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.8 Score per residue for model 8

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.9 Score per residue for model 9

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.10 Score per residue for model 10

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.11 Score per residue for model 11

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.12 Score per residue for model 12

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.13 Score per residue for model 13 (medoid)

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.14 Score per residue for model 14

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.15 Score per residue for model 15

- Molecule 1: Copper chaperone for superoxide dismutase




4.2.16 Score per residue for model 16

- Molecule 1: Copper chaperone for superoxide dismutase



4.2.17 Score per residue for model 17

- Molecule 1: Copper chaperone for superoxide dismutase



4.2.18 Score per residue for model 18

- Molecule 1: Copper chaperone for superoxide dismutase

Chain A:  71% 7% 20%



4.2.19 Score per residue for model 19

- Molecule 1: Copper chaperone for superoxide dismutase

Chain A:  62% 16% 20%



4.2.20 Score per residue for model 20

- Molecule 1: Copper chaperone for superoxide dismutase

Chain A:  67% 9% 20%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics, molecular dynamics*.

Of the 400 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	10
CYANA	structure solution	2.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	987
Number of shifts mapped to atoms	854
Number of unparsed shifts	0
Number of shifts with mapping errors	133
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	93%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.81±0.01	0±0/519 (0.0± 0.0%)	1.53±0.03	2±1/704 (0.2± 0.1%)
All	All	0.81	0/10380 (0.0%)	1.53	34/14080 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.2
All	All	0	1

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	25	CYS	CA-C-N	7.03	129.67	120.60	12	12
1	A	25	CYS	C-N-CA	7.03	129.67	120.60	12	12
1	A	16	PHE	CA-CB-CG	5.55	119.35	113.80	2	1
1	A	75	LEU	CD1-CG-CD2	-5.37	98.99	110.80	7	2
1	A	26	VAL	CA-CB-CG1	5.32	119.44	110.40	16	1
1	A	22	CYS	CA-C-N	5.24	127.30	120.28	2	1
1	A	22	CYS	C-N-CA	5.24	127.30	120.28	2	1
1	A	59	SER	CA-C-N	5.08	127.39	120.54	7	1
1	A	59	SER	C-N-CA	5.08	127.39	120.54	7	1
1	A	31	LYS	CA-CB-CG	-5.03	104.03	114.10	11	2

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	71	ARG	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	515	523	523	1±1
All	All	10320	10460	10460	14

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:44:VAL:HG22	1:A:51:VAL:HG13	0.50	1.82	14	3
1:A:18:VAL:HG21	1:A:20:MET:HE2	0.46	1.87	10	3
1:A:18:VAL:CG2	1:A:20:MET:HE2	0.43	2.44	16	3
1:A:20:MET:HE1	1:A:44:VAL:HG21	0.42	1.89	9	1
1:A:75:LEU:HD12	1:A:76:LYS:H	0.41	1.76	13	3
1:A:47:GLU:CD	1:A:47:GLU:H	0.40	2.24	15	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	69/89 (78%)	67±1 (98±1%)	2±1 (2±1%)	0±0 (0±0%)	49	83
All	All	1380/1780 (78%)	1346 (98%)	32 (2%)	2 (0%)	49	83

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	10	THR	2

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	59/73 (81%)	53±2 (90±3%)	6±2 (10±3%)	9	53
All	All	1180/1460 (81%)	1059 (90%)	121 (10%)	9	53

All 21 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	26	VAL	19
1	A	59	SER	17
1	A	11	LEU	16
1	A	51	VAL	12
1	A	39	VAL	11
1	A	48	ASP	9
1	A	22	CYS	8
1	A	46	LEU	8
1	A	72	GLN	4
1	A	67	GLU	3
1	A	27	ASP	2
1	A	52	LEU	2
1	A	44	VAL	2
1	A	50	MET	1
1	A	47	GLU	1
1	A	69	THR	1
1	A	49	GLN	1
1	A	36	VAL	1
1	A	23	GLN	1
1	A	29	VAL	1
1	A	41	ASP	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 93% for the well-defined parts and 93% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	987
Number of shifts mapped to atoms	854
Number of unparsed shifts	0
Number of shifts with mapping errors	133
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. All 133 occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	1	MET	H	8.268	0.020	1
1	A	1	MET	HA	4.376	0.020	1
1	A	1	MET	C	175.815	0.300	1
1	A	1	MET	CA	55.283	0.300	1
1	A	1	MET	CB	32.917	0.300	1
1	A	1	MET	N	123.044	0.20	1
1	A	2	ALA	H	8.301	0.020	1
1	A	2	ALA	HA	4.213	0.020	1
1	A	2	ALA	HB1	1.285	0.020	1
1	A	2	ALA	HB2	1.285	0.020	1
1	A	2	ALA	HB3	1.285	0.020	1
1	A	2	ALA	C	177.827	0.300	1
1	A	2	ALA	CA	52.53	0.300	1
1	A	2	ALA	CB	18.911	0.300	1

Continued on next page...

Continued from previous page...

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	2	ALA	N	125.557	0.20	1
1	A	3	SER	H	8.22	0.020	1
1	A	3	SER	HA	4.304	0.020	1
1	A	3	SER	HB2	3.735	0.020	.
1	A	3	SER	HB3	3.785	0.020	.
1	A	3	SER	C	174.42	0.300	1
1	A	3	SER	CA	58.25	0.300	1
1	A	3	SER	CB	63.648	0.300	1
1	A	3	SER	N	114.784	0.20	1
1	A	4	ASP	H	8.263	0.020	1
1	A	4	ASP	HA	4.561	0.020	1
1	A	4	ASP	HB2	2.598	0.020	.
1	A	4	ASP	C	176.511	0.300	1
1	A	4	ASP	CA	54.169	0.300	1
1	A	4	ASP	CB	41.143	0.300	1
1	A	4	ASP	N	122.181	0.20	1
1	A	5	SER	H	8.187	0.020	1
1	A	5	SER	HA	4.295	0.020	1
1	A	5	SER	HB2	3.813	0.020	.
1	A	5	SER	HB3	3.763	0.020	.
1	A	5	SER	C	175.298	0.300	1
1	A	5	SER	CA	58.686	0.300	1
1	A	5	SER	CB	63.718	0.300	1
1	A	5	SER	N	116.001	0.20	1
1	A	6	GLY	H	8.379	0.020	1
1	A	6	GLY	HA2	3.888	0.020	.
1	A	6	GLY	HA3	3.888	0.020	.
1	A	6	GLY	C	174.069	0.300	1
1	A	6	GLY	CA	45.093	0.300	1
1	A	6	GLY	N	110.345	0.20	1
1	A	7	ASN	H	8.171	0.020	1
1	A	7	ASN	HA	4.619	0.020	1
1	A	7	ASN	HB2	2.726	0.020	.
1	A	7	ASN	HB3	2.646	0.020	.
1	A	7	ASN	HD21	7.55	0.020	.
1	A	7	ASN	HD22	6.78	0.020	.
1	A	7	ASN	C	175.354	0.300	1
1	A	7	ASN	CA	53.171	0.300	1
1	A	7	ASN	CB	38.773	0.300	1
1	A	7	ASN	N	118.317	0.20	1
1	A	7	ASN	ND2	112.954	0.20	1

Continued on next page...

Continued from previous page...

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	8	GLN	H	8.373	0.020	1
1	A	8	GLN	HA	4.246	0.020	1
1	A	8	GLN	HB2	2.069	0.020	.
1	A	8	GLN	HB3	1.896	0.020	.
1	A	8	GLN	HE21	7.32	0.020	.
1	A	8	GLN	HE22	7.622	0.020	.
1	A	8	GLN	HG2	2.308	0.020	.
1	A	8	GLN	HG3	2.244	0.020	.
1	A	8	GLN	C	176.428	0.300	1
1	A	8	GLN	CA	55.746	0.300	1
1	A	8	GLN	CB	28.989	0.300	1
1	A	8	GLN	CG	33.843	0.300	1
1	A	8	GLN	N	120.394	0.20	1
1	A	8	GLN	NE2	112.913	0.20	1
1	A	80	SER	H	8.351	0.020	1
1	A	80	SER	HA	4.413	0.020	1
1	A	80	SER	HB2	3.823	0.020	.
1	A	80	SER	HB3	3.755	0.020	.
1	A	80	SER	C	175.257	0.300	1
1	A	80	SER	CA	58.36	0.300	1
1	A	80	SER	CB	63.949	0.300	1
1	A	80	SER	N	115.683	0.20	1
1	A	81	GLY	H	8.476	0.020	1
1	A	81	GLY	HA2	3.85	0.020	.
1	A	81	GLY	HA3	3.85	0.020	.
1	A	81	GLY	C	173.951	0.300	1
1	A	81	GLY	CA	45.245	0.300	1
1	A	81	GLY	N	110.929	0.20	1
1	A	82	GLN	H	8.137	0.020	1
1	A	82	GLN	HA	4.207	0.020	1
1	A	82	GLN	HB2	1.836	0.020	.
1	A	82	GLN	HB3	1.952	0.020	.
1	A	82	GLN	HE21	7.452	0.020	.
1	A	82	GLN	HG2	2.206	0.020	.
1	A	82	GLN	HG3	2.206	0.020	.
1	A	82	GLN	C	175.849	0.300	1
1	A	82	GLN	CA	55.513	0.300	1
1	A	82	GLN	CB	29.536	0.300	1
1	A	82	GLN	CG	33.623	0.300	1
1	A	82	GLN	N	119.529	0.20	1
1	A	82	GLN	NE2	112.301	0.20	1

Continued on next page...

Continued from previous page...

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	83	LEU	H	8.223	0.020	1
1	A	83	LEU	HA	4.218	0.020	1
1	A	83	LEU	HB2	1.525	0.020	.
1	A	83	LEU	HB3	1.458	0.020	.
1	A	83	LEU	HD11	0.793	0.020	.
1	A	83	LEU	HD12	0.793	0.020	.
1	A	83	LEU	HD13	0.793	0.020	.
1	A	83	LEU	HD21	0.745	0.020	.
1	A	83	LEU	HD22	0.745	0.020	.
1	A	83	LEU	HD23	0.745	0.020	.
1	A	83	LEU	HG	1.494	0.020	1
1	A	83	LEU	C	177.059	0.300	1
1	A	83	LEU	CA	55.092	0.300	1
1	A	83	LEU	CB	42.207	0.300	1
1	A	83	LEU	CD1	24.34	0.300	.
1	A	83	LEU	CD2	23.613	0.300	.
1	A	83	LEU	CG	26.949	0.300	1
1	A	83	LEU	N	123.317	0.20	1
1	A	84	GLN	H	8.273	0.020	1
1	A	84	GLN	HA	4.235	0.020	1
1	A	84	GLN	HB2	1.844	0.020	.
1	A	84	GLN	HB3	1.844	0.020	.
1	A	84	GLN	HG2	2.237	0.020	.
1	A	84	GLN	HG3	2.0	0.020	.
1	A	84	GLN	C	174.697	0.300	1
1	A	84	GLN	CA	55.511	0.300	1
1	A	84	GLN	CB	29.436	0.300	1
1	A	84	GLN	CG	33.389	0.300	1
1	A	84	GLN	N	121.306	0.20	1
1	A	85	ASN	H	7.967	0.020	1
1	A	85	ASN	HA	4.346	0.020	1
1	A	85	ASN	HB2	2.631	0.020	.
1	A	85	ASN	HB3	2.539	0.020	.
1	A	85	ASN	C	179.524	0.300	1
1	A	85	ASN	CA	54.609	0.300	1
1	A	85	ASN	CB	40.117	0.300	1
1	A	85	ASN	N	125.714	0.20	1

7.1.2 Chemical shift referencing ⓘ

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	85	-0.20 ± 0.07	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	76	0.45 ± 0.27	None needed (< 0.5 ppm)
$^{13}\text{C}'$	83	0.00 ± 0.17	None needed (< 0.5 ppm)
^{15}N	84	-0.37 ± 0.36	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 93%, i.e. 842 atoms were assigned a chemical shift out of a possible 910. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	346/348 (99%)	142/142 (100%)	136/138 (99%)	68/68 (100%)
Sidechain	490/538 (91%)	335/354 (95%)	148/168 (88%)	7/16 (44%)
Aromatic	6/24 (25%)	6/13 (46%)	0/9 (0%)	0/2 (0%)
Overall	842/910 (93%)	483/509 (95%)	284/315 (90%)	75/86 (87%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 93%, i.e. 854 atoms were assigned a chemical shift out of a possible 922. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	358/360 (99%)	148/148 (100%)	140/142 (99%)	70/70 (100%)
Sidechain	490/538 (91%)	335/354 (95%)	148/168 (88%)	7/16 (44%)
Aromatic	6/24 (25%)	6/13 (46%)	0/9 (0%)	0/2 (0%)
Overall	854/922 (93%)	489/515 (95%)	288/319 (90%)	77/88 (88%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

