



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:28 PM UTC

PDB ID : 7RTR / pdb_00007rtr
Title : YLQ-SG3 TCR in complex with SARS-CoV-2 Spike-derived peptide S269-277
(YLQPRTFLL) presented by HLA-A*02:01
Authors : Szeto, C.; Gras, S.
Deposited on : 2021-08-14
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	6% 61% 11% • 27%
2	B	100	3% 84% 16%
3	C	9	56% 44%
4	D	203	10% 78% 15% • •
5	E	241	3% 82% 16% • •

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6644 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	9	0
			2160	1363	392	396	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	6	0
			864	551	145	164	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			82	56	13	13			

- Molecule 4 is a protein called YLQ-SG3 TCR alpha chain (TRAV12-2).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	197	Total	C	N	O	S	0	0	0
			1513	947	251	307	8			

- Molecule 5 is a protein called YLQ-SG3 TCR beta chain (TRBV7-9).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	239	Total	C	N	O	S	0	10	0
			1974	1239	353	375	7			

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total 2	Na 2	0	0
6	D	2	Total 2	Na 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	22	Total 22	O 22	0	0
7	B	12	Total 12	O 12	0	0
7	C	1	Total 1	O 1	0	0
7	D	5	Total 5	O 5	0	0
7	E	7	Total 7	O 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class I antigen




- Molecule 2: Beta-2-microglobulin




- Molecule 3: Spike protein S1





- Molecule 4: YLQ-SG3 TCR alpha chain (TRAV12-2)





• Molecule 5: YLQ-SG3 TCR beta chain (TRBV7-9)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	225.36Å 49.62Å 91.72Å 90.00° 91.83° 90.00°	Depositor
Resolution (Å)	33.42 – 2.60 33.42 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.42-2.60) 99.9 (33.42-2.60)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.61Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.191 , 0.235 0.199 , 0.244	Depositor DCC
R_{free} test set	1507 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	41.2	Xtriage
Anisotropy	0.929	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 61.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.010 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6644	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	1/2244 (0.0%)	1.31	16/3043 (0.5%)
2	B	0.88	0/905	1.19	7/1224 (0.6%)
3	C	0.81	0/84	1.09	0/112
4	D	1.02	4/1546 (0.3%)	1.35	14/2100 (0.7%)
5	E	0.80	0/2027	1.19	4/2758 (0.1%)
All	All	0.87	5/6806 (0.1%)	1.27	41/9237 (0.4%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	7	ASN	CA-C	14.86	1.62	1.53
4	D	6	GLN	C-N	6.04	1.38	1.33
4	D	9	GLY	C-N	5.72	1.40	1.33
4	D	7	ASN	N-CA	5.41	1.49	1.46
1	A	45	MET	SD-CE	-5.36	1.66	1.79

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	16	GLY	CA-C-N	8.34	131.80	120.54
1	A	16	GLY	C-N-CA	8.34	131.80	120.54
4	D	7	ASN	CA-CB-CG	7.51	120.11	112.60
4	D	6	GLN	CA-C-N	7.28	130.91	123.04
4	D	6	GLN	C-N-CA	7.28	130.91	123.04
2	B	21	ASN	CA-CB-CG	6.79	119.39	112.60
1	A	59	TYR	CA-C-N	6.04	128.37	120.28
1	A	59	TYR	C-N-CA	6.04	128.37	120.28
1	A	18	GLY	N-CA-C	5.91	118.56	110.58
4	D	199	ASN	CA-CB-CG	5.81	118.41	112.60
4	D	191	ASN	CA-C-N	5.71	129.21	121.05

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	191	ASN	C-N-CA	5.71	129.21	121.05
5	E	26	ILE	CA-C-N	5.68	129.34	120.82
5	E	26	ILE	C-N-CA	5.68	129.34	120.82
4	D	194	ASP	CA-CB-CG	5.61	118.21	112.60
4	D	126	ILE	CA-C-N	5.57	128.78	120.31
4	D	126	ILE	C-N-CA	5.57	128.78	120.31
2	B	70	PHE	CA-CB-CG	5.56	119.36	113.80
1	A	163	THR	CA-C-N	5.50	127.59	120.44
1	A	163	THR	C-N-CA	5.50	127.59	120.44
1	A	137	ASP	CA-CB-CG	5.46	118.06	112.60
4	D	101	THR	CB-CA-C	5.42	118.69	109.80
1	A	155	GLN	CA-C-N	5.33	127.37	120.44
1	A	155	GLN	C-N-CA	5.33	127.37	120.44
4	D	152	PHE	CA-C-N	5.31	128.63	120.82
4	D	152	PHE	C-N-CA	5.31	128.63	120.82
1	A	156	LEU	CA-C-N	5.31	127.40	120.28
1	A	156	LEU	C-N-CA	5.31	127.40	120.28
5	E	18	ASN	CA-CB-CG	5.31	117.91	112.60
5	E	80	ARG	N-CA-C	-5.30	100.57	109.31
1	A	28	VAL	N-CA-CB	5.26	118.48	111.64
1	A	153	ALA	CA-C-N	5.24	127.25	120.44
1	A	153	ALA	C-N-CA	5.24	127.25	120.44
2	B	6	LYS	N-CA-C	-5.13	101.34	109.96
4	D	29	GLY	CA-C-N	5.13	128.36	120.82
4	D	29	GLY	C-N-CA	5.13	128.36	120.82
2	B	32	PRO	CA-C-N	5.11	128.07	120.31
2	B	32	PRO	C-N-CA	5.11	128.07	120.31
2	B	87	LEU	CA-C-N	5.08	127.09	120.28
2	B	87	LEU	C-N-CA	5.08	127.09	120.28
1	A	39	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2160	0	2003	10	0
2	B	864	0	846	4	0
3	C	82	0	88	0	0
4	D	1513	0	1398	9	0
5	E	1974	0	1845	17	0
6	A	2	0	0	0	0
6	D	2	0	0	0	0
7	A	22	0	0	0	0
7	B	12	0	0	0	0
7	C	1	0	0	0	0
7	D	5	0	0	0	0
7	E	7	0	0	0	0
All	All	6644	0	6180	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:101[A]:MET:HE2	5:E:117:GLY:HA3	1.61	0.83
2:B:4[A]:THR:HG22	2:B:86:THR:HB	1.72	0.72
5:E:37:ASN:HD22	5:E:58:ASN:HD21	1.38	0.72
5:E:38:ARG:HH11	5:E:57:GLN:HE21	1.36	0.72
4:D:163:ASP:HB2	4:D:166:VAL:HG22	1.75	0.68
4:D:105:ALA:HB1	4:D:112:ILE:CG2	2.24	0.66
1:A:89:GLU:H	1:A:89:GLU:CD	2.02	0.66
4:D:3:GLU:HB2	4:D:108:ARG:HD2	1.81	0.61
4:D:138:ASP:HB3	4:D:141:SER:O	2.05	0.57
5:E:139:PRO:HG3	5:E:150:ALA:HB1	1.85	0.57
1:A:97:ARG:HG3	1:A:116:TYR:CE2	2.41	0.56
4:D:105:ALA:HB1	4:D:112:ILE:HG21	1.90	0.53
1:A:126:LEU:HD22	1:A:156:LEU:HD13	1.93	0.51
5:E:101[B]:MET:HE2	5:E:103:LEU:HD21	1.94	0.50
5:E:53:LEU:HD22	5:E:72:LEU:HD13	1.94	0.49
1:A:14:ARG:HB3	1:A:17:ARG:HB2	1.96	0.48
1:A:97:ARG:HG3	1:A:116:TYR:CZ	2.48	0.48
1:A:20:PRO:HD2	1:A:75:ARG:HH11	1.78	0.48
5:E:21:PHE:CD1	5:E:118:THR:HG21	2.49	0.47
4:D:108:ARG:O	4:D:111:LYS:HG2	2.14	0.47
5:E:132:PRO:HB3	5:E:159:PHE:CD2	2.50	0.47
2:B:25:CYS:HB2	2:B:39:LEU:HD21	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:136:LEU:HD22	5:E:139:PRO:HA	1.99	0.45
5:E:12:ILE:HD13	5:E:161:PRO:HG3	1.99	0.45
5:E:76:PHE:CE2	5:E:91:ILE:HG12	2.52	0.45
2:B:2:GLN:HB3	2:B:86:THR:HG22	2.00	0.44
5:E:121:THR:HG21	5:E:161:PRO:HB3	2.00	0.43
5:E:207:ALA:O	5:E:211:GLN:HG3	2.18	0.43
2:B:2:GLN:HB3	2:B:86:THR:CG2	2.49	0.43
5:E:13[A]:THR:HG21	5:E:19:VAL:HG13	1.99	0.43
5:E:153:VAL:HG22	5:E:202:ARG:HG3	2.00	0.43
1:A:95:VAL:HG12	1:A:118:TYR:HD1	1.84	0.42
4:D:166:VAL:HG12	4:D:190:SER:HB2	2.01	0.42
1:A:133:TRP:HB2	1:A:144:LYS:HD2	2.01	0.41
1:A:6[A]:ARG:HG2	1:A:113[A]:TYR:HE1	1.85	0.41
1:A:187:THR:HA	1:A:204:TRP:O	2.21	0.41
4:D:174:LEU:HB3	5:E:180:CYS:HB3	2.03	0.41
5:E:58:ASN:O	5:E:80:ARG:HD3	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/365 (73%)	254 (96%)	10 (4%)	1 (0%)	30	51
2	B	104/100 (104%)	100 (96%)	4 (4%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
4	D	195/203 (96%)	185 (95%)	9 (5%)	1 (0%)	24	46
5	E	247/241 (102%)	233 (94%)	13 (5%)	1 (0%)	30	51
All	All	818/918 (89%)	779 (95%)	36 (4%)	3 (0%)	30	51

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	161	SER
5	E	234	GLN
1	A	252	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/299 (72%)	191 (89%)	23 (11%)	6	14
2	B	101/95 (106%)	94 (93%)	7 (7%)	14	32
3	C	9/9 (100%)	5 (56%)	4 (44%)	0	0
4	D	167/183 (91%)	145 (87%)	22 (13%)	4	8
5	E	211/213 (99%)	197 (93%)	14 (7%)	15	34
All	All	702/799 (88%)	632 (90%)	70 (10%)	7	16

All (70) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	SER
1	A	14	ARG
1	A	35	ARG
1	A	73	THR
1	A	75	ARG
1	A	89	GLU
1	A	97	ARG
1	A	110	LEU
1	A	141[A]	GLN
1	A	141[B]	GLN
1	A	156	LEU
1	A	176	LYS
1	A	181	ARG
1	A	189	MET
1	A	200	THR
1	A	213	ILE
1	A	216	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	234	ARG
1	A	251	SER
1	A	254	GLU
1	A	255	GLN
1	A	264	GLU
1	A	271	THR
2	B	1	ILE
2	B	19	LYS
2	B	58	LYS
2	B	68	THR
2	B	88	SER
2	B	89[A]	GLN
2	B	89[B]	GLN
3	C	2	LEU
3	C	5	ARG
3	C	8	LEU
3	C	9	LEU
4	D	3	GLU
4	D	6	GLN
4	D	7	ASN
4	D	11	LEU
4	D	38	SER
4	D	67	GLU
4	D	82	LYS
4	D	87	VAL
4	D	90	LEU
4	D	92	ARG
4	D	95	GLN
4	D	99	SER
4	D	101	THR
4	D	108	ARG
4	D	119	ARG
4	D	127	GLN
4	D	145	SER
4	D	148	LEU
4	D	153	ASP
4	D	161	SER
4	D	163	ASP
4	D	209	ASP
5	E	12	ILE
5	E	15	ARG
5	E	19	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	E	22	ARG
5	E	38	ARG
5	E	63	GLU
5	E	67	GLU
5	E	83	LYS
5	E	92	GLN
5	E	165	GLU
5	E	189	GLN
5	E	192	LEU
5	E	231	GLU
5	E	251	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	115	GLN
1	A	145	HIS
3	C	3	GLN
4	D	37	GLN
4	D	95	GLN
5	E	10	HIS
5	E	18	ASN
5	E	57	GLN
5	E	58	ASN
5	E	112	GLN
5	E	146	HIS
5	E	193	ASN
5	E	220	GLN
5	E	222	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	266/365 (72%)	0.27	23 (8%) 16 12	21, 44, 98, 115	11 (4%)
2	B	100/100 (100%)	-0.09	3 (3%) 52 47	19, 40, 71, 81	6 (6%)
3	C	9/9 (100%)	-0.71	0 100 100	31, 32, 35, 37	0
4	D	197/203 (97%)	0.55	20 (10%) 12 9	15, 48, 96, 109	4 (2%)
5	E	239/241 (99%)	0.27	7 (2%) 53 48	21, 53, 88, 101	10 (4%)
All	All	811/918 (88%)	0.28	53 (6%) 25 19	15, 47, 91, 115	31 (3%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	141	SER	11.4
4	D	203	ASN	10.2
4	D	195	PHE	9.5
1	A	193	ALA	7.8
1	A	195	SER	7.3
1	A	218	GLN	5.4
4	D	143	ASP	4.5
4	D	214	SER	4.1
4	D	9	GLY	4.0
1	A	256	ARG	3.8
4	D	161	SER	3.6
1	A	258	THR	3.4
5	E	228	GLU	3.3
4	D	144	LYS	3.2
1	A	248	VAL	3.1
4	D	140	LYS	3.1
1	A	251	SER	3.1
5	E	253	ASP	3.0
4	D	142	SER	3.0
4	D	191	ASN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	0	MET	2.9
4	D	10	PRO	2.9
1	A	198	GLU	2.8
5	E	191	ALA	2.8
1	A	196	ASP	2.7
1	A	226	GLN	2.7
4	D	3	GLU	2.7
1	A	225	THR	2.6
4	D	192	LYS	2.6
1	A	224	GLN	2.6
4	D	205	ILE	2.6
1	A	254	GLU	2.6
4	D	4	VAL	2.6
1	A	259	CYS	2.5
1	A	223	ASP	2.5
2	B	80	CYS	2.4
2	B	75	LYS	2.4
1	A	270	LEU	2.4
1	A	274	TRP	2.4
5	E	193	ASN	2.2
4	D	160	GLN	2.2
1	A	194	VAL	2.2
1	A	201	LEU	2.2
4	D	153	ASP	2.2
1	A	199	ALA	2.2
1	A	271	THR	2.2
5	E	235	ASP	2.2
5	E	79[A]	GLU	2.1
4	D	197	CYS	2.1
1	A	255	GLN	2.0
4	D	162	LYS	2.0
5	E	192	LEU	2.0
1	A	197	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NA	A	402	1/1	0.85	0.32	53,53,53,53	0
6	NA	A	401	1/1	0.94	0.09	48,48,48,48	0
6	NA	D	301	1/1	0.95	0.15	46,46,46,46	0
6	NA	D	302	1/1	0.95	0.20	57,57,57,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.