



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 02:47 AM UTC

PDB ID : 3RU8 / pdb_00003ru8
Title : Structure of an HIV epitope scaffold in complex with neutralizing antibody b12 Fab
Authors : Carrico, C.T.D.; Strong, R.K.
Deposited on : 2011-05-04
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	282	5% 94%
2	H	230	11% 90% 7%
3	L	215	6% 93% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5881 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epitope Scaffold 2bodx43.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	277	Total	C	N	O	S	0	14	0
			2120	1326	375	410	9			

- Molecule 2 is a protein called Antibody b12, Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	225	Total	C	N	O	S	0	11	0
			1765	1121	299	337	8			

- Molecule 3 is a protein called Antibody b12, Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	213	Total	C	N	O	S	0	2	0
			1627	1014	286	323	4			

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	X	1	Total	C	O	0	0
			6	3	3		
4	X	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	X	169	Total	O	0	2
			171	171		
5	H	99	Total	O	0	0
			99	99		
5	L	75	Total	O	0	0
			75	75		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epitope Scaffold 2bodx43



- Molecule 2: Antibody b12, Heavy Chain



- Molecule 3: Antibody b12, Light Chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.45Å 93.10Å 94.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.23 – 2.07 45.23 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.2 (45.23-2.07) 99.3 (45.23-2.07)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.07Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.168 , 0.203 (Not available) , 0.221	Depositor DCC
R_{free} test set	2483 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.013 for -h,l,k 0.013 for -l,-k,-h 0.015 for k,h,-l 0.000 for k,l,h 0.000 for l,h,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5881	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	0.60	1/2214 (0.0%)	0.73	0/3022
2	H	0.52	0/1845	0.66	0/2514
3	L	0.52	0/1669	0.67	0/2267
All	All	0.55	1/5728 (0.0%)	0.69	0/7803

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	221	ASP	C-O	-5.14	1.21	1.23

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	2120	0	2057	7	0
2	H	1765	0	1729	12	0
3	L	1627	0	1551	8	0
4	L	12	0	16	1	0
4	X	12	0	16	2	0
5	H	99	0	0	0	0
5	L	75	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	X	171	0	0	0	0
All	All	5881	0	5369	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:100(J):MET:HE3	3:L:90:GLN:HE22	1.38	0.88
1:X:156[B]:MET:HE1	4:X:283:GOL:H31	1.75	0.68
3:L:48:VAL:HG12	3:L:49:ILE:HG12	1.85	0.57
2:H:17[A]:SER:OG	2:H:82(A)[A]:ARG:HG2	2.07	0.54
2:H:98:TYR:CE2	2:H:100(B):ASP:HB2	2.45	0.52
2:H:187:LEU:C	2:H:187:LEU:HD12	2.35	0.52
3:L:152:ASP:HA	3:L:192:VAL:HG12	1.93	0.50
3:L:121:PRO:HD3	3:L:133:VAL:HG22	1.93	0.50
1:X:251:GLU:OE1	4:X:284:GOL:H32	2.12	0.50
2:H:80[B]:MET:HE1	2:H:90:TYR:HB3	1.96	0.47
3:L:152:ASP:HA	3:L:192:VAL:CG1	2.44	0.47
2:H:100(J):MET:HG3	2:H:103:TRP:CZ2	2.50	0.47
2:H:127:SER:O	2:H:133:THR:HG23	2.15	0.47
1:X:156[A]:MET:HG3	1:X:188:TYR:OH	2.14	0.46
2:H:100(J):MET:HE2	2:H:100(J):MET:N	2.30	0.46
2:H:100(I):TYR:C	2:H:100(J):MET:HE2	2.41	0.46
3:L:179:THR:OG1	4:L:217:GOL:H12	2.17	0.45
2:H:11:VAL:HG21	2:H:149:PRO:HG3	1.98	0.45
1:X:145:ALA:C	1:X:156[A]:MET:HE2	2.41	0.45
1:X:99:ASN:HA	1:X:136:SER:HA	2.00	0.43
2:H:24:ALA:HB1	2:H:27:TYR:CE2	2.53	0.43
1:X:231:SER:HA	1:X:243:PHE:O	2.19	0.42
1:X:208:SER:HA	1:X:245:TRP:CD1	2.54	0.42
2:H:205:THR:HG23	2:H:222:LYS:HE3	2.01	0.42
3:L:192:VAL:HG13	3:L:192:VAL:O	2.21	0.41
3:L:8:PRO:HG3	3:L:11:LEU:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	289/282 (102%)	283 (98%)	6 (2%)	0	100	100
2	H	232/230 (101%)	228 (98%)	4 (2%)	0	100	100
3	L	213/215 (99%)	209 (98%)	4 (2%)	0	100	100
All	All	734/727 (101%)	720 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	219/214 (102%)	218 (100%)	1 (0%)	81	85
2	H	199/196 (102%)	198 (100%)	1 (0%)	81	85
3	L	178/187 (95%)	176 (99%)	2 (1%)	65	68
All	All	596/597 (100%)	592 (99%)	4 (1%)	76	79

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	79	MET
2	H	187	LEU
3	L	106	GLU
3	L	180	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	21	ASN
1	X	167	ASN
2	H	39	GLN
2	H	64	GLN
3	L	38	GLN
3	L	43	GLN
3	L	90	GLN
3	L	148	GLN
3	L	161	GLN
3	L	200	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	L	217	-	5,5,5	0.34	0	5,5,5	0.51	0
4	GOL	X	283	-	5,5,5	0.36	0	5,5,5	0.46	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	X	284	-	5,5,5	0.37	0	5,5,5	0.46	0
4	GOL	L	216	-	5,5,5	0.45	0	5,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	L	217	-	-	2/4/4/4	-
4	GOL	X	283	-	-	2/4/4/4	-
4	GOL	X	284	-	-	2/4/4/4	-
4	GOL	L	216	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	X	283	GOL	C1-C2-C3-O3
4	X	284	GOL	C1-C2-C3-O3
4	L	216	GOL	C1-C2-C3-O3
4	L	217	GOL	O1-C1-C2-O2
4	L	217	GOL	O1-C1-C2-C3
4	X	284	GOL	O2-C2-C3-O3
4	X	283	GOL	O2-C2-C3-O3
4	L	216	GOL	O2-C2-C3-O3
4	L	216	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	217	GOL	1	0
4	X	283	GOL	1	0
4	X	284	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	X	277/282 (98%)	0.37	13 (4%)	36 36	18, 35, 53, 72	14 (5%)
2	H	225/230 (97%)	0.77	26 (11%)	9 9	21, 40, 62, 106	11 (4%)
3	L	213/215 (99%)	0.63	13 (6%)	27 27	21, 42, 70, 87	2 (0%)
All	All	715/727 (98%)	0.57	52 (7%)	21 21	18, 38, 66, 106	27 (3%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	125	ALA	6.9
2	H	124	LEU	6.9
2	H	122	PHE	5.6
2	H	121	VAL	5.1
2	H	123	PRO	4.8
2	H	126	PRO	4.2
2	H	133	THR	3.7
3	L	107[A]	ARG	3.7
2	H	100	TRP	3.6
1	X	277	HIS	3.5
2	H	118	GLY	3.5
3	L	213	GLY	3.4
1	X	79	MET	3.4
2	H	119	PRO	3.4
1	X	82	GLY	3.4
2	H	128	SER	3.3
1	X	81	GLN	3.3
3	L	155	LEU	3.3
2	H	100(A)	ASP	3.3
2	H	120	SER	3.2
1	X	275	LEU	3.1
3	L	205	PRO	3.1
2	H	65	ASP	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	76	THR	3.0
3	L	189	LYS	3.0
1	X	80	SER	2.9
3	L	153	ASN	2.8
1	X	44	HIS	2.7
1	X	46	PRO	2.6
2	H	198	LEU	2.6
3	L	203	ARG	2.6
2	H	199	GLY	2.5
1	X	45	ASN	2.5
2	H	127	SER	2.4
2	H	116	THR	2.4
2	H	1	GLN	2.3
1	X	1	ASP	2.3
2	H	168	SER	2.3
1	X	217	ASN	2.3
2	H	117	LYS	2.3
1	X	83	GLU	2.2
3	L	195	CYS	2.2
3	L	209	SER	2.2
2	H	165	ALA	2.2
2	H	142	CYS	2.1
3	L	204	SER	2.1
3	L	154	ALA	2.1
2	H	115	SER	2.1
3	L	198	THR	2.1
3	L	118	ILE	2.1
2	H	216	ASN	2.0
2	H	82(A)[A]	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	L	217	6/6	0.77	0.17	52,55,57,60	0
4	GOL	X	283	6/6	0.87	0.16	51,52,55,58	0
4	GOL	X	284	6/6	0.88	0.14	49,51,52,54	0
4	GOL	L	216	6/6	0.89	0.17	38,45,46,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.