



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 04:29 PM UTC

PDB ID : 6RV6 / pdb_00006rv6
Title : Structure of properdin lacking TSR3 based on anomalous data
Authors : Pedersen, D.V.; Andersen, G.R.
Deposited on : 2019-05-31
Resolution : 3.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1085 (3.54-3.46)
Clashscore	190562	1140 (3.54-3.46)
Ramachandran outliers	187476	1113 (3.54-3.46)
Sidechain outliers	187428	1114 (3.54-3.46)
RSRZ outliers	180081	1084 (3.54-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	
2	B	221	
3	C	2	
3	D	2	
3	E	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	3	67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FUC	F	3	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3191 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Properdin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	165	Total	C	N	O	S	0	0	0
			1244	756	229	238	21			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	192	GLU	-	expression tag	UNP P27918
A	193	ASN	-	expression tag	UNP P27918
A	194	LEU	-	expression tag	UNP P27918
A	195	TYR	-	expression tag	UNP P27918
A	196	PHE	-	expression tag	UNP P27918
A	197	GLN	-	expression tag	UNP P27918


- Molecule 2 is a protein called Properdin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	220	Total	C	N	O	S	0	0	0
			1714	1063	319	310	22			

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	255	GLY	-	expression tag	UNP P27918
B	470	GLU	-	expression tag	UNP P27918
B	471	ASN	-	expression tag	UNP P27918
B	472	LEU	-	expression tag	UNP P27918
B	473	TYR	-	expression tag	UNP P27918
B	474	PHE	-	expression tag	UNP P27918
B	475	GLN	-	expression tag	UNP P27918

- Molecule 3 is an oligosaccharide called beta-D-glucopyranose-(1-3)-alpha-L-fucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	2	Total	C	O	0	0	0
			21	12	9			
3	D	2	Total	C	O	0	0	0
			21	12	9			
3	E	2	Total	C	O	0	0	0
			21	12	9			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 5 is alpha-D-mannopyranose (CCD ID: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 1: Properdin



- | |
|------|
| G265 | A256 | W260 | P265 | V266 | M279 | E280 | T283 | P287 | G293 | P294 | F295 | C296 | A297 | T308 | A309 | V310 | P311 | C312 | P313 | V314 | I328 | C337 | R359 | P394 | L411 | M420 | V421 | E422 | T430 | L442 | V449 | L472 | V473 | F474 | G475 |
|------|

- FUC1
BGC2

- FUC1
BGC2

- FUC1
BGC2

- 

Chain F:

67%

33%



4 Data and refinement statistics

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, α , β , γ	219.79Å 219.79Å 47.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	77.71 – 3.51 77.71 – 3.51	Depositor EDS
% Data completeness (in resolution range)	99.8 (77.71-3.51) 99.9 (77.71-3.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 3.49Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.264 , 0.283 (Not available) , 0.262	Depositor DCC
R_{free} test set	717 reflections (2.57%)	wwPDB-VP
Wilson B-factor (Å ²)	134.3	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 96.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.012 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3191	wwPDB-VP
Average B, all atoms (Å ²)	149.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, NAG, MAN, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/1278	0.75	0/1734
2	B	0.29	0/1770	0.74	2/2410 (0.1%)
All	All	0.29	0/3048	0.75	2/4144 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	394	PRO	CA-C-N	5.29	131.63	121.54
2	B	394	PRO	C-N-CA	5.29	131.63	121.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1244	0	1135	7	0
2	B	1714	0	1613	5	0
3	C	21	0	19	0	0
3	D	21	0	19	0	0
3	E	21	0	19	0	0
4	F	38	0	34	0	0
5	A	55	0	50	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	77	0	70	0	0
All	All	3191	0	2959	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 12 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:TYR:HB2	1:A:63:ALA:HB2	1.89	0.54
1:A:97:SER:HB3	1:A:124:LEU:HD11	1.88	0.54
1:A:159:ARG:NH1	1:A:180:GLU:OE1	2.41	0.53
2:B:257:ALA:HA	2:B:294:PRO:HG2	1.93	0.50
1:A:102:ARG:NH2	1:A:121:GLU:OE1	2.44	0.47

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/170 (96%)	147 (90%)	14 (9%)	2 (1%)	10	41
2	B	218/221 (99%)	204 (94%)	12 (6%)	2 (1%)	14	47
All	All	381/391 (97%)	351 (92%)	26 (7%)	4 (1%)	12	44

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	ARG
1	A	191	PRO
2	B	293	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	287	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/141 (96%)	128 (94%)	8 (6%)	18	44
2	B	189/190 (100%)	175 (93%)	14 (7%)	13	37
All	All	325/331 (98%)	303 (93%)	22 (7%)	14	40

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	314	VAL
2	B	359	ARG
2	B	337	CYS
2	B	411	LEU
1	A	170	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	FUC	C	1	1,3	10,10,11	0.89	0	14,14,16	1.04	1 (7%)
3	BGC	C	2	3	11,11,12	1.64	3 (27%)	15,15,17	1.02	1 (6%)
3	FUC	D	1	1,3	10,10,11	1.10	1 (10%)	14,14,16	1.03	1 (7%)
3	BGC	D	2	3	11,11,12	1.71	3 (27%)	15,15,17	1.02	1 (6%)
3	FUC	E	1	2,3	10,10,11	0.81	0	14,14,16	0.86	1 (7%)
3	BGC	E	2	3	11,11,12	1.70	3 (27%)	15,15,17	1.04	1 (6%)
4	NAG	F	1	2,4	14,14,15	0.44	0	17,19,21	0.64	0
4	NAG	F	2	4	14,14,15	0.61	1 (7%)	17,19,21	0.95	1 (5%)
4	FUC	F	3	4	10,10,11	1.20	0	14,14,16	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FUC	C	1	1,3	-	-	0/1/1/1
3	BGC	C	2	3	-	2/2/19/22	0/1/1/1
3	FUC	D	1	1,3	-	-	0/1/1/1
3	BGC	D	2	3	-	2/2/19/22	0/1/1/1
3	FUC	E	1	2,3	-	-	0/1/1/1
3	BGC	E	2	3	-	1/2/19/22	0/1/1/1
4	NAG	F	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	F	2	4	-	4/6/23/26	0/1/1/1
4	FUC	F	3	4	1/1/4/5	-	0/1/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	BGC	O5-C1	4.54	1.51	1.43
3	E	2	BGC	O5-C1	4.43	1.51	1.43
3	C	2	BGC	O5-C1	4.24	1.50	1.43
3	D	1	FUC	C2-C3	2.25	1.55	1.52
3	E	2	BGC	O5-C5	2.24	1.47	1.43

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	2	NAG	C2-N2-C7	2.81	126.67	122.90
3	C	2	BGC	C1-C2-C3	2.42	113.17	109.64
3	E	2	BGC	C3-C4-C5	2.14	114.11	110.23
3	D	2	BGC	C2-C3-C4	2.13	114.60	110.86
3	D	1	FUC	C1-C2-C3	2.09	112.69	109.64

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	F	3	FUC	C1

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	F	1	NAG	O5-C5-C6-O6
4	F	2	NAG	O5-C5-C6-O6
3	C	2	BGC	O5-C5-C6-O6
4	F	1	NAG	C4-C5-C6-O6
4	F	2	NAG	C4-C5-C6-O6

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.


Oligosaccharide Chain C



Oligosaccharide Chain D







5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MAN	B	512	2	11,11,12	0.90	0	15,15,17	1.85	1 (6%)
5	MAN	B	509	2	11,11,12	0.94	0	15,15,17	1.70	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MAN	B	510	2	11,11,12	0.80	0	15,15,17	1.78	3 (20%)
5	MAN	B	511	2	11,11,12	1.00	0	15,15,17	1.71	2 (13%)
5	MAN	A	208	1	11,11,12	1.13	0	15,15,17	1.69	2 (13%)
5	MAN	A	204	1	11,11,12	0.99	0	15,15,17	1.96	2 (13%)
5	MAN	A	203	1	11,11,12	0.83	0	15,15,17	1.61	2 (13%)
5	MAN	B	505	2	11,11,12	1.07	0	15,15,17	1.84	4 (26%)
5	MAN	B	503	2	11,11,12	1.01	0	15,15,17	1.51	1 (6%)
5	MAN	A	207	1	11,11,12	1.06	0	15,15,17	1.59	3 (20%)
5	MAN	A	209	1	11,11,12	1.05	0	15,15,17	1.59	1 (6%)
5	MAN	B	504	2	11,11,12	0.94	0	15,15,17	1.57	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MAN	B	512	2	-	0/2/19/22	0/1/1/1
5	MAN	B	509	2	-	0/2/19/22	0/1/1/1
5	MAN	B	510	2	-	0/2/19/22	0/1/1/1
5	MAN	B	511	2	-	0/2/19/22	0/1/1/1
5	MAN	A	208	1	-	0/2/19/22	0/1/1/1
5	MAN	A	204	1	-	0/2/19/22	0/1/1/1
5	MAN	A	203	1	-	0/2/19/22	0/1/1/1
5	MAN	B	505	2	-	0/2/19/22	0/1/1/1
5	MAN	B	503	2	-	0/2/19/22	0/1/1/1
5	MAN	A	207	1	-	0/2/19/22	0/1/1/1
5	MAN	A	209	1	-	0/2/19/22	0/1/1/1
5	MAN	B	504	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	512	MAN	C1-O5-C5	6.06	120.31	112.19
5	A	204	MAN	C1-O5-C5	5.90	120.09	112.19
5	B	510	MAN	C1-O5-C5	5.29	119.27	112.19
5	B	505	MAN	C1-O5-C5	5.12	119.05	112.19
5	A	208	MAN	C1-O5-C5	5.03	118.93	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	165/170 (97%)	0.12	2 (1%) 76 52	100, 137, 203, 209	0
2	B	220/221 (99%)	0.18	2 (0%) 81 58	99, 137, 216, 241	0
All	All	385/391 (98%)	0.16	4 (1%) 79 56	99, 137, 208, 241	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	309	ALA	3.9
1	A	113	GLY	2.8
2	B	265	PRO	2.6
1	A	39	SER	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BGC	E	2	11/12	0.51	0.17	136,150,159,162	0
4	NAG	F	2	14/15	0.64	0.10	205,243,253,254	0
4	NAG	F	1	14/15	0.70	0.12	199,220,229,232	0
3	BGC	C	2	11/12	0.73	0.11	136,143,146,147	0
3	BGC	D	2	11/12	0.75	0.07	149,157,164,167	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FUC	E	1	10/11	0.83	0.15	132,146,151,158	0
4	FUC	F	3	10/11	0.83	0.13	145,157,184,186	0
3	FUC	D	1	10/11	0.84	0.09	154,160,167,171	0
3	FUC	C	1	10/11	0.97	0.09	133,140,143,144	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain D:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain E:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MAN	A	209	11/12	0.60	0.10	182,189,193,194	0
5	MAN	B	511	11/12	0.67	0.10	200,208,214,221	0
5	MAN	A	204	11/12	0.72	0.08	150,158,162,164	0
5	MAN	B	512	11/12	0.75	0.09	187,197,208,209	0
5	MAN	A	208	11/12	0.76	0.07	182,189,197,200	0
5	MAN	B	505	11/12	0.76	0.10	138,145,154,156	0
5	MAN	B	510	11/12	0.79	0.09	151,159,169,170	0
5	MAN	B	503	11/12	0.79	0.09	119,131,136,136	0
5	MAN	A	203	11/12	0.79	0.12	141,149,158,159	0
5	MAN	B	504	11/12	0.86	0.11	129,134,145,146	0
5	MAN	A	207	11/12	0.86	0.06	182,188,196,197	0
5	MAN	B	509	11/12	0.86	0.08	128,136,144,146	0

6.5 Other polymers [i](#)

There are no such residues in this entry.