



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:24 PM UTC

PDB ID : 6RV9 / pdb_00006rv9
Title : Crystal Structure of Glucuronoyl Esterase from *Cerrena unicolor* inactive S270A variant in complex with the aldouronic acid XUXXR
Authors : Ernst, H.A.; Mosbech, C.; Langkilde, A.; Westh, P.; Meyer, A.; Agger, J.W.; Larsen, S.
Deposited on : 2019-05-31
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	2% 91% 5%
1	B	401	17% 84% 11% 5%
2	C	5	80% 20%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6398 atoms, of which 51 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-O-methyl-glucuronoyl methylesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	380	Total	C	N	O	S	0	2	0
			2872	1814	492	551	15			
1	B	380	Total	C	N	O	S	0	1	0
			2866	1810	491	550	15			

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	73	GLU	-	expression tag	UNP A0A0A7EQR3
A	74	ALA	-	expression tag	UNP A0A0A7EQR3
A	75	GLU	-	expression tag	UNP A0A0A7EQR3
A	76	ALA	-	expression tag	UNP A0A0A7EQR3
A	77	GLU	-	expression tag	UNP A0A0A7EQR3
A	78	PHE	-	expression tag	UNP A0A0A7EQR3
A	270	ALA	SER	engineered mutation	UNP A0A0A7EQR3
A	459	GLU	-	expression tag	UNP A0A0A7EQR3
A	460	ASN	-	expression tag	UNP A0A0A7EQR3
A	461	LEU	-	expression tag	UNP A0A0A7EQR3
A	462	TYR	-	expression tag	UNP A0A0A7EQR3
A	463	PHE	-	expression tag	UNP A0A0A7EQR3
A	464	GLN	-	expression tag	UNP A0A0A7EQR3
A	465	GLY	-	expression tag	UNP A0A0A7EQR3
A	466	VAL	-	expression tag	UNP A0A0A7EQR3
A	467	ASP	-	expression tag	UNP A0A0A7EQR3
A	468	HIS	-	expression tag	UNP A0A0A7EQR3
A	469	HIS	-	expression tag	UNP A0A0A7EQR3
A	470	HIS	-	expression tag	UNP A0A0A7EQR3
A	471	HIS	-	expression tag	UNP A0A0A7EQR3
A	472	HIS	-	expression tag	UNP A0A0A7EQR3
A	473	HIS	-	expression tag	UNP A0A0A7EQR3
B	73	GLU	-	expression tag	UNP A0A0A7EQR3
B	74	ALA	-	expression tag	UNP A0A0A7EQR3
B	75	GLU	-	expression tag	UNP A0A0A7EQR3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	76	ALA	-	expression tag	UNP A0A0A7EQR3
B	77	GLU	-	expression tag	UNP A0A0A7EQR3
B	78	PHE	-	expression tag	UNP A0A0A7EQR3
B	270	ALA	SER	engineered mutation	UNP A0A0A7EQR3
B	459	GLU	-	expression tag	UNP A0A0A7EQR3
B	460	ASN	-	expression tag	UNP A0A0A7EQR3
B	461	LEU	-	expression tag	UNP A0A0A7EQR3
B	462	TYR	-	expression tag	UNP A0A0A7EQR3
B	463	PHE	-	expression tag	UNP A0A0A7EQR3
B	464	GLN	-	expression tag	UNP A0A0A7EQR3
B	465	GLY	-	expression tag	UNP A0A0A7EQR3
B	466	VAL	-	expression tag	UNP A0A0A7EQR3
B	467	ASP	-	expression tag	UNP A0A0A7EQR3
B	468	HIS	-	expression tag	UNP A0A0A7EQR3
B	469	HIS	-	expression tag	UNP A0A0A7EQR3
B	470	HIS	-	expression tag	UNP A0A0A7EQR3
B	471	HIS	-	expression tag	UNP A0A0A7EQR3
B	472	HIS	-	expression tag	UNP A0A0A7EQR3
B	473	HIS	-	expression tag	UNP A0A0A7EQR3

- Molecule 2 is an oligosaccharide called 4-O-methyl-alpha-D-glucopyranuronic acid-(1-2)-[beta-D-xylopyranose-(1-4)]beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-Xylitol.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	5	Total	C	H	O	0	0	0
			95	27	45	23			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	302	Total 302	O 302	0	0
5	B	225	Total 225	O 225	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 4-O-methyl-glucuronoyl methylesterase



- Molecule 1: 4-O-methyl-glucuronoyl methylesterase



- Molecule 2: 4-O-methyl-alpha-D-glucopyranuronic acid-(1-2)-[beta-D-xylopyranose-(1-4)]beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-Xylitol



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	84.48Å 84.48Å 261.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.50 – 1.64 44.50 – 1.64	Depositor EDS
% Data completeness (in resolution range)	99.5 (44.50-1.64) 99.5 (44.50-1.64)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 1.64Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.186 , 0.210 0.186 , 0.211	Depositor DCC
R_{free} test set	5754 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6398	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XYL, EDO, XYP, GCV, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/2956	0.49	0/4045
1	B	0.24	0/2947	0.45	0/4033
All	All	0.26	0/5903	0.47	0/8078

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2872	0	2760	9	0
1	B	2866	0	2752	25	0
2	C	50	45	20	1	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
4	A	4	6	6	0	0
5	A	302	0	0	1	0
5	B	225	0	0	1	0
All	All	6347	51	5564	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:174:PRO:HG2	1:B:183:TRP:CE3	2.36	0.60
1:B:252:GLU:HB3	5:B:627:HOH:O	2.01	0.60
1:B:270:ALA:HB1	2:C:4:GCV:C6	2.32	0.59
1:B:350:ALA:HB1	1:B:358[A]:MET:HE1	1.85	0.58
1:A:378:MET:HA	1:A:378:MET:HE2	1.91	0.52
1:B:142:ALA:HA	1:B:154:ALA:O	2.10	0.52
1:A:429:THR:HG21	1:B:338:LYS:NZ	2.24	0.51
1:B:138:PRO:HD2	1:B:159:LEU:HA	1.92	0.51
1:B:179:PRO:HG2	1:B:182:GLY:O	2.10	0.51
1:B:186:ILE:HD12	1:B:196:ILE:HD12	1.93	0.50
1:B:190:GLU:HG3	1:B:207:ASN:OD1	2.13	0.48
1:B:153:LEU:O	1:B:168:SER:HA	2.13	0.48
1:B:144:PHE:CZ	1:B:151:GLY:HA3	2.49	0.47
1:B:248:ILE:HG21	1:B:285:ARG:HD2	1.95	0.47
1:B:153:LEU:O	1:B:169:PRO:HD2	2.14	0.47
1:A:385:TRP:CD2	1:A:394:HIS:HB2	2.50	0.47
1:A:314:GLN:NE2	5:A:603:HOH:O	2.48	0.46
1:B:165:ILE:HD12	1:B:223:LEU:HB3	1.97	0.46
1:B:174:PRO:HB2	1:B:183:TRP:CH2	2.51	0.45
1:A:178:PRO:HB3	1:A:183:TRP:NE1	2.32	0.44
1:B:242:TRP:O	1:B:246:ARG:HG2	2.17	0.44
1:A:177:THR:HB	1:B:229:GLY:HA2	2.01	0.42
1:B:140:VAL:HA	1:B:156:THR:O	2.19	0.42
1:B:174:PRO:HB2	1:B:183:TRP:CZ3	2.54	0.42
1:B:146:LYS:HE2	1:B:173:TYR:CD1	2.55	0.42
1:A:350:ALA:HB1	1:A:358[A]:MET:HE1	2.02	0.42
1:A:373:SER:O	1:A:377:CYS:HB2	2.20	0.42
1:B:137:PRO:HG3	1:B:242:TRP:HZ2	1.85	0.42
1:B:203:LEU:HD23	1:B:203:LEU:C	2.44	0.41
1:A:304:LEU:CD2	1:A:449:TRP:CD2	3.04	0.40
1:B:187:ILE:O	1:B:266:VAL:HA	2.21	0.40
1:B:385:TRP:CD2	1:B:394:HIS:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/401 (95%)	365 (96%)	15 (4%)	0	100	100
1	B	379/401 (94%)	366 (97%)	13 (3%)	0	100	100
All	All	759/802 (95%)	731 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/320 (95%)	304 (100%)	0	100	100
1	B	303/320 (95%)	299 (99%)	4 (1%)	61	38
All	All	607/640 (95%)	603 (99%)	4 (1%)	76	62

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	175	SER
1	B	256	THR
1	B	370	SER
1	B	433	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	331	ASN
1	A	334	ASN
1	B	316	GLN
1	B	443	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	XYL	C	1	2	9,9,9	0.83	0	11,11,11	1.09	1 (9%)
2	XYP	C	2	2	9,9,10	1.89	3 (33%)	10,12,14	1.40	2 (20%)
2	XYP	C	3	2	9,9,10	1.85	2 (22%)	10,12,14	1.28	1 (10%)
2	GCV	C	4	2	13,13,14	1.62	3 (23%)	14,18,20	1.48	3 (21%)
2	XYP	C	5	2	9,9,10	1.88	2 (22%)	10,12,14	1.11	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYL	C	1	2	-	2/12/12/12	-
2	XYP	C	2	2	-	-	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYP	C	3	2	-	-	0/1/1/1
2	GCV	C	4	2	-	4/6/23/26	0/1/1/1
2	XYP	C	5	2	-	-	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	5	XYP	O5-C1	4.05	1.50	1.43
2	C	3	XYP	O5-C1	4.00	1.50	1.43
2	C	2	XYP	O5-C1	3.99	1.50	1.43
2	C	4	GCV	O5-C1	3.49	1.49	1.43
2	C	4	GCV	O5-C5	2.91	1.49	1.43
2	C	5	XYP	O5-C5	2.75	1.48	1.43
2	C	3	XYP	O5-C5	2.71	1.48	1.43
2	C	2	XYP	O5-C5	2.56	1.47	1.43
2	C	4	GCV	C2-C3	-2.34	1.48	1.52
2	C	2	XYP	C2-C3	-2.23	1.49	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	XYP	C5-C4-C3	3.58	114.86	109.64
2	C	4	GCV	O3-C3-C2	-2.91	104.11	110.05
2	C	4	GCV	O2-C2-C1	-2.80	102.81	109.22
2	C	2	XYP	C1-C2-C3	2.58	113.40	109.64
2	C	5	XYP	C5-C4-C3	2.53	113.33	109.64
2	C	1	XYL	C5-C4-C3	-2.46	107.17	112.17
2	C	2	XYP	C4-C3-C2	2.33	113.68	110.92
2	C	4	GCV	O2-C2-C3	2.03	114.35	110.15

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	4	GCV	C3-C4-O4-C7
2	C	4	GCV	C5-C4-O4-C7
2	C	1	XYL	C1-C2-C3-C4
2	C	1	XYL	O2-C2-C3-C4
2	C	4	GCV	O5-C5-C6-O6A
2	C	4	GCV	O5-C5-C6-O6B

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	4	GCV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	501	1	14,14,15	0.29	0	17,19,21	0.55	0
3	NAG	A	501	1	14,14,15	0.42	0	17,19,21	0.57	0
4	EDO	A	502	-	3,3,3	0.43	0	2,2,2	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	501	1	-	0/6/23/26	0/1/1/1
3	NAG	A	501	1	-	0/6/23/26	0/1/1/1
4	EDO	A	502	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	380/401 (94%)	-0.12	8 (2%) 63 70	13, 24, 39, 62	2 (0%)
1	B	380/401 (94%)	0.78	70 (18%) 3 4	17, 32, 67, 84	1 (0%)
All	All	760/802 (94%)	0.33	78 (10%) 12 13	13, 27, 61, 84	3 (0%)

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	250	ALA	5.3
1	B	144	PHE	4.9
1	B	183	TRP	4.9
1	B	159	LEU	4.9
1	B	178	PRO	4.8
1	B	257	ALA	4.6
1	B	177	THR	4.6
1	B	171	ILE	4.6
1	B	180	ALA	4.4
1	B	259	ILE	4.3
1	B	153	LEU	4.2
1	B	256	THR	3.8
1	B	179	PRO	3.8
1	B	261	THR	3.8
1	B	155	ILE	3.7
1	B	251	LEU	3.7
1	B	255	PRO	3.6
1	B	254	THR	3.6
1	B	181	ASN	3.6
1	B	165	ILE	3.6
1	B	154	ALA	3.6
1	B	151	GLY	3.6
1	B	79	GLY	3.5
1	B	173	TYR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	162	SER	3.4
1	B	175	SER	3.4
1	B	167	PHE	3.3
1	B	148	GLY	3.3
1	B	152	THR	3.3
1	B	149	ASN	3.3
1	B	142	ALA	3.2
1	B	139	VAL	3.2
1	A	79	GLY	3.1
1	A	176	GLY	3.1
1	B	182	GLY	3.1
1	B	248	ILE	3.1
1	B	170	THR	3.1
1	B	141	THR	3.1
1	A	80	ALA	3.0
1	B	163	GLN	3.0
1	B	140	VAL	2.9
1	B	84	ILE	2.9
1	B	161	ASN	2.9
1	B	158	GLY	2.8
1	B	156	THR	2.8
1	B	145	SER	2.7
1	B	80	ALA	2.7
1	A	90	ASN	2.7
1	B	258	GLN	2.7
1	B	253	MET	2.7
1	B	174	PRO	2.6
1	B	164	THR	2.6
1	B	184	PRO	2.6
1	B	150	THR	2.6
1	B	168	SER	2.6
1	B	176	GLY	2.5
1	B	160	SER	2.5
1	A	458	THR	2.4
1	A	180	ALA	2.4
1	B	135	PRO	2.3
1	B	185	LEU	2.3
1	B	247	ILE	2.3
1	B	109	ARG	2.3
1	B	205	TYR	2.3
1	B	166	LYS	2.3
1	B	201	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	431	ASN	2.2
1	B	157	ALA	2.2
1	B	223	LEU	2.2
1	B	169	PRO	2.2
1	B	172	SER	2.2
1	B	186	ILE	2.2
1	B	137	PRO	2.1
1	B	199	GLY	2.1
1	B	147	SER	2.1
1	A	392	ASP	2.0
1	B	458	THR	2.0
1	B	83	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	XYL	C	1	10/10	0.88	0.11	36,48,60,62	0
2	XYP	C	5	9/10	0.89	0.09	35,42,48,50	0
2	XYP	C	2	9/10	0.95	0.07	25,33,43,49	0
2	GCV	C	4	13/14	0.97	0.06	23,28,33,33	0
2	XYP	C	3	9/10	0.97	0.06	24,30,41,41	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	B	501	14/15	0.92	0.10	24,29,34,36	0
4	EDO	A	502	4/4	0.92	0.11	21,35,44,44	0
3	NAG	A	501	14/15	0.95	0.07	22,27,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.