



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 07:28 AM UTC

PDB ID : 6RVH / pdb_00006rvh
Title : NADH-dependent Coenzyme A Disulfide Reductase soaked with Menadione
Authors : Koepke, J.; Preu, J.
Deposited on : 2019-05-31
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	2% 43% 34% 18% 5%
1	B	443	2% 43% 35% 16% 6%
1	C	443	2% 41% 36% 18% 5%
1	D	443	2% 41% 37% 16% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:


Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	VK3	D	503	-	-	-	X

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total 3379	C 2147	N 608	O 617	S 7	0	0	0
1	B	443	Total 3379	C 2147	N 608	O 617	S 7	0	0	0
1	C	443	Total 3378	C 2146	N 608	O 617	S 7	0	0	0
1	D	443	Total 3377	C 2146	N 608	O 616	S 7	0	0	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	B	1	Total 53	C 27	N 9	O 15	P 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is COENZYME A (CCD ID: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	16	3		
3	B	1	Total	C	N	O	P	0	0
			48	21	7	16	3		
3	C	1	Total	C	N	O	P	0	0
			48	21	7	16	3		
3	D	1	Total	C	N	O	P	0	0
			48	21	7	16	3		

- Molecule 4 is MENADIONE (CCD ID: VK3) (formula: C₁₁H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	11	2		
4	B	1	Total	C	O	0	0
			13	11	2		
4	C	1	Total	C	O	0	0
			13	11	2		
4	D	1	Total	C	O	0	0
			13	11	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	35	Total	O	0	0
			35	35		
5	B	35	Total	O	0	0
			35	35		
5	C	36	Total	O	0	0
			36	36		
5	D	30	Total	O	0	0
			30	30		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NADH oxidase




• Molecule 1: NADH oxidase







● Molecule 1: NADH oxidase



● Molecule 1: NADH oxidase





4 Data and refinement statistics

Property	Value	Source
Space group	P 4 ₂ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	159.95Å 159.95Å 255.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.35 – 3.00 58.35 – 3.00	Depositor EDS
% Data completeness (in resolution range)	95.0 (58.35-3.00) 95.0 (58.35-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.29	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.6.4_486	Depositor
R, R_{free}	0.215 , 0.259 0.207 , 0.250	Depositor DCC
R_{free} test set	2000 reflections (3.14%)	wwPDB-VP
Wilson B-factor (Å ²)	58.0	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	14105	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, FAD, VK3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.13	107/3451 (3.1%)	1.91	106/4686 (2.3%)
1	B	1.97	75/3451 (2.2%)	1.92	108/4686 (2.3%)
1	C	1.98	73/3450 (2.1%)	1.87	102/4683 (2.2%)
1	D	1.92	61/3449 (1.8%)	1.92	128/4683 (2.7%)
All	All	2.00	316/13801 (2.3%)	1.91	444/18738 (2.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8
1	B	0	5
1	C	0	5
1	D	0	6
All	All	0	24

The worst 5 of 316 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	122	ILE	CA-CB	-14.26	1.43	1.54
1	B	93	VAL	CA-CB	12.88	1.72	1.54
1	A	406	ALA	CA-CB	-12.38	1.33	1.53
1	A	437	ILE	CA-CB	12.37	1.70	1.54
1	D	396	HIS	C-O	12.21	1.40	1.24

The worst 5 of 444 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	282	VAL	N-CA-C	-13.13	99.95	112.29
1	D	400	ARG	N-CA-C	-11.25	99.51	113.01
1	B	400	ARG	N-CA-C	-11.11	99.67	113.01
1	C	54	ILE	CA-C-N	10.53	133.00	119.84
1	C	54	ILE	C-N-CA	10.53	133.00	119.84

There are no chirality outliers.

5 of 24 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	127	GLN	Peptide
1	A	90	THR	Peptide
1	A	91	LEU	Peptide
1	A	93	VAL	Peptide
1	A	98	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3379	0	3407	213	0
1	B	3379	0	3407	229	0
1	C	3378	0	3402	222	0
1	D	3377	0	3402	221	0
2	A	53	0	31	3	0
2	B	53	0	31	5	0
2	C	53	0	31	5	0
2	D	53	0	31	5	0
3	A	48	0	31	4	0
3	B	48	0	31	10	0
3	C	48	0	30	11	0
3	D	48	0	31	5	0
4	A	13	0	8	4	0
4	B	13	0	8	1	0
4	C	13	0	8	2	0
4	D	13	0	8	2	0
5	A	35	0	0	10	0
5	B	35	0	0	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	36	0	0	25	0
5	D	30	0	0	22	0
All	All	14105	0	13897	872	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 872 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:439:ALA:O	1:A:441:GLN:HB2	1.11	1.28
1:D:439:ALA:O	1:D:441:GLN:HB2	1.30	1.28
1:C:439:ALA:O	1:C:441:GLN:HB2	1.21	1.26
1:B:439:ALA:O	1:B:441:GLN:HB2	1.02	1.19
1:A:396:HIS:ND1	1:B:396:HIS:CE1	2.12	1.17

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/443 (100%)	378 (86%)	44 (10%)	19 (4%)	2	12
1	B	441/443 (100%)	381 (86%)	40 (9%)	20 (4%)	2	12
1	C	441/443 (100%)	385 (87%)	38 (9%)	18 (4%)	2	13
1	D	441/443 (100%)	386 (88%)	36 (8%)	19 (4%)	2	12
All	All	1764/1772 (100%)	1530 (87%)	158 (9%)	76 (4%)	2	12

5 of 76 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	THR
1	A	149	ALA
1	A	398	ALA
1	A	441	GLN
1	A	442	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/338 (100%)	275 (81%)	63 (19%)	1	9
1	B	338/338 (100%)	276 (82%)	62 (18%)	2	9
1	C	337/338 (100%)	279 (83%)	58 (17%)	2	11
1	D	337/338 (100%)	277 (82%)	60 (18%)	2	10
All	All	1350/1352 (100%)	1107 (82%)	243 (18%)	2	10

5 of 243 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	394	ARG
1	D	237	VAL
1	C	133	LEU
1	D	227	THR
1	D	408	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	94	HIS
1	C	287	HIS
1	D	287	HIS
1	C	272	ASN
1	C	359	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	COA	B	503	1	47,50,50	1.62	5 (10%)	69,75,75	2.42	21 (30%)
4	VK3	C	503	-	14,14,14	2.08	2 (14%)	20,20,20	2.28	6 (30%)
4	VK3	B	501	-	14,14,14	2.12	3 (21%)	20,20,20	2.63	11 (55%)
4	VK3	D	503	-	14,14,14	2.29	4 (28%)	20,20,20	2.74	8 (40%)
3	COA	C	502	1	47,50,50	1.68	9 (19%)	69,75,75	2.43	23 (33%)
2	FAD	C	501	-	58,58,58	2.81	24 (41%)	85,89,89	2.26	29 (34%)
4	VK3	A	503	-	14,14,14	2.20	3 (21%)	20,20,20	2.86	11 (55%)
2	FAD	B	502	-	58,58,58	3.12	28 (48%)	85,89,89	2.38	34 (40%)
2	FAD	D	501	-	58,58,58	2.78	27 (46%)	85,89,89	2.28	31 (36%)
2	FAD	A	501	-	58,58,58	2.83	26 (44%)	85,89,89	2.24	26 (30%)
3	COA	A	502	1	47,50,50	1.76	7 (14%)	69,75,75	2.48	23 (33%)
3	COA	D	502	1	47,50,50	1.80	7 (14%)	69,75,75	2.35	24 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COA	B	503	1	-	20/48/64/64	0/3/3/3
4	VK3	C	503	-	-	-	0/2/2/2
4	VK3	B	501	-	-	-	0/2/2/2
4	VK3	D	503	-	-	-	0/2/2/2
3	COA	C	502	1	-	19/48/64/64	0/3/3/3
2	FAD	C	501	-	-	15/34/50/50	0/6/6/6
4	VK3	A	503	-	-	-	0/2/2/2
2	FAD	B	502	-	-	16/34/50/50	0/6/6/6
2	FAD	D	501	-	-	18/34/50/50	0/6/6/6
2	FAD	A	501	-	-	17/34/50/50	0/6/6/6
3	COA	A	502	1	-	19/48/64/64	0/3/3/3
3	COA	D	502	1	-	19/48/64/64	0/3/3/3

The worst 5 of 145 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	FAD	C5A-C4A	9.10	1.55	1.39
2	A	501	FAD	PA-O3P	9.02	1.69	1.59
2	A	501	FAD	C5A-C4A	8.84	1.54	1.39
2	B	502	FAD	PA-O3P	8.84	1.69	1.59
2	D	501	FAD	C5A-C4A	8.74	1.54	1.39

The worst 5 of 247 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	502	COA	C5A-C4A-N3A	-8.53	114.98	126.72
3	B	503	COA	C5A-C4A-N3A	-8.30	115.29	126.72
3	A	502	COA	C5A-C4A-N3A	-7.67	116.16	126.72
2	D	501	FAD	O4B-C1B-N9A	7.37	122.24	108.09
3	D	502	COA	C7P-C6P-C5P	-7.32	100.20	112.39

There are no chirality outliers.

5 of 143 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	FAD	C5B-O5B-PA-O3P
2	A	501	FAD	C1'-C2'-C3'-O3'
2	A	501	FAD	C3'-C4'-C5'-O5'

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	A	501	FAD	O4'-C4'-C5'-O5'
2	A	501	FAD	C5'-O5'-P-O1P


There are no ring outliers.


12 monomers are involved in 51 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	503	COA	10	0
4	C	503	VK3	2	0
4	B	501	VK3	1	0
4	D	503	VK3	2	0
3	C	502	COA	11	0
2	C	501	FAD	5	0
4	A	503	VK3	4	0
2	B	502	FAD	5	0
2	D	501	FAD	5	0
2	A	501	FAD	3	0
3	A	502	COA	4	0
3	D	502	COA	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.












Ligand FAD A 501



Ligand COA A 502





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	443/443 (100%)	-0.08	8 (1%) 67 44	28, 47, 74, 134	0
1	B	443/443 (100%)	-0.04	10 (2%) 61 38	29, 48, 75, 137	0
1	C	443/443 (100%)	-0.08	9 (2%) 65 41	28, 48, 74, 136	0
1	D	443/443 (100%)	-0.05	10 (2%) 61 38	27, 49, 76, 137	0
All	All	1772/1772 (100%)	-0.06	37 (2%) 63 40	27, 48, 76, 137	0

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	6.1
1	D	1	MET	6.0
1	A	1	MET	4.6
1	A	396	HIS	4.4
1	D	88	LEU	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	VK3	D	503	13/13	0.51	0.44	80,101,140,141	0
4	VK3	A	503	13/13	0.65	0.33	80,100,137,143	0
4	VK3	B	501	13/13	0.72	0.34	79,106,136,140	0
4	VK3	C	503	13/13	0.75	0.31	81,103,139,139	0
2	FAD	B	502	53/53	0.80	0.19	55,83,109,156	0
2	FAD	D	501	53/53	0.84	0.18	54,81,111,141	0
2	FAD	C	501	53/53	0.86	0.17	50,79,108,131	0
2	FAD	A	501	53/53	0.87	0.16	54,81,113,152	0
3	COA	A	502	48/48	0.94	0.09	38,52,70,94	0
3	COA	B	503	48/48	0.95	0.09	39,52,75,91	0
3	COA	C	502	48/48	0.96	0.09	37,52,74,99	0
3	COA	D	502	48/48	0.96	0.09	36,52,73,94	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FAD B 502:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 501:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around COA A 502:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around COA B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around COA C 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.