



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:28 AM UTC

PDB ID : 4RWY / pdb_00004rwy
Title : Crystal structure of VH1-46 germline-derived CD4-binding site-directed anti-body 8ANC131 in complex with HIV-1 clade B YU2 gp120
Authors : Acharya, P.; Luongo, T.S.; Kwong, P.D.
Deposited on : 2014-12-08
Resolution : 2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	6% 86% 10%
2	H	227	6% 93% 6%
3	L	213	6% 91% 7%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 12412 atoms, of which 6038 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 YU2 gp120.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	337	5256	1667	2600	461	507	21	0	5	0


- Molecule 2 is a protein called Antibody 8ANC131 Heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	H	227	3408	1077	1699	296	329	7	0	0	0

- Molecule 3 is a protein called Antibody 8ANC131 Light chain.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	L	213	3270	1036	1612	291	326	5	0	1	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
4	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
4	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	S	0	0
			32	8	17	2	4	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			10	2	6	2		
6	H	1	Total	C	H	O	0	0
			10	2	6	2		
6	H	1	Total	C	H	O	0	0
			10	2	6	2		
6	H	1	Total	C	H	O	0	0
			10	2	6	2		
6	H	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	H	1	Total	Na	0	0
			1	1		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	80	Total	O	0	0
			80	80		
8	H	82	Total	O	0	0
			82	82		
8	L	69	Total	O	0	0
			69	69		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HIV-1 YU2 gp120

Chain A: 




- Molecule 2: Antibody 8ANC131 Heavy chain

Chain H: 



- Molecule 3: Antibody 8ANC131 Light chain

Chain L: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.25Å 67.14Å 214.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.84 – 2.13 28.84 – 2.13	Depositor EDS
% Data completeness (in resolution range)	79.3 (28.84-2.13) 73.6 (28.84-2.13)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 2.14Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1690)	Depositor
R, R_{free}	0.219 , 0.254 0.224 , 0.252	Depositor DCC
R_{free} test set	2139 reflections (3.92%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.714	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 36.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12412	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, EDO, NA, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/2738	0.58	0/3716
2	H	0.27	0/1749	0.58	0/2378
3	L	0.29	0/1700	0.59	0/2308
All	All	0.28	0/6187	0.58	0/8402

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	L	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	L	50	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2656	2600	2583	6	0
2	H	1709	1699	1696	10	0
3	L	1658	1612	1611	11	0
4	A	84	80	78	1	0
5	A	15	17	17	0	0
6	A	4	6	6	0	0
6	H	16	24	24	0	0
7	H	1	0	0	0	0
8	A	80	0	0	1	0
8	H	82	0	0	3	0
8	L	69	0	0	1	0
All	All	6374	6038	6015	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:189:HIS:O	3:L:211:ARG:NH1	2.29	0.66
2:H:134:GLY:O	2:H:186:SER:N	2.33	0.61
3:L:142:ARG:NH1	8:L:323:HOH:O	2.33	0.61
2:H:131:THR:OG1	2:H:135:THR:O	2.15	0.59
1:A:234:ASN:HD22	4:A:502:NAG:H83	1.69	0.57
3:L:50:ALA:HB1	3:L:51:PRO:CD	2.38	0.54
1:A:463:ASN:O	1:A:463:ASN:ND2	2.40	0.53
3:L:2:ILE:O	3:L:97:ASN:ND2	2.42	0.53
1:A:231:LYS:NZ	8:A:663:HOH:O	2.43	0.51
2:H:143:LYS:NZ	2:H:171:GLN:OE1	2.43	0.51
2:H:129:LYS:NZ	2:H:133:GLY:O	2.44	0.50
2:H:188:SER:O	2:H:192:GLN:N	2.45	0.49
3:L:51:PRO:HA	3:L:52:SER:CB	2.43	0.49
2:H:80:MET:O	8:H:458:HOH:O	2.20	0.48
1:A:327:ARG:NH2	1:A:422:GLN:OE1	2.48	0.45
1:A:461:ASP:N	1:A:461:ASP:OD1	2.49	0.45
2:H:80:MET:N	8:H:458:HOH:O	2.43	0.44
2:H:100(F):TYR:HB2	3:L:50:ALA:HB2	2.00	0.43
3:L:108:ARG:NE	3:L:170:ASP:O	2.50	0.43
1:A:257:THR:O	1:A:375:SER:N	2.52	0.42
3:L:2:ILE:HG22	3:L:3:VAL:N	2.35	0.42
3:L:13:LEU:CD1	3:L:19:ALA:HB2	2.50	0.42
3:L:202:ARG:H	3:L:202:ARG:HD2	1.85	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:55:ALA:HB1	3:L:56:PRO:HD2	2.03	0.41
2:H:20:ILE:N	8:H:458:HOH:O	2.53	0.41
2:H:134:GLY:C	2:H:185:PRO:HA	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	336/376 (89%)	325 (97%)	9 (3%)	2 (1%)	21	18
2	H	225/227 (99%)	218 (97%)	6 (3%)	1 (0%)	30	28
3	L	212/213 (100%)	204 (96%)	6 (3%)	2 (1%)	14	10
All	All	773/816 (95%)	747 (97%)	21 (3%)	5 (1%)	18	18

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	45	TRP
2	H	129	LYS
3	L	2	ILE
1	A	268	GLU
3	L	51	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/328 (92%)	296 (98%)	7 (2%)	44	50
2	H	190/190 (100%)	188 (99%)	2 (1%)	65	74
3	L	186/185 (100%)	182 (98%)	4 (2%)	45	52
All	All	679/703 (97%)	666 (98%)	13 (2%)	50	57

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	230	ASP
1	A	267	GLU
1	A	340	ASN
1	A	360	ILE
1	A	416	LEU
1	A	461	ASP
1	A	463	ASN
2	H	131	THR
2	H	193	THR
3	L	20	THR
3	L	48	ILE
3	L	52	SER
3	L	90	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	HIS
3	L	97	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	501	1	14,14,15	0.20	0	17,19,21	0.49	0
4	NAG	A	502	1	14,14,15	0.19	0	17,19,21	0.47	0
6	EDO	H	302	-	3,3,3	0.43	0	2,2,2	0.35	0
6	EDO	H	304	-	3,3,3	0.42	0	2,2,2	0.35	0
6	EDO	H	301	-	3,3,3	0.43	0	2,2,2	0.33	0
5	EPE	A	507	-	15,15,15	0.80	1 (6%)	19,20,20	1.54	2 (10%)
4	NAG	A	503	1	14,14,15	0.28	0	17,19,21	0.49	0
4	NAG	A	504	1	14,14,15	0.22	0	17,19,21	0.50	0
6	EDO	A	508	-	3,3,3	0.43	0	2,2,2	0.28	0
4	NAG	A	506	1	14,14,15	0.33	0	17,19,21	0.44	0
6	EDO	H	303	-	3,3,3	0.42	0	2,2,2	0.37	0
4	NAG	A	505	1	14,14,15	0.24	0	17,19,21	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	501	1	-	2/6/23/26	0/1/1/1
4	NAG	A	502	1	-	2/6/23/26	0/1/1/1
6	EDO	H	302	-	-	0/1/1/1	-
6	EDO	H	304	-	-	0/1/1/1	-
6	EDO	H	301	-	-	0/1/1/1	-
5	EPE	A	507	-	-	4/9/19/19	0/1/1/1
4	NAG	A	503	1	-	2/6/23/26	0/1/1/1
4	NAG	A	504	1	-	0/6/23/26	0/1/1/1
6	EDO	A	508	-	-	0/1/1/1	-
4	NAG	A	506	1	-	0/6/23/26	0/1/1/1
6	EDO	H	303	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	505	1	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	507	EPE	C10-S	2.77	1.81	1.77

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	507	EPE	C5-N4-C3	3.82	117.07	108.84
5	A	507	EPE	C7-N4-C5	3.01	119.27	111.24

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	507	EPE	C9-C10-S-O2S
5	A	507	EPE	C8-C7-N4-C5
4	A	501	NAG	C8-C7-N2-C2
4	A	501	NAG	O7-C7-N2-C2
4	A	502	NAG	C8-C7-N2-C2
4	A	502	NAG	O7-C7-N2-C2
5	A	507	EPE	C9-C10-S-O3S
5	A	507	EPE	C9-C10-S-O1S
4	A	503	NAG	C4-C5-C6-O6
4	A	505	NAG	C1-C2-N2-C7
4	A	503	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	502	NAG	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	337/376 (89%)	0.54	22 (6%)	25 27	29, 59, 102, 144	5 (1%)
2	H	227/227 (100%)	0.35	14 (6%)	26 29	30, 49, 103, 215	1 (0%)
3	L	213/213 (100%)	0.47	12 (5%)	30 33	34, 58, 91, 148	1 (0%)
All	All	777/816 (95%)	0.47	48 (6%)	26 29	29, 56, 101, 215	7 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	73	ARG	7.8
3	L	51	PRO	5.8
3	L	50	ALA	5.4
2	H	2	GLY	5.3
1	A	411	GLY	5.1
2	H	127	SER	4.5
3	L	202	ARG	4.5
2	H	133	GLY	3.9
2	H	131	THR	3.8
1	A	44	VAL	3.7
3	L	52	SER	3.7
3	L	2	ILE	3.5
2	H	134	GLY	3.4
1	A	464	GLY	3.1
1	A	45	TRP	3.0
2	H	216	CYS	2.8
1	A	491[A]	ILE	2.8
1	A	265	LEU	2.7
1	A	266	ALA	2.7
1	A	257	THR	2.7
1	A	301	ASN	2.7
3	L	74	VAL	2.7
3	L	68	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	215	SER	2.5
1	A	231	LYS	2.4
3	L	49[A]	HIS	2.4
2	H	128	SER	2.4
3	L	67	SER	2.4
1	A	242	VAL	2.3
1	A	416	LEU	2.3
2	H	135	THR	2.3
1	A	72	HIS	2.3
3	L	20	THR	2.3
1	A	234	ASN	2.3
1	A	463	ASN	2.3
1	A	336	THR	2.2
2	H	161	SER	2.2
1	A	340	ASN	2.2
1	A	459	GLY	2.1
1	A	358	THR	2.1
1	A	439	ILE	2.1
3	L	30	LEU	2.1
1	A	239	CYS	2.1
2	H	136	ALA	2.1
1	A	92	ASN	2.0
2	H	1	GLN	2.0
2	H	191	THR	2.0
3	L	1	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	EDO	H	301	4/4	0.55	0.18	70,85,85,85	0
6	EDO	H	304	4/4	0.60	0.21	80,96,96,97	0
4	NAG	A	506	14/15	0.61	0.16	75,100,129,129	0
4	NAG	A	501	14/15	0.67	0.16	85,93,111,113	0
6	EDO	H	302	4/4	0.72	0.16	70,84,84,84	0
4	NAG	A	505	14/15	0.72	0.19	102,118,140,144	0
4	NAG	A	502	14/15	0.79	0.11	54,62,73,74	0
4	NAG	A	504	14/15	0.80	0.09	51,57,68,68	0
4	NAG	A	503	14/15	0.83	0.10	40,46,55,55	0
6	EDO	A	508	4/4	0.85	0.24	91,110,113,113	0
7	NA	H	305	1/1	0.85	0.52	60,60,60,60	0
6	EDO	H	303	4/4	0.91	0.23	51,61,63,63	0
5	EPE	A	507	15/15	0.95	0.09	42,50,54,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.