



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 04:22 PM UTC

PDB ID : 6RZ3 / pdb_00006rz3
Title : Crystal structure of a complex between the DNA-binding domain of p53 and the carboxyl-terminal conserved region of iASPP
Authors : Chen, S.; Ren, J.; Jones, E.Y.; Lu, X.
Deposited on : 2019-06-12
Resolution : 4.23 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1042 (4.60-3.88)
Clashscore	190562	1043 (4.58-3.90)
Ramachandran outliers	187476	1031 (4.62-3.86)
Sidechain outliers	187428	1016 (4.62-3.86)
RSRZ outliers	180081	1039 (4.60-3.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	
2	B	238	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2867 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellular tumor antigen p53.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	201	Total	C	N	O	S	0	0	0
			1584	979	293	296	16			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	MET	-	initiating methionine	UNP P04637

- Molecule 2 is a protein called RelA-associated inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	167	Total	C	N	O	S	0	0	0
			1282	808	213	252	9			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	624	MET	-	initiating methionine	UNP Q8WUF5
B	829	LEU	-	expression tag	UNP Q8WUF5
B	830	GLU	-	expression tag	UNP Q8WUF5
B	831	GLY	-	expression tag	UNP Q8WUF5
B	832	GLY	-	expression tag	UNP Q8WUF5
B	833	SER	-	expression tag	UNP Q8WUF5
B	834	GLY	-	expression tag	UNP Q8WUF5
B	835	GLY	-	expression tag	UNP Q8WUF5
B	836	SER	-	expression tag	UNP Q8WUF5
B	837	GLY	-	expression tag	UNP Q8WUF5
B	838	LEU	-	expression tag	UNP Q8WUF5
B	839	ASN	-	expression tag	UNP Q8WUF5
B	840	ASP	-	expression tag	UNP Q8WUF5
B	841	ILE	-	expression tag	UNP Q8WUF5
B	842	PHE	-	expression tag	UNP Q8WUF5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	843	GLU	-	expression tag	UNP Q8WUF5
B	844	ALA	-	expression tag	UNP Q8WUF5
B	845	GLN	-	expression tag	UNP Q8WUF5
B	846	LYS	-	expression tag	UNP Q8WUF5
B	847	ILE	-	expression tag	UNP Q8WUF5
B	848	GLU	-	expression tag	UNP Q8WUF5
B	849	TRP	-	expression tag	UNP Q8WUF5
B	850	HIS	-	expression tag	UNP Q8WUF5
B	851	GLU	-	expression tag	UNP Q8WUF5
B	852	GLY	-	expression tag	UNP Q8WUF5
B	853	ARG	-	expression tag	UNP Q8WUF5
B	854	THR	-	expression tag	UNP Q8WUF5
B	855	LYS	-	expression tag	UNP Q8WUF5
B	856	HIS	-	expression tag	UNP Q8WUF5
B	857	HIS	-	expression tag	UNP Q8WUF5
B	858	HIS	-	expression tag	UNP Q8WUF5
B	859	HIS	-	expression tag	UNP Q8WUF5
B	860	HIS	-	expression tag	UNP Q8WUF5
B	861	HIS	-	expression tag	UNP Q8WUF5


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 1: Cellular tumor antigen p53

LYS	MET GLU ALA PRO ARG MET PRO GLU ALA ALA PRO PRO VAL ALA PRO ALA ALA ALA PRO THR PRO ALA ALA PRO ALA PRO SER W91 P98 S99 Q100 R110 L114 L115 S116 G117 T123 I162 S241 C242 M243 R248 P278 R283 E287 W301
-----	---

Chain B: 59% 11% 30%



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	71.09Å 71.09Å 255.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.80 – 4.23 29.80 – 4.23	Depositor EDS
% Data completeness (in resolution range)	93.2 (29.80-4.23) 93.2 (29.80-4.23)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 4.26Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.243 , 0.292 0.255 , 0.313	Depositor DCC
R_{free} test set	473 reflections (9.15%)	wwPDB-VP
Wilson B-factor (Å ²)	160.5	Xtriage
Anisotropy	0.535	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 126.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2867	wwPDB-VP
Average B, all atoms (Å ²)	198.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/1622	0.77	0/2200
2	B	0.57	0/1318	0.91	3/1800 (0.2%)
All	All	0.54	0/2940	0.84	3/4000 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	725	SER	N-CA-C	9.65	121.88	111.36
2	B	726	ASP	N-CA-C	-7.43	102.14	113.89
2	B	817	LEU	N-CA-C	-5.94	105.53	112.89

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1584	0	1544	10	0
2	B	1282	0	1175	17	0
3	A	1	0	0	0	0
All	All	2867	0	2719	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:766:LEU:O	2:B:767:TRP:HD1	1.40	1.05
2:B:766:LEU:O	2:B:767:TRP:CD1	2.23	0.90
1:A:123:THR:O	1:A:278:PRO:HG3	1.94	0.68
2:B:687:ASN:HD21	2:B:718:ALA:H	1.43	0.66
1:A:241:SER:HB2	1:A:248:ARG:NH2	2.20	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/232 (86%)	194 (98%)	4 (2%)	1 (0%)	24	62
2	B	165/238 (69%)	160 (97%)	5 (3%)	0	100	100
All	All	364/470 (77%)	354 (97%)	9 (2%)	1 (0%)	36	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	117	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/201 (90%)	178 (98%)	3 (2%)	53	67
2	B	131/191 (69%)	131 (100%)	0	100	100
All	All	312/392 (80%)	309 (99%)	3 (1%)	68	75

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ARG
1	A	243	MET
1	A	287	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	HIS
1	A	235	ASN
1	A	239	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/232 (86%)	-0.34	0 100 100	147, 206, 248, 285	0
2	B	167/238 (70%)	-0.22	0 100 100	145, 179, 240, 340	0
All	All	368/470 (78%)	-0.28	0 100 100	145, 195, 248, 340	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ZN	A	301	1/1	0.97	0.04	210,210,210,210	0

6.5 Other polymers [i](#)

There are no such residues in this entry.