



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:18 AM UTC

PDB ID : 6RZ4 / pdb_00006rz4
Title : Crystal structure of cysteinyl leukotriene receptor 1 in complex with pranlukast
Authors : Luginina, A.; Gusach, A.; Marin, E.; Mishin, A.; Brouillette, R.; Popov, P.; Shiryaeva, A.; Besserer-Offroy, E.; Longpre, J.M.; Lyapina, E.; Ishchenko, A.; Patel, N.; Polovinkin, V.; Safronova, N.; Bogorodskiy, A.; Edelweiss, E.; Liu, W.; Batyuk, A.; Gordeliy, V.; Han, G.W.; Sarret, P.; Katritch, V.; Borshchevskiy, V.; Cherezov, V.
Deposited on : 2019-06-12
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	5% 87% 6% 7%

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cysteinyl leukotriene receptor 1, Soluble cytochrome b562, Cysteinyl leukotriene receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	395	Total	C	N	O	S	0	2	0
			3020	1974	488	533	25			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q9Y271
A	-1	GLY	-	expression tag	UNP Q9Y271
A	0	THR	-	expression tag	UNP Q9Y271
A	1000	SER	-	linker	UNP Q9Y271
A	1007	TRP	MET	conflict	UNP P0ABE7
A	1102	ILE	HIS	conflict	UNP P0ABE7
A	1106	LEU	-	linker	UNP P0ABE7
A	?	SER	-	linker	UNP P0ABE7
A	?	GLY	-	linker	UNP P0ABE7

- Molecule 2 is pranlukast (CCD ID: KNT) (formula: C₂₇H₂₃N₅O₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			36	27	5	4		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0

- Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (CCD ID: OLC) (formula: $C_{21}H_{40}O_4$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 13 9 4	0	0
4	A	1	Total C O 12 8 4	0	0
4	A	1	Total C O 25 21 4	0	0
4	A	1	Total C 7 7	0	0
4	A	1	Total C O 21 17 4	0	0
4	A	1	Total C 17 17	0	0
4	A	1	Total C 6 6	0	0
4	A	1	Total C O 10 6 4	0	0
4	A	1	Total C 11 11	0	0

- Molecule 5 is OLEIC ACID (CCD ID: OLA) (formula: $C_{18}H_{34}O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 11 9 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	16	Total	O	0	0
			16	16		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cysteinyl leukotriene receptor 1, Soluble cytochrome b562, Cysteinyl leukotriene receptor 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.63Å 45.35Å 86.43Å 90.00° 91.47° 90.00°	Depositor
Resolution (Å)	29.30 – 2.70 29.30 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.30-2.70) 99.8 (29.30-2.70)	Depositor EDS
R_{merge}	0.27	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.68Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.228 , 0.254 (Not available) , 0.271	Depositor DCC
R_{free} test set	641 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	42.9	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 56.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.052 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3206	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OLC, OLA, NA, KNT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/3091	1.39	0/4213

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3020	0	2900	11	0
2	A	36	0	0	0	0
3	A	1	0	0	0	0
4	A	122	0	169	2	0
5	A	11	0	14	0	0
6	A	16	0	0	0	0
All	All	3206	0	3083	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1083:LYS:HB3	1:A:1086:GLU:HB2	1.85	0.59
1:A:76:LEU:HD23	1:A:79:ARG:HD2	1.85	0.58
1:A:73:VAL:HA	1:A:76:LEU:HD12	1.90	0.53
1:A:196:VAL:HG13	4:A:2005:OLC:H17	1.94	0.50
1:A:255:ILE:HG22	1:A:273:MET:HE1	1.95	0.47
1:A:70:LEU:HA	1:A:73:VAL:HG22	2.00	0.44
1:A:242:PHE:O	1:A:247:MET:HB2	2.19	0.42
1:A:177:PRO:HG2	1:A:186:VAL:HG21	2.02	0.42
1:A:195:PHE:HE2	4:A:2005:OLC:H13A	1.85	0.41
1:A:44:VAL:HG21	1:A:292:PRO:HA	2.03	0.41
1:A:73:VAL:O	1:A:77:PRO:HD3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/423 (93%)	387 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/375 (82%)	297 (97%)	10 (3%)	33	63

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	ILE
1	A	45	LEU
1	A	71	LEU
1	A	103	LEU
1	A	134	LEU
1	A	205	ILE
1	A	1010	LEU
1	A	1014	LEU
1	A	1054	ASP
1	A	294	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	1025	GLN
1	A	1103	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	OLC	A	2009	-	5,5,24	0.59	0	4,4,25	0.54	0
4	OLC	A	2006	-	6,6,24	0.32	0	5,5,25	0.69	0
4	OLC	A	2004	-	11,11,24	0.81	0	12,12,25	1.03	1 (8%)
4	OLC	A	2010	-	9,9,24	0.86	0	10,10,25	1.17	1 (10%)
4	OLC	A	2011	-	10,10,24	0.39	0	9,9,25	0.59	0
4	OLC	A	2008	-	16,16,24	0.36	0	15,15,25	0.72	0
4	OLC	A	2003	-	12,12,24	0.78	0	13,13,25	1.02	1 (7%)
2	KNT	A	2001	-	39,40,40	0.59	1 (2%)	52,54,54	1.22	9 (17%)
5	OLA	A	2012	-	10,10,19	0.64	0	10,10,19	0.96	0
4	OLC	A	2005	-	24,24,24	0.62	0	25,25,25	0.90	1 (4%)
4	OLC	A	2007	-	20,20,24	0.67	0	21,21,25	0.89	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLC	A	2009	-	-	0/3/3/24	-
4	OLC	A	2006	-	-	4/4/4/24	-
4	OLC	A	2004	-	-	6/11/11/24	-
4	OLC	A	2010	-	-	6/9/9/24	-
4	OLC	A	2011	-	-	2/8/8/24	-
4	OLC	A	2008	-	-	9/14/14/24	-
4	OLC	A	2003	-	-	7/12/12/24	-
2	KNT	A	2001	-	-	2/16/20/20	0/5/5/5
5	OLA	A	2012	-	-	7/8/8/17	-
4	OLC	A	2005	-	-	15/24/24/24	-
4	OLC	A	2007	-	-	9/20/20/24	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	KNT	C7-C6	-2.01	1.40	1.44

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	KNT	C10-C5-C6	-3.96	116.18	119.52
4	A	2010	OLC	O20-C1-C2	3.02	120.31	111.15
2	A	2001	KNT	O2-C8-C7	-2.88	119.78	123.84
2	A	2001	KNT	C4-C5-C6	2.66	124.89	120.73
4	A	2007	OLC	O20-C1-C2	2.66	119.95	111.83
4	A	2005	OLC	O20-C1-C2	2.64	119.88	111.83
4	A	2004	OLC	O20-C1-C2	2.59	119.74	111.83
4	A	2003	OLC	O20-C1-C2	2.51	119.50	111.83
2	A	2001	KNT	O2-C10-C1	2.41	118.87	115.43
2	A	2001	KNT	O1-C6-C5	-2.40	118.00	121.38
2	A	2001	KNT	C10-O2-C8	2.34	123.20	118.94
2	A	2001	KNT	C5-C6-C7	2.34	121.26	116.55
2	A	2001	KNT	C9-N1-N2	-2.21	105.21	108.48
2	A	2001	KNT	N1-N2-N3	2.07	109.62	106.68

There are no chirality outliers.

All (67) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2003	OLC	C21-C22-C24-O25
4	A	2003	OLC	O20-C21-C22-C24
4	A	2003	OLC	O20-C21-C22-O23
4	A	2004	OLC	C21-C22-C24-O25
4	A	2005	OLC	C21-C22-C24-O25
4	A	2007	OLC	C21-C22-C24-O25
4	A	2007	OLC	O23-C22-C24-O25
4	A	2010	OLC	O20-C21-C22-C24
4	A	2004	OLC	C2-C1-O20-C21
4	A	2005	OLC	C2-C1-O20-C21
4	A	2010	OLC	C2-C1-O20-C21
4	A	2004	OLC	O19-C1-O20-C21
4	A	2005	OLC	O19-C1-O20-C21
4	A	2010	OLC	O19-C1-O20-C21
4	A	2007	OLC	C2-C1-O20-C21
4	A	2003	OLC	O19-C1-O20-C21
4	A	2007	OLC	O19-C1-O20-C21
4	A	2003	OLC	C2-C1-O20-C21
4	A	2010	OLC	O20-C21-C22-O23
4	A	2005	OLC	C1-C2-C3-C4
4	A	2005	OLC	C11-C12-C13-C14
4	A	2008	OLC	C13-C14-C15-C16
4	A	2011	OLC	C11-C12-C13-C14
5	A	2012	OLA	C4-C5-C6-C7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	2012	OLA	C5-C6-C7-C8
4	A	2005	OLC	O23-C22-C24-O25
4	A	2008	OLC	C4-C5-C6-C7
4	A	2004	OLC	C1-C2-C3-C4
4	A	2006	OLC	C4-C5-C6-C7
4	A	2003	OLC	C1-C2-C3-C4
4	A	2008	OLC	C11-C12-C13-C14
4	A	2005	OLC	C14-C15-C16-C17
4	A	2005	OLC	C10-C11-C12-C13
4	A	2005	OLC	C5-C6-C7-C8
4	A	2005	OLC	C6-C7-C8-C9
4	A	2008	OLC	C14-C15-C16-C17
4	A	2011	OLC	C10-C11-C12-C13
4	A	2007	OLC	C1-C2-C3-C4
4	A	2008	OLC	C3-C4-C5-C6
4	A	2008	OLC	C2-C3-C4-C5
4	A	2006	OLC	C5-C6-C7-C8
5	A	2012	OLA	C6-C7-C8-C9
4	A	2004	OLC	O23-C22-C24-O25
5	A	2012	OLA	C2-C3-C4-C5
4	A	2003	OLC	O23-C22-C24-O25
4	A	2007	OLC	C6-C7-C8-C9
4	A	2008	OLC	C6-C7-C8-C9
2	A	2001	KNT	C15-C16-C17-C18
4	A	2010	OLC	O20-C1-C2-C3
2	A	2001	KNT	C16-C17-C18-C19
4	A	2005	OLC	C9-C10-C11-C12
4	A	2004	OLC	C2-C3-C4-C5
4	A	2006	OLC	C2-C3-C4-C5
4	A	2010	OLC	O19-C1-C2-C3
5	A	2012	OLA	C3-C4-C5-C6
5	A	2012	OLA	O2-C1-C2-C3
4	A	2007	OLC	C2-C3-C4-C5
5	A	2012	OLA	O1-C1-C2-C3
4	A	2005	OLC	C7-C8-C9-C10
4	A	2008	OLC	C9-C10-C11-C12
4	A	2007	OLC	C7-C8-C9-C10
4	A	2005	OLC	O20-C1-C2-C3
4	A	2006	OLC	C3-C4-C5-C6
4	A	2007	OLC	C3-C4-C5-C6
4	A	2005	OLC	O19-C1-C2-C3
4	A	2005	OLC	C12-C13-C14-C15

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	2008	OLC	C5-C6-C7-C8



There are no ring outliers.




1 monomer is involved in 2 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2005	OLC	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/423 (93%)	0.76	21 (5%) 32 28	33, 64, 93, 110	2 (0%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1017	ILE	3.6
1	A	1087	ALA	3.5
1	A	1078	LEU	2.8
1	A	53	HIS	2.6
1	A	158	PHE	2.6
1	A	1007[A]	TRP	2.6
1	A	1084	VAL	2.6
1	A	263	GLU	2.5
1	A	1091	ALA	2.5
1	A	88	ILE	2.4
1	A	1048	LEU	2.3
1	A	1094	LEU	2.3
1	A	1093	GLN	2.3
1	A	50[A]	LYS	2.3
1	A	18	ILE	2.2
1	A	290	PHE	2.2
1	A	294	LEU	2.1
1	A	1044	THR	2.1
1	A	258	HIS	2.1
1	A	64	ASN	2.0
1	A	44	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	OLC	A	2009	6/25	0.71	0.17	34,36,37,37	0
4	OLC	A	2010	10/25	0.73	0.14	70,73,76,76	0
4	OLC	A	2003	13/25	0.78	0.16	74,76,78,78	0
4	OLC	A	2008	17/25	0.79	0.11	71,72,74,74	0
4	OLC	A	2011	11/25	0.79	0.15	64,65,66,66	0
4	OLC	A	2005	25/25	0.80	0.16	52,54,58,58	0
4	OLC	A	2007	21/25	0.80	0.15	64,66,71,71	0
4	OLC	A	2004	12/25	0.80	0.14	71,72,72,73	0
5	OLA	A	2012	11/20	0.86	0.14	57,59,62,62	0
4	OLC	A	2006	7/25	0.88	0.15	45,46,46,46	0
2	KNT	A	2001	36/36	0.91	0.12	47,48,58,58	0
3	NA	A	2002	1/1	0.95	0.09	52,52,52,52	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OLC A 2010:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 2003:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 2008:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around OLC A 2011:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around OLC A 2005:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 2007:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 2004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLA A 2012:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 2006:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around KNT A 2001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.