



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 02:19 AM EDT

PDB ID : 8S0H / pdb_00008s0h
Title : A fragment-based inhibitor of SHP2
Authors : Cleasby, A.; Price, A.
Deposited on : 2024-02-14
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	537	
1	B	537	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9030 atoms, of which 40 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	505	Total	C	N	O	S	0	2	0
			4098	2572	735	772	19			
1	B	500	Total	C	N	O	S	0	2	0
			4055	2547	726	762	20			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP Q06124
A	529	LEU	-	expression tag	UNP Q06124
A	530	GLU	-	expression tag	UNP Q06124
A	531	HIS	-	expression tag	UNP Q06124
A	532	HIS	-	expression tag	UNP Q06124
A	533	HIS	-	expression tag	UNP Q06124
A	534	HIS	-	expression tag	UNP Q06124
A	535	HIS	-	expression tag	UNP Q06124
A	536	HIS	-	expression tag	UNP Q06124
B	0	HIS	-	expression tag	UNP Q06124
B	529	LEU	-	expression tag	UNP Q06124
B	530	GLU	-	expression tag	UNP Q06124
B	531	HIS	-	expression tag	UNP Q06124
B	532	HIS	-	expression tag	UNP Q06124
B	533	HIS	-	expression tag	UNP Q06124
B	534	HIS	-	expression tag	UNP Q06124
B	535	HIS	-	expression tag	UNP Q06124
B	536	HIS	-	expression tag	UNP Q06124

- Molecule 2 is 5-(aminomethyl)-N-(3-chloranyl-1-methyl-indol-7-yl)-1,3-dihydroisindole-2-sulfonamide (CCD ID: A1H4Q) (formula: C₁₈H₁₉ClN₄O₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	C	Cl	H	N	O	S	0	0
			46	18	1	20	4	2	1		
2	B	1	Total	C	Cl	H	N	O	S	0	0
			46	18	1	20	4	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	403	Total	O	0	0
			403	403		
3	B	382	Total	O	0	0
			382	382		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.10Å 215.84Å 56.51Å 90.00° 97.05° 90.00°	Depositor
Resolution (Å)	107.92 – 1.99 107.92 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.7 (107.92-1.99) 99.7 (107.92-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.204 , 0.254 0.212 , 0.260	Depositor DCC
R_{free} test set	3751 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	30.8	Xtriage
Anisotropy	0.485	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9030	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1H4Q

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	0/4186	0.91	4/5645 (0.1%)
1	B	0.79	0/4139	0.91	2/5578 (0.0%)
All	All	0.78	0/8325	0.91	6/11223 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	458	HIS	CA-C-N	5.84	132.70	121.54
1	B	458	HIS	C-N-CA	5.84	132.70	121.54
1	A	458	HIS	CA-C-N	5.47	131.99	121.54
1	A	458	HIS	C-N-CA	5.47	131.99	121.54
1	A	299	GLU	CA-C-N	5.22	124.67	119.24
1	A	299	GLU	C-N-CA	5.22	124.67	119.24

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	265	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4098	0	4025	20	0
1	B	4055	0	3985	23	0
2	A	26	20	0	0	0
2	B	26	20	0	1	0
3	A	403	0	0	3	0
3	B	382	0	0	4	0
All	All	8990	40	8010	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:171[A]:MET:HE3	1:B:171[A]:MET:HA	1.68	0.73
1:B:129:LYS:NZ	3:B:707:HOH:O	2.39	0.55
1:B:171[A]:MET:HE3	1:B:171[A]:MET:CA	2.37	0.53
1:A:520:HIS:O	1:A:524:THR:HG23	2.09	0.52
1:A:366:LYS:HE3	3:A:945:HOH:O	2.11	0.51
1:B:483:GLY:O	1:B:486:CYS:SG	2.63	0.51
1:B:518:VAL:O	1:B:522:ILE:HG12	2.11	0.50
1:A:131:LYS:HB2	1:A:156:ASP:OD2	2.12	0.49
1:A:11:ILE:HD11	1:A:19:LEU:HD12	1.95	0.49
1:A:134:SER:HA	1:A:214:GLN:O	2.13	0.49
1:A:271:GLN:NE2	3:A:703:HOH:O	2.34	0.49
1:B:382:VAL:HG12	3:B:1071:HOH:O	2.12	0.48
1:B:222:ASN:ND2	1:B:484:VAL:O	2.45	0.48
1:B:148:VAL:HG22	1:B:171[A]:MET:HE1	1.95	0.48
1:A:278:ARG:HD3	1:A:279:TYR:CE2	2.49	0.48
1:B:28:SER:HA	1:B:100:TYR:O	2.14	0.47
1:B:118:SER:OG	1:B:141:GLN:OE1	2.33	0.47
1:A:255:GLN:OE1	1:A:498:ARG:NH1	2.48	0.47
1:B:520:HIS:O	1:B:524:THR:HG23	2.16	0.46
1:B:30:LEU:HD12	1:B:30:LEU:C	2.41	0.46
1:A:521:TYR:O	1:A:525:LEU:HD13	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:LYS:HG3	1:A:281:ASN:OD1	2.16	0.45
1:B:495:GLN:HG2	2:B:601:A1H4Q:CL5	2.54	0.45
1:B:222:ASN:HD22	1:B:224:ALA:H	1.65	0.44
1:B:120:LYS:HA	1:B:123:GLU:HB2	1.99	0.44
1:B:134:SER:HA	1:B:214:GLN:O	2.18	0.44
1:A:369:LYS:HE2	1:A:371:TRP:NE1	2.33	0.44
1:A:524:THR:HG22	3:A:806:HOH:O	2.17	0.44
1:A:193:LEU:C	1:A:193:LEU:HD23	2.42	0.43
1:B:421:ARG:NH2	3:B:738:HOH:O	2.51	0.43
1:B:235:LYS:O	1:B:245:GLN:N	2.52	0.43
1:A:80:TYR:OH	1:A:280:LYS:NZ	2.51	0.42
1:A:90:GLU:HB3	1:A:96:ILE:HD11	2.01	0.42
1:B:410:ASN:ND2	3:B:729:HOH:O	2.48	0.42
1:B:109:SER:HB2	1:B:256:GLN:HG3	2.01	0.42
1:A:28:SER:HA	1:A:100:TYR:O	2.20	0.42
1:B:11:ILE:CD1	1:B:19:LEU:HD12	2.50	0.41
1:A:178:LYS:HE2	1:A:188:ASP:OD1	2.20	0.41
1:A:56:ILE:HG12	1:A:65:LEU:HD13	2.01	0.41
1:B:118:SER:HB2	1:B:121:GLU:OE1	2.21	0.41
1:A:80:TYR:CD1	1:A:281:ASN:ND2	2.89	0.40
1:A:203:VAL:HG12	1:A:204:GLU:O	2.22	0.40
1:B:290:VAL:HG11	1:B:344:MET:HG3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	499/537 (93%)	485 (97%)	12 (2%)	2 (0%)	30	27
1	B	490/537 (91%)	476 (97%)	12 (2%)	2 (0%)	30	27
All	All	989/1074 (92%)	961 (97%)	24 (2%)	4 (0%)	30	27

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	459	CYS
1	B	459	CYS
1	A	505	VAL
1	B	505	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	451/479 (94%)	447 (99%)	4 (1%)	70	78
1	B	446/479 (93%)	437 (98%)	9 (2%)	48	54
All	All	897/958 (94%)	884 (99%)	13 (1%)	61	66

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	ARG
1	A	151	VAL
1	A	335	GLN
1	A	458	HIS
1	B	95	VAL
1	B	151	VAL
1	B	259[A]	CYS
1	B	259[B]	CYS
1	B	280	LYS
1	B	313	GLU
1	B	481	GLU
1	B	508	GLU
1	B	525	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	48	ASN
1	A	116	HIS
1	A	141	GLN
1	A	196	HIS
1	B	196	HIS
1	B	214	GLN
1	B	222	ASN
1	B	293	HIS
1	B	408	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A1H4Q	B	601	-	28,29,29	0.30	0	37,44,44	0.73	1 (2%)
2	A1H4Q	A	601	-	28,29,29	0.33	0	37,44,44	0.78	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1H4Q	B	601	-	-	1/11/21/21	0/4/4/4
2	A1H4Q	A	601	-	-	2/11/21/21	0/4/4/4

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	A1H4Q	C6-C26-N2	3.21	108.28	107.18
2	B	601	A1H4Q	C6-C26-N2	2.05	107.88	107.18

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	A1H4Q	C25-N15-S12-O13
2	B	601	A1H4Q	C25-N15-S12-O13
2	A	601	A1H4Q	C16-N15-S12-O13

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	A1H4Q	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	505/537 (94%)	0.81	66 (13%) 7 6	17, 40, 80, 108	2 (0%)
1	B	500/537 (93%)	0.88	72 (14%) 6 5	19, 42, 79, 110	2 (0%)
All	All	1005/1074 (93%)	0.84	138 (13%) 6 6	17, 41, 80, 110	4 (0%)

All (138) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	484	VAL	5.1
1	B	301	VAL	4.8
1	A	484	VAL	4.7
1	A	301	VAL	4.6
1	A	486	CYS	4.6
1	A	224	ALA	4.5
1	B	248	TRP	4.4
1	B	486	CYS	4.3
1	A	117	LEU	4.1
1	A	526	GLN	4.0
1	B	36	SER	4.0
1	B	275	ASN	3.9
1	A	177	LEU	3.9
1	A	66	TYR	3.8
1	A	525	LEU	3.7
1	A	206	LEU	3.7
1	B	177	LEU	3.7
1	B	522	ILE	3.7
1	A	86	GLY	3.6
1	A	323	PRO	3.6
1	B	206	LEU	3.5
1	A	95	VAL	3.5
1	A	210	LEU	3.5
1	B	529	LEU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	86	GLY	3.5
1	A	85	HIS	3.4
1	B	37	ASN	3.4
1	B	143	HIS	3.3
1	A	91	LYS	3.2
1	B	226	ILE	3.2
1	A	92	ASN	3.2
1	B	409	GLY	3.2
1	A	248	TRP	3.2
1	B	117	LEU	3.2
1	B	155	ASP	3.0
1	A	311	MET	3.0
1	B	85	HIS	3.0
1	A	38	PRO	3.0
1	B	530	GLU	3.0
1	A	203	VAL	2.9
1	A	300	PRO	2.9
1	A	212	LEU	2.9
1	A	48	ASN	2.9
1	A	226	ILE	2.9
1	A	294	ASP	2.8
1	A	59	THR	2.8
1	A	181	VAL	2.8
1	B	181	VAL	2.8
1	A	299	GLU	2.8
1	B	109	SER	2.8
1	A	527	ARG	2.8
1	A	144	PRO	2.8
1	B	311	MET	2.8
1	A	275[A]	ASN	2.7
1	B	93	GLY	2.7
1	A	524	THR	2.7
1	A	485	ASP	2.7
1	B	485	ASP	2.7
1	A	25	VAL	2.7
1	A	222	ASN	2.6
1	A	143	HIS	2.6
1	B	221	ILE	2.6
1	A	93	GLY	2.6
1	A	157	LYS	2.6
1	A	209	VAL	2.6
1	A	450	MET	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	35	LYS	2.5
1	B	224	ALA	2.5
1	B	89	LYS	2.5
1	A	52	THR	2.5
1	B	426	HIS	2.5
1	B	203	VAL	2.5
1	B	120	LYS	2.5
1	B	38	PRO	2.5
1	A	145	GLY	2.4
1	B	172	ILE	2.4
1	B	476	ILE	2.4
1	B	324	LYS	2.4
1	B	84	HIS	2.4
1	B	293	HIS	2.4
1	A	34	SER	2.4
1	A	512	ARG	2.4
1	B	313	GLU	2.4
1	A	142	SER	2.4
1	B	144	PRO	2.4
1	B	298	ASN	2.4
1	B	524	THR	2.3
1	B	526	GLN	2.3
1	A	410	ASN	2.3
1	B	482	LYS	2.3
1	B	205	THR	2.3
1	A	10	ASN	2.3
1	B	223	ALA	2.3
1	B	450	MET	2.3
1	B	197	TYR	2.3
1	B	142	SER	2.3
1	A	437	ASP	2.3
1	B	151	VAL	2.3
1	B	167	VAL	2.3
1	B	235	LYS	2.3
1	B	88	LEU	2.2
1	A	522	ILE	2.2
1	B	96	ILE	2.2
1	A	120	LYS	2.2
1	B	527	ARG	2.2
1	B	225	GLU	2.2
1	A	483	GLY	2.2
1	B	231	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	14	VAL	2.2
1	B	336	ASN	2.1
1	A	11	ILE	2.1
1	B	95	VAL	2.1
1	B	168	THR	2.1
1	A	297	PRO	2.1
1	B	171[A]	MET	2.1
1	B	141	GLN	2.1
1	A	488	ILE	2.1
1	A	207	GLY	2.1
1	B	207	GLY	2.1
1	A	90	GLU	2.1
1	B	125	LEU	2.1
1	B	525	LEU	2.1
1	A	172	ILE	2.1
1	A	296	ASP	2.1
1	A	36	SER	2.1
1	B	34	SER	2.1
1	B	108	THR	2.1
1	B	271	GLN	2.1
1	A	482	LYS	2.0
1	A	196	HIS	2.0
1	B	483	GLY	2.0
1	A	518	VAL	2.0
1	B	265	ARG	2.0
1	A	171	MET	2.0
1	B	11	ILE	2.0
1	A	119	GLY	2.0
1	B	209	VAL	2.0
1	B	230	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	A1H4Q	B	601	26/26	0.93	0.09	25,32,38,40	0
2	A1H4Q	A	601	26/26	0.94	0.09	23,31,36,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1H4Q B 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1H4Q A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.