



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:12 AM UTC

PDB ID : 4S1I / pdb_00004s1i
Title : Pyridoxal Kinase of Entamoeba histolytica with PLP
Authors : Tarique, K.F.; Devi, S.; Abdul Rehman, S.A.; Gourinath, S.
Deposited on : 2015-01-14
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	287	10% 86% 9% ...
2	B	287	7% 84% 11% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4628 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridoxal kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	0	0
			2182	1404	351	417	10			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	LEU	-	expression tag	UNP C4LVZ4
A	281	GLU	-	expression tag	UNP C4LVZ4
A	282	HIS	-	expression tag	UNP C4LVZ4
A	283	HIS	-	expression tag	UNP C4LVZ4
A	284	HIS	-	expression tag	UNP C4LVZ4
A	285	HIS	-	expression tag	UNP C4LVZ4
A	286	HIS	-	expression tag	UNP C4LVZ4
A	287	HIS	-	expression tag	UNP C4LVZ4


- Molecule 2 is a protein called Pyridoxal kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	275	Total	C	N	O	S	0	0	0
			2151	1387	344	408	12			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	280	LEU	-	expression tag	UNP C4LVZ4
B	281	GLU	-	expression tag	UNP C4LVZ4
B	282	HIS	-	expression tag	UNP C4LVZ4
B	283	HIS	-	expression tag	UNP C4LVZ4
B	284	HIS	-	expression tag	UNP C4LVZ4
B	285	HIS	-	expression tag	UNP C4LVZ4
B	286	HIS	-	expression tag	UNP C4LVZ4
B	287	HIS	-	expression tag	UNP C4LVZ4

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
3	B	1	Total	C	N	O	P	0	0
			16	8	1	6	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	110	Total	O	0	0
			110	110		
5	B	151	Total	O	0	0
			151	151		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyridoxal kinase



• Molecule 2: Pyridoxal kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	78.17Å 79.01Å 90.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.60 50.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-1.60) 99.3 (50.00-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0102	Depositor
R, R_{free}	0.195 , 0.225 0.199 , 0.230	Depositor DCC
R_{free} test set	3764 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	23.8	Xtriage
Anisotropy	0.444	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.006 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4628	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CME, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.37	9/2219 (0.4%)	1.21	4/3013 (0.1%)
2	B	1.55	15/2176 (0.7%)	1.34	8/2952 (0.3%)
All	All	1.47	24/4395 (0.5%)	1.28	12/5965 (0.2%)

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	187	ILE	C-N	-8.62	1.21	1.33
2	B	136	TYR	N-CA	7.62	1.56	1.46
2	B	24	ILE	N-CA	-7.07	1.38	1.46
2	B	173	ILE	CA-C	6.97	1.60	1.52
2	B	142	PRO	N-CA	6.46	1.54	1.47
2	B	11	TYR	C-O	6.20	1.31	1.24
2	B	146	GLU	CA-C	-6.04	1.45	1.52
2	B	187	ILE	N-CA	6.01	1.57	1.46
1	A	102	ASP	N-CA	5.92	1.53	1.46
2	B	237	ALA	CA-C	5.86	1.60	1.52
2	B	191	CYS	CA-C	-5.72	1.45	1.52
1	A	2	THR	C-O	-5.60	1.16	1.23
1	A	187	ILE	C-O	-5.56	1.18	1.24
2	B	20	ARG	CZ-NH2	-5.48	1.26	1.33
1	A	20	ARG	CA-C	5.45	1.59	1.52
1	A	41	LEU	N-CA	-5.42	1.39	1.45
1	A	181	ILE	C-O	-5.41	1.18	1.24
1	A	26	LEU	N-CA	5.36	1.52	1.46
2	B	241	ALA	CA-CB	-5.24	1.45	1.53
1	A	161	ALA	N-CA	-5.21	1.40	1.46
1	A	146	GLU	N-CA	5.12	1.52	1.46
2	B	151	THR	CA-C	5.09	1.59	1.52
2	B	192	SER	N-CA	-5.04	1.39	1.46
2	B	228	ILE	C-O	5.01	1.29	1.24

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	183	GLU	N-CA-C	6.46	118.17	107.20
2	B	187	ILE	O-C-N	-6.14	113.18	123.00
1	A	279	ILE	CB-CA-C	5.94	119.35	110.98
2	B	32	GLN	CA-CB-CG	-5.74	102.62	114.10
2	B	27	ASP	O-C-N	-5.68	116.10	122.12
2	B	179	THR	CA-C-N	-5.44	113.19	122.33
2	B	179	THR	C-N-CA	-5.44	113.19	122.33
2	B	145	THR	N-CA-C	-5.37	105.50	111.36
2	B	187	ILE	N-CA-C	5.33	125.93	111.00
1	A	73	ASP	N-CA-C	5.09	119.18	112.92
2	B	253	VAL	N-CA-C	5.06	115.60	108.36
1	A	20	ARG	CG-CD-NE	-5.02	100.95	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2182	0	2233	14	0
2	B	2151	0	2210	14	0
3	A	16	0	7	0	0
3	B	16	0	7	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	110	0	0	0	0
5	B	151	0	0	2	0
All	All	4628	0	4457	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:GLN:HE22	2:B:259:ASN:ND2	1.63	0.95
1:A:32:GLN:NE2	2:B:259:ASN:HD21	1.64	0.95
1:A:79:LEU:HD21	1:A:97:VAL:HG22	1.76	0.67
2:B:79:LEU:HD21	2:B:97:VAL:HG22	1.79	0.64
2:B:255:THR:HG22	2:B:255:THR:O	2.05	0.56
1:A:80:VAL:HG21	1:A:225:LEU:HD22	1.89	0.55
1:A:177:LEU:HD11	1:A:189:LEU:HD21	1.89	0.53
1:A:79:LEU:CD2	1:A:97:VAL:HG22	2.40	0.51
1:A:80:VAL:CG2	1:A:225:LEU:HD22	2.41	0.50
2:B:109:LEU:C	2:B:109:LEU:HD23	2.36	0.50
2:B:173:ILE:HD13	5:B:417:HOH:O	2.10	0.50
2:B:177:LEU:HD11	2:B:189:LEU:HD21	1.94	0.49
1:A:109:LEU:HD23	1:A:109:LEU:C	2.39	0.48
2:B:80:VAL:HG21	2:B:225:LEU:HD22	1.94	0.47
1:A:127:GLN:HB3	1:A:149:PHE:CE2	2.49	0.47
2:B:79:LEU:CD2	2:B:97:VAL:HG22	2.45	0.46
2:B:80:VAL:CG2	2:B:225:LEU:HD22	2.45	0.46
2:B:235:GLU:HG2	5:B:419:HOH:O	2.14	0.46
2:B:255:THR:O	2:B:255:THR:CG2	2.64	0.46
1:A:127:GLN:HB3	1:A:149:PHE:CZ	2.51	0.45
1:A:182:LYS:HA	1:A:182:LYS:HD3	1.40	0.43
2:B:189:LEU:C	2:B:189:LEU:HD23	2.44	0.42
1:A:183:GLU:O	1:A:184:GLY:C	2.62	0.42
2:B:103:ASN:O	2:B:103:ASN:CG	2.63	0.42
1:A:110:CYS:HB3	1:A:140:ILE:HG22	2.01	0.41
1:A:220:LEU:HB2	1:A:245:LEU:CD2	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/287 (97%)	272 (98%)	6 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	270/287 (94%)	264 (98%)	6 (2%)	0	100	100
All	All	548/574 (96%)	536 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/260 (97%)	244 (97%)	8 (3%)	34	12
2	B	248/259 (96%)	245 (99%)	3 (1%)	63	43
All	All	500/519 (96%)	489 (98%)	11 (2%)	45	22

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	43	ASN
1	A	58	ASN
1	A	119	LYS
1	A	124	SER
1	A	182	LYS
1	A	187	ILE
1	A	245	LEU
2	B	43	ASN
2	B	197	LEU
2	B	245	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	262	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CME	B	263	2	8,9,10	1.11	1 (12%)	6,9,11	1.60	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CME	B	263	2	-	1/5/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	263	CME	CB-SG	-2.19	1.74	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	263	CME	CB-SG-SD	2.16	109.46	103.86
2	B	263	CME	CE-SD-SG	2.16	112.93	103.46

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	263	CME	SD-CE-CZ-OH

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	PLP	A	301	-	16,16,16	2.14	4 (25%)	20,23,23	2.72	9 (45%)
3	PLP	B	302	-	16,16,16	2.43	6 (37%)	20,23,23	2.95	8 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PLP	A	301	-	-	2/8/8/8	0/1/1/1
3	PLP	B	302	-	-	3/8/8/8	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	PLP	C6-N1	-5.75	1.22	1.34
3	A	301	PLP	C6-N1	-5.36	1.23	1.34
3	B	302	PLP	O3-C3	-5.00	1.25	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	PLP	O3-C3	-3.69	1.28	1.36
3	B	302	PLP	C2A-C2	-2.93	1.45	1.50
3	A	301	PLP	C5A-C5	-2.46	1.44	1.50
3	B	302	PLP	C3-C2	2.27	1.43	1.41
3	A	301	PLP	P-O3P	-2.15	1.46	1.54
3	B	302	PLP	C4-C3	2.05	1.44	1.41
3	B	302	PLP	P-O2P	-2.03	1.47	1.54

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	PLP	C3-C4-C5	-8.70	111.29	118.28
3	A	301	PLP	C3-C4-C5	-7.35	112.38	118.28
3	B	302	PLP	O4P-C5A-C5	-6.62	96.94	109.36
3	A	301	PLP	O4P-C5A-C5	-5.45	99.14	109.36
3	A	301	PLP	C5A-C5-C6	-4.36	112.25	119.36
3	B	302	PLP	O3-C3-C2	3.16	124.13	117.58
3	A	301	PLP	O3P-P-O4P	-2.98	98.89	106.67
3	B	302	PLP	O3P-P-O1P	2.93	122.25	110.83
3	B	302	PLP	O4P-P-O1P	-2.91	98.57	106.44
3	A	301	PLP	C6-C5-C4	2.78	122.99	118.21
3	B	302	PLP	O2P-P-O4P	2.62	113.50	106.67
3	A	301	PLP	O3-C3-C2	2.60	122.97	117.58
3	B	302	PLP	C6-C5-C4	2.52	122.54	118.21
3	B	302	PLP	O3P-P-O4P	-2.48	100.19	106.67
3	A	301	PLP	C2A-C2-C3	-2.28	118.13	120.80
3	A	301	PLP	O3P-P-O2P	2.22	116.13	107.80
3	A	301	PLP	C4-C3-C2	-2.14	118.94	120.14

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	PLP	C5A-O4P-P-O3P
3	B	302	PLP	C5A-O4P-P-O2P
3	B	302	PLP	C5A-O4P-P-O3P
3	B	302	PLP	C5A-O4P-P-O1P
3	A	301	PLP	C3-C4-C4A-O4A

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	280/287 (97%)	0.68	30 (10%)	11 11	18, 32, 62, 91	0
2	B	274/287 (95%)	0.50	21 (7%)	19 20	17, 27, 49, 71	0
All	All	554/574 (96%)	0.59	51 (9%)	14 14	17, 30, 55, 91	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	185	ASN	6.0
1	A	197	LEU	5.4
1	A	186	ASP	5.3
1	A	254	GLY	5.0
1	A	213	PHE	4.3
1	A	187	ILE	4.1
2	B	209	ILE	3.8
1	A	1	MET	3.8
1	A	280	LEU	3.6
2	B	197	LEU	3.5
2	B	187	ILE	3.4
2	B	156	ASN	3.3
1	A	124	SER	3.3
2	B	102	ASP	3.2
2	B	213	PHE	3.1
2	B	1	MET	3.1
1	A	118	GLY	3.1
2	B	280	LEU	3.0
1	A	116	ASP	3.0
1	A	253	VAL	2.9
1	A	209	ILE	2.9
2	B	206	ILE	2.8
1	A	182	LYS	2.8
1	A	102	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	184	GLY	2.8
2	B	207	PRO	2.7
2	B	157	SER	2.7
2	B	154	GLU	2.7
2	B	181	ILE	2.7
2	B	103	ASN	2.6
2	B	253	VAL	2.6
1	A	255	THR	2.5
1	A	263	CYS	2.5
1	A	119	LYS	2.4
1	A	149	PHE	2.4
2	B	155	VAL	2.4
1	A	256	ALA	2.4
2	B	210	GLU	2.3
1	A	99	ARG	2.3
1	A	104	LYS	2.3
1	A	100	ILE	2.2
1	A	250	ARG	2.2
1	A	180	SER	2.2
2	B	117	ASN	2.2
2	B	153	LEU	2.1
1	A	189	LEU	2.1
2	B	265	PRO	2.1
1	A	278	TYR	2.1
2	B	158	VAL	2.1
1	A	120	MET	2.0
1	A	208	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CME	B	263	10/11	0.90	0.14	25,26,46,53	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PLP	B	302	16/16	0.79	0.15	23,46,69,82	0
3	PLP	A	301	16/16	0.85	0.14	23,46,85,86	0
4	MG	B	301	1/1	0.98	0.12	25,25,25,25	0
4	MG	A	302	1/1	0.99	0.11	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.