



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:31 AM UTC

PDB ID : 6S1Z / pdb_00006s1z
Title : Crystal structure of Anopheles gambiae AnoACE2 in complex with fosinoprilat
Authors : Cozier, G.E.; Acharya, K.R.; Cashman, J.S.
Deposited on : 2019-06-19
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	621	19% 80% 14% 6%
2	B	3	100%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9558 atoms, of which 4687 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Angiotensin-converting enzyme.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	585	9397	3067	4621	802	888	19	0	0	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	639	LYS	-	expression tag	UNP A0NFU8
A	640	HIS	-	expression tag	UNP A0NFU8
A	641	HIS	-	expression tag	UNP A0NFU8
A	642	HIS	-	expression tag	UNP A0NFU8
A	643	HIS	-	expression tag	UNP A0NFU8
A	644	HIS	-	expression tag	UNP A0NFU8
A	645	HIS	-	expression tag	UNP A0NFU8
A	646	HIS	-	expression tag	UNP A0NFU8
A	647	HIS	-	expression tag	UNP A0NFU8

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O				
2	B	3	73	22	34	2	15		0	0	0

- Molecule 3 is fosinoprilat (CCD ID: KS8) (formula: C₂₃H₃₄NO₅P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	0	0
			62	23	32	1	5	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	25	Total	O	0	0
			25	25		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Angiotensin-converting enzyme



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	166.72Å 166.72Å 117.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	72.19 – 2.50 72.19 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (72.19-2.50) 100.0 (72.19-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.17 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.294 , 0.337 0.297 , 0.338	Depositor DCC
R_{free} test set	1695 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	54.3	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 63.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.36$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	9558	wwPDB-VP
Average B, all atoms (Å ²)	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, KS8, BMA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/4905	0.27	0/6648

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4776	4621	4622	53	0
2	B	39	34	34	0	0
3	A	30	32	0	1	0
4	A	1	0	0	0	0
5	A	25	0	0	4	0
All	All	4871	4687	4656	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:GLU:OE2	1:A:112:ARG:NH2	2.18	0.76
1:A:329:THR:HG21	1:A:392:LEU:HD13	1.67	0.76
1:A:351:VAL:O	1:A:368:LYS:NZ	2.19	0.75
1:A:373:VAL:O	5:A:801:HOH:O	2.05	0.74
1:A:360:PHE:O	5:A:802:HOH:O	2.07	0.71
1:A:617:GLY:O	5:A:803:HOH:O	2.10	0.68
1:A:535:ARG:NH2	1:A:590:GLU:OE2	2.29	0.66
1:A:289:PHE:O	5:A:804:HOH:O	2.15	0.63
1:A:389:GLN:OE1	1:A:393:GLN:NE2	2.30	0.62
1:A:315:MET:HE1	1:A:373:VAL:HA	1.85	0.59
1:A:523:TYR:OH	3:A:704:KS8:O16	2.20	0.59
1:A:194:ARG:NH2	1:A:503:GLU:O	2.36	0.57
1:A:331:LEU:HD21	1:A:335:PHE:CD1	2.41	0.56
1:A:127:ASP:OD1	1:A:128:GLN:N	2.40	0.54
1:A:507:ASP:N	1:A:508:PRO:CD	2.71	0.53
1:A:425:LYS:HD2	1:A:557:ASP:HB3	1.91	0.53
1:A:331:LEU:HD23	1:A:332:PRO:O	2.09	0.53
1:A:424:PRO:O	1:A:428:LYS:N	2.37	0.52
1:A:301:VAL:HG12	1:A:301:VAL:O	2.10	0.51
1:A:124:LEU:HD23	1:A:210:ARG:HD2	1.92	0.50
1:A:444:ILE:O	1:A:448:TYR:N	2.40	0.50
1:A:420:SER:O	1:A:423:THR:OG1	2.24	0.48
1:A:303:ASP:OD1	1:A:303:ASP:N	2.47	0.48
1:A:221:LEU:HD21	1:A:469:TRP:CH2	2.49	0.48
1:A:111:ARG:NH1	1:A:577:SER:O	2.48	0.47
1:A:233:VAL:HG13	1:A:524:PHE:CE2	2.49	0.47
1:A:446:GLN:NE2	1:A:555:ASN:OD1	2.47	0.47
1:A:375:MET:HE2	1:A:379:PHE:CE2	2.49	0.47
1:A:45:VAL:O	1:A:49:GLU:N	2.47	0.46
1:A:200:TYR:OH	1:A:517:ASP:OD1	2.22	0.46
1:A:580:TRP:N	1:A:581:PRO:CD	2.79	0.46
1:A:297:LYS:HG2	1:A:436:TYR:HD1	1.80	0.46
1:A:243:LEU:HD12	1:A:243:LEU:N	2.32	0.44
1:A:329:THR:CG2	1:A:392:LEU:HD13	2.42	0.44
1:A:535:ARG:HG3	1:A:544:TYR:CE2	2.52	0.44
1:A:331:LEU:HD22	1:A:336:TRP:CE2	2.52	0.44
1:A:365:VAL:HB	1:A:388:ILE:HG23	2.00	0.44
1:A:296:LYS:NZ	1:A:439:ASP:OD2	2.37	0.44
1:A:300:ASP:OD1	1:A:301:VAL:N	2.51	0.44
1:A:554:ASN:ND2	1:A:601:TYR:O	2.47	0.43
1:A:281:GLN:OE1	1:A:511:LYS:NZ	2.42	0.43
1:A:430:VAL:O	1:A:432:LEU:N	2.47	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:322:PHE:CE1	1:A:417:LEU:HD22	2.54	0.42
1:A:460:PHE:CE1	1:A:464:LEU:HD11	2.54	0.42
1:A:547:GLY:O	1:A:548:ASP:C	2.62	0.42
1:A:111:ARG:O	1:A:115:LYS:N	2.45	0.42
1:A:375:MET:HE2	1:A:379:PHE:HE2	1.83	0.42
1:A:612:GLU:OE2	1:A:615:ARG:NH2	2.53	0.42
1:A:424:PRO:HB3	1:A:433:LEU:HD21	2.01	0.41
1:A:409:PHE:O	1:A:413:VAL:HG23	2.20	0.41
1:A:149:CYS:HA	1:A:157:CYS:HA	2.03	0.41
1:A:508:PRO:N	1:A:509:PRO:CD	2.84	0.41
1:A:286:ILE:C	1:A:286:ILE:HD12	2.45	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	581/621 (94%)	538 (93%)	42 (7%)	1 (0%)	43 63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	549	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	505/538 (94%)	502 (99%)	3 (1%)	78 91

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	166	THR
1	A	303	ASP
1	A	410	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	59	GLN
1	A	142	ASN
1	A	426	HIS
1	A	446	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	0.50	0	17,19,21	0.61	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	2	2	14,14,15	0.29	0	17,19,21	0.42	0
2	BMA	B	3	2	11,11,12	0.63	0	15,15,17	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	O5-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	KS8	A	704	4	29,32,32	3.12	9 (31%)	34,44,44	1.17	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KS8	A	704	4	-	5/25/45/45	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	704	KS8	C20-C21	-8.56	1.38	1.53
3	A	704	KS8	C18-N17	-8.48	1.34	1.47
3	A	704	KS8	C21-N17	5.55	1.57	1.47
3	A	704	KS8	C15-N17	5.45	1.51	1.35
3	A	704	KS8	C14-C15	4.80	1.57	1.51
3	A	704	KS8	C18-C19	3.12	1.57	1.53
3	A	704	KS8	O24-C22	-3.10	1.20	1.30
3	A	704	KS8	C07-C08	3.03	1.60	1.51
3	A	704	KS8	C05-C04	2.25	1.59	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	704	KS8	C18-N17-C21	-3.77	106.83	111.83

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	704	KS8	C05-C04-P02-O03
3	A	704	KS8	C04-C05-C06-C07
3	A	704	KS8	C15-C14-P02-O01
3	A	704	KS8	C05-C04-P02-O01
3	A	704	KS8	P02-C14-C15-N17


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	704	KS8	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	585/621 (94%)	1.24	121 (20%) 2 2	26, 95, 146, 182	0

All (121) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	335	PHE	6.4
1	A	573	ALA	6.1
1	A	298	LEU	5.2
1	A	362	ILE	5.1
1	A	399	VAL	4.9
1	A	332	PRO	4.8
1	A	537	ALA	4.7
1	A	365	VAL	4.7
1	A	116	LEU	4.6
1	A	113	ILE	4.5
1	A	360	PHE	4.3
1	A	109	LEU	4.3
1	A	117	ALA	4.3
1	A	229	PHE	4.2
1	A	616	LEU	4.2
1	A	47	TRP	4.1
1	A	630	SER	3.9
1	A	299	LEU	3.8
1	A	536	ALA	3.7
1	A	357	TRP	3.6
1	A	45	VAL	3.5
1	A	83	ALA	3.5
1	A	444	ILE	3.5
1	A	568	LEU	3.5
1	A	301	VAL	3.3
1	A	126	ALA	3.3
1	A	422	SER	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	293	PHE	3.3
1	A	122	ALA	3.3
1	A	339	SER	3.3
1	A	361	ALA	3.2
1	A	316	PHE	3.2
1	A	533	PHE	3.2
1	A	250	TYR	3.2
1	A	552	THR	3.1
1	A	461	ALA	3.1
1	A	396	HIS	3.1
1	A	439	ASP	3.1
1	A	599	LEU	3.1
1	A	129	PHE	3.0
1	A	583	ALA	3.0
1	A	52	TYR	3.0
1	A	561	SER	3.0
1	A	41	ILE	2.9
1	A	313	ILE	2.9
1	A	476	ILE	2.9
1	A	558	ILE	2.9
1	A	328	MET	2.9
1	A	336	TRP	2.9
1	A	119	LEU	2.9
1	A	310	TYR	2.9
1	A	467	TYR	2.9
1	A	112	ARG	2.9
1	A	430	VAL	2.9
1	A	528	VAL	2.9
1	A	96	LEU	2.9
1	A	341	LEU	2.9
1	A	323	PHE	2.9
1	A	338	LYS	2.8
1	A	364	ASP	2.8
1	A	436	TYR	2.8
1	A	416	VAL	2.7
1	A	331	LEU	2.7
1	A	253	TYR	2.7
1	A	415	ASP	2.7
1	A	388	ILE	2.7
1	A	424	PRO	2.7
1	A	609	LEU	2.7
1	A	405	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	378	PHE	2.6
1	A	340	ILE	2.6
1	A	133	LEU	2.6
1	A	432	LEU	2.6
1	A	55	THR	2.6
1	A	524	PHE	2.6
1	A	460	PHE	2.5
1	A	442	VAL	2.5
1	A	334	THR	2.5
1	A	358	ASP	2.4
1	A	553	LEU	2.4
1	A	580	TRP	2.4
1	A	433	LEU	2.4
1	A	238	GLU	2.4
1	A	404	GLY	2.4
1	A	401	PHE	2.4
1	A	593	MET	2.4
1	A	88	VAL	2.4
1	A	555	ASN	2.4
1	A	124	LEU	2.3
1	A	366	ARG	2.3
1	A	587	LEU	2.3
1	A	107	ALA	2.3
1	A	311	THR	2.3
1	A	64	TRP	2.3
1	A	417	LEU	2.3
1	A	44	ILE	2.2
1	A	322	PHE	2.2
1	A	312	PRO	2.2
1	A	48	ILE	2.2
1	A	480	GLU	2.2
1	A	564	ALA	2.2
1	A	562	THR	2.2
1	A	36	PRO	2.1
1	A	123	ALA	2.1
1	A	597	ALA	2.1
1	A	431	GLY	2.1
1	A	425	LYS	2.1
1	A	359	PHE	2.1
1	A	435	ASP	2.1
1	A	241	ARG	2.1
1	A	243	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	427	LEU	2.1
1	A	302	THR	2.1
1	A	297	LYS	2.1
1	A	337	ASP	2.1
1	A	379	PHE	2.0
1	A	121	TYR	2.0
1	A	391	TYR	2.0
1	A	544	TYR	2.0
1	A	601	TYR	2.0
1	A	154	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BMA	B	3	11/12	0.77	0.10	74,113,133,137	0
2	NAG	B	2	14/15	0.79	0.12	83,103,129,135	0
2	NAG	B	1	14/15	0.80	0.13	71,93,116,133	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	KS8	A	704	30/30	0.89	0.15	63,94,116,118	0
4	ZN	A	705	1/1	0.98	0.04	87,87,87,87	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around KS8 A 704:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.