



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:04 AM UTC

PDB ID : 3S2D / pdb_00003s2d
Title : RNA Polymerase II Initiation Complex with a 5-nt RNA containing a 5Br-U
Authors : Liu, X.; Bushnell, D.A.; Silva, D.A.; Huang, X.; Kornberg, R.D.
Deposited on : 2011-05-16
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)
RNA backbone	3983	1222 (3.50-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1733	 2% 57% 20% 19%
2	B	1224	 3% 63% 24% 9%
3	C	318	 59% 20% 16%
4	E	215	 2% 79% 20%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	155	 53% 45%
6	H	146	 5% 65% 20% 5% 9%
7	I	122	 3% 80% 16%
8	J	70	 % 57% 26% 10% 7%
9	K	120	 % 59% 33% 5%
10	L	70	 9% 39% 17% 9% 34%
11	R	5	 40% 60%
12	T	29	 3% 28% 17% 55%

2 Entry composition [i](#)

There are 14 unique types of molecules in this entry. The entry contains 28672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1405	11043	6965	1936	2081	61	0	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	1114	8861	5610	1549	1647	55	0	0	0

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	266	2095	1317	348	417	13	0	0	0

- Molecule 4 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	E	214	1752	1111	309	321	11	0	0	0

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	F	85	688	439	116	130	3	0	0	0

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	H	133	1068	673	180	211	4	0	0	0

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	I	119	971	596	179	186	10	0	0	0

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	J	65	532	339	93	94	6	0	0	0

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	K	114	919	590	156	171	2	0	0	0

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	L	46	363	224	72	63	4	0	0	0

- Molecule 11 is a RNA chain called RNA (5'-R(*AP*GP*GP*(5BU)P*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	Br	C	N	O	P			
11	R	5	109	1	49	22	33	4	0	0	0

- Molecule 12 is a DNA chain called DNA (5'-D(*CP*TP*AP*CP*CP*GP*AP*TP*AP*AP*GP*CP*AP*GP*AP*CP*GP*AP*TP*CP*AP*CP*CP*TP*CP*GP*AP*TP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
12	T	13	262	125	46	78	13	0	0	0

- Molecule 13 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	2	Total	Zn	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	B	1	Total 1	Zn 1	0	0
13	C	1	Total 1	Zn 1	0	0
13	I	2	Total 2	Zn 2	0	0
13	J	1	Total 1	Zn 1	0	0
13	L	1	Total 1	Zn 1	0	0


- Molecule 14 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	A	1	Total 1	Mg 1	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-directed RNA polymerase II subunit RPB1








• Molecule 3: DNA-directed RNA polymerase II subunit RPB3




• Molecule 4: DNA-directed RNA polymerases I, II, and III subunit RPABC1





• Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC2






- Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC3




- Molecule 7: DNA-directed RNA polymerase II subunit RPB9




- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC5



- Molecule 9: DNA-directed RNA polymerase II subunit RPB11



- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC4





- Molecule 11: RNA (5'-R(*AP*GP*GP*(5BU)P*G)-3')





- Molecule 12: DNA (5'-D(*CP*TP*AP*CP*CP*GP*AP*TP*AP*AP*GP*CP*AP*GP*AP*C
P*GP*AP*TP*CP*AP*CP*CP*TP*CP*GP*AP*TP*G)-3')



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	158.29Å 220.77Å 192.24Å 90.00° 97.69° 90.00°	Depositor
Resolution (Å)	47.76 – 3.20 47.76 – 3.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (47.76-3.20) 99.0 (47.76-3.20)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 3.19Å)	Xtrriage
Refinement program	BUSTER-TNT BUSTER 2.8.0, BUSTER 2.8.0	Depositor
R, R_{free}	0.183 , 0.231 0.202 , 0.247	Depositor DCC
R_{free} test set	5360 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	67.2	Xtrriage
Anisotropy	0.685	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 92.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	28672	wwPDB-VP
Average B, all atoms (Å ²)	94.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 5BU, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.80	2/11241 (0.0%)	1.44	61/15199 (0.4%)
2	B	0.81	1/9033 (0.0%)	1.42	61/12181 (0.5%)
3	C	0.76	0/2133	1.43	17/2891 (0.6%)
4	E	0.77	0/1788	1.40	9/2406 (0.4%)
5	F	0.77	0/700	1.33	0/945
6	H	0.74	0/1086	1.29	14/1470 (1.0%)
7	I	0.75	0/989	1.31	5/1331 (0.4%)
8	J	0.83	0/541	1.55	6/727 (0.8%)
9	K	0.71	0/937	1.34	6/1265 (0.5%)
10	L	0.86	0/365	1.43	9/485 (1.9%)
11	R	0.58	0/99	1.32	1/154 (0.6%)
12	T	0.43	0/292	1.14	1/447 (0.2%)
All	All	0.79	3/29204 (0.0%)	1.41	190/39501 (0.5%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	487	MET	SD-CE	-5.96	1.64	1.79
2	B	563	MET	SD-CE	-5.91	1.64	1.79
1	A	249	SER	CA-C	5.67	1.55	1.52

The worst 5 of 190 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	647	GLY	CA-C-N	10.56	140.72	121.70
2	B	647	GLY	C-N-CA	10.56	140.72	121.70
3	C	172	PRO	CA-C-N	9.02	138.77	121.54
3	C	172	PRO	C-N-CA	9.02	138.77	121.54
1	A	1097	GLY	CA-C-N	8.63	125.68	120.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11043	0	11133	181	0
2	B	8861	0	8884	152	0
3	C	2095	0	2051	39	0
4	E	1752	0	1776	17	0
5	F	688	0	707	1	0
6	H	1068	0	1040	11	0
7	I	971	0	927	8	0
8	J	532	0	542	18	0
9	K	919	0	929	19	0
10	L	363	0	386	5	0
11	R	109	0	54	1	0
12	T	262	0	147	3	0
13	A	2	0	0	0	0
13	B	1	0	0	0	0
13	C	1	0	0	0	0
13	I	2	0	0	0	0
13	J	1	0	0	0	0
13	L	1	0	0	0	0
14	A	1	0	0	0	0
All	All	28672	0	28576	397	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 397 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1123:GLY:HA3	1:A:1124:HIS:HB2	1.59	0.83
1:A:869:GLY:O	4:E:204:THR:HG21	1.79	0.82
8:J:48:ARG:O	8:J:52:THR:HB	1.80	0.82
2:B:1013:ASN:HD21	2:B:1015:HIS:HD2	1.28	0.81
2:B:783:THR:HG22	8:J:63:TYR:HE1	1.45	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1395/1733 (80%)	1234 (88%)	124 (9%)	37 (3%)	4	25
2	B	1096/1224 (90%)	962 (88%)	96 (9%)	38 (4%)	3	20
3	C	264/318 (83%)	239 (90%)	21 (8%)	4 (2%)	8	37
4	E	212/215 (99%)	203 (96%)	7 (3%)	2 (1%)	14	47
5	F	83/155 (54%)	75 (90%)	7 (8%)	1 (1%)	10	42
6	H	129/146 (88%)	103 (80%)	16 (12%)	10 (8%)	1	5
7	I	117/122 (96%)	105 (90%)	11 (9%)	1 (1%)	14	47
8	J	63/70 (90%)	58 (92%)	2 (3%)	3 (5%)	2	14
9	K	112/120 (93%)	107 (96%)	4 (4%)	1 (1%)	14	47
10	L	44/70 (63%)	30 (68%)	7 (16%)	7 (16%)	0	0
All	All	3515/4173 (84%)	3116 (89%)	295 (8%)	104 (3%)	3	23

5 of 104 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	40	THR
1	A	54	ASN
1	A	55	ASP
1	A	250	ILE
1	A	399	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1225/1520 (81%)	1058 (86%)	167 (14%)	3	18
2	B	967/1061 (91%)	851 (88%)	116 (12%)	5	23
3	C	234/274 (85%)	207 (88%)	27 (12%)	5	24
4	E	196/197 (100%)	183 (93%)	13 (7%)	15	47
5	F	75/137 (55%)	74 (99%)	1 (1%)	61	78
6	H	117/128 (91%)	100 (86%)	17 (14%)	3	16
7	I	113/116 (97%)	103 (91%)	10 (9%)	9	35
8	J	60/65 (92%)	52 (87%)	8 (13%)	4	19
9	K	99/102 (97%)	85 (86%)	14 (14%)	3	17
10	L	40/57 (70%)	31 (78%)	9 (22%)	1	5
All	All	3126/3657 (86%)	2744 (88%)	382 (12%)	5	22

5 of 382 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	762	ASN
3	C	21	ILE
2	B	844	SER
2	B	1019	SER
3	C	144	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 77 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	1015	HIS
6	H	35	GLN
2	B	1176	ASN
3	C	167	HIS
9	K	110	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
11	R	5/5 (100%)	4 (80%)	1 (20%)

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
11	R	7	G
11	R	8	G
11	R	9	5BU
11	R	10	G

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
11	R	6	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
11	5BU	R	9	11,12	19,22,23	1.10	2 (10%)	27,32,35	1.30	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	5BU	R	9	11,12	-	2/7/25/26	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	R	9	5BU	C2-N1	2.64	1.42	1.38
11	R	9	5BU	C4-C5	2.44	1.49	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	R	9	5BU	O4-C4-C5	-2.46	122.66	125.80
11	R	9	5BU	O3'-C3'-C4'	-2.43	104.10	111.08
11	R	9	5BU	O4'-C4'-C3'	-2.29	100.62	105.15
11	R	9	5BU	C3'-C2'-C1'	-2.09	97.50	101.46
11	R	9	5BU	C2'-C3'-C4'	-2.04	98.66	102.61

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	R	9	5BU	C3'-C4'-C5'-O5'
11	R	9	5BU	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1405/1733 (81%)	0.15	40 (2%) 55 35	42, 85, 175, 219	0
2	B	1114/1224 (91%)	0.11	37 (3%) 49 30	37, 78, 147, 189	0
3	C	266/318 (83%)	-0.11	0 100 100	49, 75, 114, 155	0
4	E	214/215 (99%)	0.27	4 (1%) 66 46	61, 115, 162, 175	0
5	F	85/155 (54%)	-0.10	0 100 100	66, 96, 133, 152	0
6	H	133/146 (91%)	0.45	8 (6%) 27 17	80, 119, 154, 166	0
7	I	119/122 (97%)	0.19	4 (3%) 48 30	55, 93, 129, 142	0
8	J	65/70 (92%)	-0.10	1 (1%) 72 52	49, 67, 102, 116	0
9	K	114/120 (95%)	-0.14	1 (0%) 81 64	51, 77, 104, 120	0
10	L	46/70 (65%)	0.85	6 (13%) 7 6	61, 116, 143, 158	0
11	R	4/5 (80%)	0.03	0 100 100	78, 91, 109, 113	0
12	T	13/29 (44%)	0.80	1 (7%) 19 13	84, 115, 174, 181	0
All	All	3578/4207 (85%)	0.13	102 (2%) 53 35	37, 84, 162, 219	0

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	883	LEU	5.9
1	A	1081	LEU	5.7
6	H	85	GLY	5.5
2	B	715	ALA	5.1
1	A	1087	ALA	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
11	5BU	R	9	21/22	0.91	0.09	77,80,90,102	1

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
13	ZN	A	1734	1/1	0.82	0.09	300,300,300,300	0
13	ZN	A	1735	1/1	0.95	0.07	135,135,135,135	0
13	ZN	B	1307	1/1	0.99	0.03	149,149,149,149	0
13	ZN	C	319	1/1	1.00	0.03	67,67,67,67	0
13	ZN	I	203	1/1	1.00	0.03	87,87,87,87	0
13	ZN	I	204	1/1	1.00	0.04	66,66,66,66	0
13	ZN	J	101	1/1	1.00	0.03	58,58,58,58	0
13	ZN	L	105	1/1	1.00	0.02	102,102,102,102	0
14	MG	A	2001	1/1	1.00	0.04	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.