



Full wwPDB EM Validation Report ⓘ

Mar 27, 2026 – 09:32 PM UTC

PDB ID : 8S2T / pdb_00008s2t
EMDB ID : EMD-19683
Title : Cryo-EM structure of cell-free synthesized Human beta-1 adrenergic receptor cotranslationally inserted into a lipidic nanodiscs
Authors : Merino, F.; Koeck, Z.; Ermel, U.; Dahlhaus, P.; Doetsch, V.; Kubicek, J.; Frangakis, A.S.; Hilger, D.; Bernhard, F.
Deposited on : 2024-02-19
Resolution : 3.30 Å (reported)
Based on initial models : ., 6E3Y

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15087 (2.80 - 3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	
2	B	345	
3	C	68	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	N	134	
5	R	477	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7955 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoform Gnas-2 of Guanine nucleotide-binding protein G(s) subunit alpha isoforms short.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	196	Total	C	N	O	S	0	0
			1673	1062	307	299	5		

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	338	Total	C	N	O	S	0	0
			2601	1604	467	509	21		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP P62873
B	-3	PRO	-	expression tag	UNP P62873
B	-2	GLY	-	expression tag	UNP P62873
B	-1	SER	-	expression tag	UNP P62873
B	0	SER	-	expression tag	UNP P62873
B	1	GLY	-	expression tag	UNP P62873

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	57	Total	C	N	O	S	0	0
			436	273	77	83	3		

- Molecule 4 is a protein called Nanobody-35.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	N	126	Total	C	N	O	S	0	0
			961	599	168	188	6		


- Molecule 5 is a protein called Beta-1 adrenergic receptor.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	R	285	Total	C	N	O	S	0	0
			2269	1496	380	375	18		

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	85	SER	LYS	conflict	UNP P08588
R	107	VAL	MET	conflict	UNP P08588
R	146	VAL	ILE	conflict	UNP P08588
R	147	TRP	GLU	conflict	UNP P08588
R	316	LEU	ALA	conflict	UNP P08588
R	361	ALA	PHE	conflict	UNP P08588
R	372	MET	PHE	conflict	UNP P08588
R	392	ALA	CYS	conflict	UNP P08588
R	393	ALA	CYS	conflict	UNP P08588

- Molecule 6 is ISOPRENALINE (CCD ID: 5FW) (formula: $C_{11}H_{17}NO_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
6	R	1	Total	C	N	O	0
			15	11	1	3	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Isoform Gnas-2 of Guanine nucleotide-binding protein G(s) subunit alpha isoforms short




- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1



- Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2




- Molecule 4: Nanobody-35





• Molecule 5: Beta-1 adrenergic receptor



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	74266	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	65	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	165000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.073	Depositor
Minimum map value	-0.032	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.012	Depositor
Map size (Å)	339.906, 339.906, 339.906	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.13302, 1.13302, 1.13302	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5FW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.28	5/1706 (0.3%)	1.27	13/2290 (0.6%)
2	B	1.44	6/2648 (0.2%)	1.33	15/3589 (0.4%)
3	C	1.17	1/442 (0.2%)	1.29	9/597 (1.5%)
4	N	1.35	5/981 (0.5%)	1.34	4/1329 (0.3%)
5	R	1.21	2/2327 (0.1%)	1.32	20/3170 (0.6%)
All	All	1.32	19/8104 (0.2%)	1.31	61/10975 (0.6%)

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	225	HIS	CB-CG	-9.03	1.37	1.50
3	C	62	ARG	NE-CZ	6.75	1.40	1.33
1	A	218	ARG	CB-CG	-6.11	1.34	1.52
2	B	205	ASP	CB-CG	-6.02	1.36	1.52
2	B	142	HIS	CB-CG	-5.97	1.41	1.50
2	B	211	TRP	NE1-CE2	-5.93	1.30	1.37
5	R	138	ASP	CB-CG	-5.79	1.37	1.52
2	B	61	MET	CG-SD	-5.77	1.66	1.80
2	B	190	LEU	CB-CG	-5.68	1.42	1.53
4	N	98	ARG	CG-CD	-5.62	1.35	1.52
4	N	72	ARG	CD-NE	-5.51	1.38	1.46
4	N	108	PHE	CB-CG	-5.48	1.38	1.50
1	A	208	PHE	CB-CG	-5.39	1.38	1.50
1	A	218	ARG	CG-CD	-5.37	1.36	1.52
4	N	98	ARG	CZ-NH2	-5.26	1.26	1.33
1	A	41	HIS	CB-CG	-5.18	1.42	1.50
5	R	243	VAL	CA-C	-5.18	1.46	1.52
1	A	193	ILE	CB-CG1	5.16	1.63	1.53
4	N	38	ARG	CB-CG	-5.13	1.37	1.52

All (61) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	197	ARG	N-CA-C	-10.50	101.02	114.04
5	R	162	SER	CA-C-N	8.59	128.23	119.56
5	R	162	SER	C-N-CA	8.59	128.23	119.56
4	N	110	VAL	N-CA-C	-8.20	99.00	109.58
3	C	48	ASP	CA-C-N	7.99	127.71	119.56
3	C	48	ASP	C-N-CA	7.99	127.71	119.56
1	A	206	HIS	CA-CB-CG	7.86	121.66	113.80
2	B	161	SER	N-CA-C	7.28	119.98	109.14
5	R	212	ASP	CA-C-N	7.24	126.94	119.56
5	R	212	ASP	C-N-CA	7.24	126.94	119.56
5	R	198	TRP	N-CA-C	-7.00	104.21	112.89
5	R	196	MET	N-CA-C	-6.93	104.82	113.28
1	A	208	PHE	CA-CB-CG	-6.83	106.97	113.80
1	A	215	ASP	CA-CB-CG	6.67	119.28	112.60
1	A	36	VAL	N-CA-C	-6.62	105.38	111.67
2	B	237	ASN	N-CA-C	-6.32	104.81	114.16
2	B	38	ASP	CA-C-N	6.29	126.26	120.03
2	B	38	ASP	C-N-CA	6.29	126.26	120.03
2	B	235	PHE	N-CA-C	-6.27	101.87	109.83
5	R	373	ASN	CA-C-N	6.20	125.88	119.56
5	R	373	ASN	C-N-CA	6.20	125.88	119.56
5	R	86	THR	CA-C-N	6.20	126.66	119.47
5	R	86	THR	C-N-CA	6.20	126.66	119.47
5	R	189	VAL	N-CA-C	-6.17	106.76	112.43
5	R	132	GLU	N-CA-C	-6.15	104.13	111.69
5	R	354	VAL	CA-C-N	6.03	125.99	119.78
5	R	354	VAL	C-N-CA	6.03	125.99	119.78
1	A	317	ASP	CA-C-N	5.93	126.35	119.47
1	A	317	ASP	C-N-CA	5.93	126.35	119.47
3	C	54	VAL	CA-C-N	5.91	125.87	119.78
3	C	54	VAL	C-N-CA	5.91	125.87	119.78
3	C	59	ASN	CA-C-N	5.90	126.01	119.87
3	C	59	ASN	C-N-CA	5.90	126.01	119.87
5	R	380	SER	N-CA-C	-5.84	100.42	109.87
3	C	52	THR	CA-C-N	5.63	125.63	119.89
3	C	52	THR	C-N-CA	5.63	125.63	119.89
2	B	290	ASP	CA-CB-CG	5.50	118.10	112.60
2	B	289	TYR	N-CA-C	5.47	117.73	109.41
4	N	99	CYS	CA-C-N	5.42	126.62	119.84
4	N	99	CYS	C-N-CA	5.42	126.62	119.84
3	C	54	VAL	N-CA-CB	-5.35	108.14	111.83
1	A	205	PHE	CA-CB-CG	5.34	119.14	113.80
2	B	151	PHE	CA-CB-CG	5.31	119.11	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	298	PHE	CA-C-N	5.29	125.60	119.47
1	A	298	PHE	C-N-CA	5.29	125.60	119.47
2	B	199	PHE	CA-CB-CG	5.29	119.09	113.80
5	R	76	ASN	CA-CB-CG	-5.26	107.34	112.60
1	A	11	ASP	CA-CB-CG	5.23	117.83	112.60
2	B	189	SER	N-CA-C	5.21	117.66	108.75
5	R	196	MET	CA-C-N	5.16	129.52	122.34
5	R	196	MET	C-N-CA	5.16	129.52	122.34
1	A	225	ASN	CA-CB-CG	5.14	117.74	112.60
2	B	78	LYS	N-CA-C	5.13	117.60	109.50
5	R	130	PHE	CA-CB-CG	-5.12	108.68	113.80
4	N	109	ASP	CA-CB-CG	-5.08	107.52	112.60
1	A	218	ARG	CB-CA-C	-5.07	102.37	110.79
5	R	95	LEU	N-CA-C	-5.07	105.83	111.36
2	B	256	ARG	N-CA-C	-5.06	106.28	112.90
1	A	218	ARG	N-CA-C	5.05	116.78	111.28
2	B	316	SER	CA-C-N	-5.01	115.42	122.63
2	B	316	SER	C-N-CA	-5.01	115.42	122.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1673	0	1649	4	0
2	B	2601	0	2505	6	0
3	C	436	0	448	0	0
4	N	961	0	928	5	0
5	R	2269	0	2326	7	0
6	R	15	0	16	0	0
All	All	7955	0	7872	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:ASP:OD2	2:B:78:LYS:NZ	2.37	0.58
5:R:356:ASP:N	5:R:356:ASP:OD1	2.33	0.57
1:A:214:ARG:NH2	4:N:114:THR:OG1	2.40	0.55
5:R:190:SER:OG	5:R:191:PHE:N	2.41	0.54
5:R:204:ASP:N	5:R:204:ASP:OD1	2.38	0.53
2:B:163:ASP:N	2:B:163:ASP:OD1	2.45	0.46
4:N:104:THR:OG1	4:N:105:ARG:N	2.48	0.46
1:A:218:ARG:HG3	4:N:109:ASP:OD2	2.16	0.45
4:N:50:ASP:OD1	4:N:50:ASP:C	2.61	0.44
5:R:373:ASN:N	5:R:374:PRO:CD	2.82	0.43
5:R:112:VAL:N	5:R:113:PRO:HD2	2.34	0.42
2:B:316:SER:O	2:B:317:CYS:C	2.62	0.42
2:B:155:ASN:O	2:B:155:ASN:CG	2.61	0.42
1:A:213:GLN:O	1:A:214:ARG:HB2	2.19	0.42
2:B:102:THR:OG1	2:B:103:CYS:N	2.51	0.42
2:B:76:ASP:N	2:B:76:ASP:OD1	2.40	0.41
4:N:47:TRP:O	4:N:61:THR:OG1	2.39	0.41
5:R:221:ASN:OD1	5:R:221:ASN:C	2.63	0.41
5:R:235:VAL:HB	5:R:236:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/380 (49%)	183 (98%)	3 (2%)	0	100	100
2	B	336/345 (97%)	327 (97%)	9 (3%)	0	100	100
3	C	55/68 (81%)	55 (100%)	0	0	100	100
4	N	124/134 (92%)	119 (96%)	5 (4%)	0	100	100
5	R	281/477 (59%)	273 (97%)	8 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	982/1404 (70%)	957 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/341 (53%)	180 (100%)	0	100	100
2	B	281/285 (99%)	281 (100%)	0	100	100
3	C	46/56 (82%)	46 (100%)	0	100	100
4	N	104/112 (93%)	104 (100%)	0	100	100
5	R	243/378 (64%)	243 (100%)	0	100	100
All	All	854/1172 (73%)	854 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	GLN
1	A	225	ASN
1	A	253	GLN
2	B	91	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	5FW	R	501	-	15,15,15	1.80	6 (40%)	19,20,20	0.88	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	5FW	R	501	-	-	1/9/9/9	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	R	501	5FW	CAK-CAH	3.44	1.44	1.39
6	R	501	5FW	CAM-CAO	3.09	1.45	1.40
6	R	501	5FW	CAK-CAM	2.44	1.42	1.38
6	R	501	5FW	CAL-CAH	2.37	1.42	1.39
6	R	501	5FW	CAH-CAE	2.22	1.55	1.51
6	R	501	5FW	CAN-CAL	2.18	1.42	1.38

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	R	501	5FW	CAF-NAD-CAG	2.27	121.34	114.38

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	R	501	5FW	OAA-CAE-CAF-NAD

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-19683. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 149




Y Index: 156




Z Index: 190


6.3.2 Raw map



X Index: 149



Y Index: 155




Z Index: 190


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.012. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_19683_msk_1.map [i](#)



X




Y




Z


6.6.2 emd_19683_msk_2.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 46 nm³; this corresponds to an approximate mass of 41 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.303 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.303 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.30	-	-
Author-provided FSC curve	3.33	3.79	3.37
Unmasked-calculated*	3.82	7.42	3.88

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.82 differs from the reported value 3.3 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-19683 and PDB model 8S2T. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.012 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.012).

9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 71% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.012) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7130	0.4470
A	0.7030	0.4500
B	0.7620	0.4820
C	0.6920	0.4220
N	0.7720	0.4720
R	0.6420	0.4010

